

EQUATE NAPROXEN SODIUM- naproxen sodium tablet, film coated
Wal-Mart Stores Inc

Wal-Mart Naproxen Sodium Tablets, 200 mg Drug Facts

Active ingredient (in each caplet)

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen,

or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - you have difficulty swallowing
 - it feels like the pill is stuck in your throat
 - redness or swelling is present in the painful area
 - any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older	<ul style="list-style-type: none">• take 1 caplet every 8 to 12 hours while symptoms last• for the first dose you may take 2 caplets within the first hour• do not exceed 2 caplets in any 8- to 12-hour period• do not exceed 3 caplets in a 24-hour period
Children under 12 years	<ul style="list-style-type: none">• ask a doctor

Other information

- each caplet contains: sodium 21 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients

FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments?

1-888-287-1915

Principal Display Panel

EASY OPEN CAP

Compare to Aleve® Caplets Active Ingredient

Naproxen Sodium Tablets, 220 mg

Pain Reliever/Fever Reducer (NSAID)

Strength to Last 12 Hours

PACKAGE NOT CHILD RESISTANT

220 mg

225 CAPLETS†

†Capsule Shaped Tablets

Actual Size

Drug Facts (continued)
Inactive Ingredients: FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, titanium dioxide
Questions or comments?
1-888-281-1915
Gluten-Free

Satisfaction guaranteed - For quantities or comments please call 1-888-281-1915

Contains: Wheat, Soy, Yeast, Lactose, Egg, Milk, Sesame, Peanuts, Tree Nuts, Fish, Shellfish, and Milk Protein
Manufactured by: Watson, LLC, a Division of Watson Pharmaceuticals, Inc.
K452

EASY OPEN CAP

NDC 49035-140-83

Compare to
Aleve®
Capsule
Active
Ingredients™

equate™

Naproxen Sodium

Tablets, 220 mg
Pain Reliever/Fever Reducer (NSAID)
Strength to Last 12 Hours

PACKAGE NOT CHILD RESISTANT

220
mg

225
CAPSULE

[†]Capsule
Shaped
Tablet

Actual Size

DO NOT USE IF PRINTED FOIL LINER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each caplet): Naproxen sodium 220 mg (naproxen 200 mg (NSAID))...over reducer

Purpose: Pain reliever/fever reducer
Temporarily reduces fever

Uses: Temporarily relieves minor aches and pains due to:

- minor pain of arthritis
- muscle aches
- headache
- backache
- menstrual cramps
- the common cold
- temporarily reduces fever

Warnings

Alert: Naproxen sodium may cause a severe allergic reaction in some people. Stop use and seek medical help if an allergic reaction occurs, stop use and seek medical help if an allergic reaction occurs, stop use and seek medical help if an allergic reaction occurs, stop use and seek medical help if an allergic reaction occurs.

Stop use and ask a doctor if: you have had stomach ulcers or bleeding problems, you have had stomach ulcers or bleeding problems, you have had stomach ulcers or bleeding problems, you have had stomach ulcers or bleeding problems.

Do not use: if you have ever had an allergic reaction to any other pain reliever/fever reducer.

Ask a doctor before use if you are: taking any other medicine, you are pregnant, you are breastfeeding, you are taking any other medicine, you are pregnant, you are breastfeeding, you are taking any other medicine, you are pregnant, you are breastfeeding.

When using this product: Take with food or milk. If stomach upset occurs, the risk of heart attack or stroke may increase if you use more than directed or for longer than directed.

Stop use and ask a doctor if: you experience any of the following signs of stomach bleeding: ■ feel faint, ■ vomit blood, ■ have bloody or black stools.

Keep this and all other medicines out of the reach of children.

Keep this and all other medicines out of the reach of children.

Keep this and all other medicines out of the reach of children.

Keep this and all other medicines out of the reach of children.

Keep this and all other medicines out of the reach of children.

Keep this and all other medicines out of the reach of children.

Keep this and all other medicines out of the reach of children.

Keep this and all other medicines out of the reach of children.

Keep this and all other medicines out of the reach of children.

Keep this and all other medicines out of the reach of children.

Keep this and all other medicines out of the reach of children.

Keep this and all other medicines out of the reach of children.

Keep this and all other medicines out of the reach of children.



SEEK BACK HERE

Drug Facts (continued)

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If **pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

(1-800-222-1222)

- Directions**
- do not take more than directed
 - the smallest effective dose should be used
 - drink a full glass of water with each dose

Adults and children 12 years and older

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12-hour period
- do not exceed 3 caplets in a 24-hour period

Children under 12 years

- ask a doctor

Other information

- each caplet contains: sodium 21 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).



EQUATE NAPROXEN SODIUM

naproxen sodium tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49035-140
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	200 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
POVIDONES (UNII: FZ989GH94E)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	BLUE (light blue)	Score	no score
Shape	CAPSULE	Size	12mm
Flavor		Imprint Code	L368
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49035-140-83	225 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074661	08/09/2013	

Labeler - Wal-Mart Stores Inc (051957769)