

**XTRA-7 (PART B)- chlorine dioxide teat dip solution**  
**AgroChem Inc**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click [here](#).*

-----

**XTRA-7 (Part B)**

**XTRA-7 (PART B)**

**XTRA-7 (PART B)**

**TEAT DIP**

**WARNING**

**AVOID FREEZING.**

**KEEP OUT OF DIRECT SUNLIGHT**

**CAUSES SERIOUS EYE IRRITATION.**

**KEEP OUT OF REACH OF CHILDREN.**

**FOR FARM AND INDUSTRIAL USE ONLY**

.

**PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.**

**FIRST AID / RESPONSE:**

**IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.**

**STORAGE: Store locked up.**

**DISPOSAL: Dispose of contents/ container to an approved waste disposal plant**

**EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494**

**ACTIVE INGREDIENT:**

**Sodium Chlorite - 0.65 %**

**XTRA-7 is a two-part Chlorine Dioxide teat dip that can be effectively used to reduce the spread of mastitis causing organisms. Recommended for pre and post milking applications.**

**Directions:**

**NOTE: MUST BE MIXED PRIOR TO USE. MIX PART A & PART B in equal parts.**

**CREATES CHLORINE DIOXIDE. Reaches minimum Chlorine Dioxide levels within 15 minutes of mixing. Will maintain minimum Chlorine Dioxide levels for up to 24 hours. It is recommended to replace with freshly mixed XTRA-7 after 24 hours.**

**Directions:**

**General Use**

**Use a teat dip cup to apply. Clean dip cup just prior to use.**

**If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.**

**If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.**

**If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.**

**Teat Sprays can also be foamed or sprayed using recommended equipment.**

#### **Pre Dipping Use**

**Perform forestripping and inspect teats for clinical mastitis.**

**Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels**

**Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product.**

**Allow product to stand for 30 seconds.**

**Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.**

#### **Post Dipping Use**

**Immediately after each milking, use product.**

**Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe.**

**If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.**

#### **Extended Use**

**When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking.**

**Dipping should be resumed at least twice daily one week before calving.**

# XTRA-7 (PART B)

## CHLORINE DIOXIDE TEAT DIP & SPRAY

TEAT DIP

### WARNING

**AVOID FREEZING.**  
**KEEP OUT OF DIRECT SUNLIGHT.**  
**CAUSES SERIOUS EYE IRRITATION.**  
**KEEP OUT OF REACH OF CHILDREN.**  
**FOR FARM AND INDUSTRIAL USE ONLY.**



**PREVENTION:** Wash skin thoroughly after handling. Wear eye protection / face protection.  
**FIRST AID / RESPONSE:**  
**IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice / attention.  
**STORAGE:** Store locked up.  
**DISPOSAL:** Dispose of contents / container to an approved waste disposal plant

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494

**ACTIVE INGREDIENT:**  
 Sodium Chlorite - 0.65%

**NET CONTENTS:** 55 GALLON



3653S

XTRA-7 is a two-part Chlorine Dioxide teat dip and can be effectively used to reduce the spread of mastitis causing organisms. Recommended for pre and post milking applications.

**Directions:**

**NOTE: MUST BE MIXED PRIOR TO USE. MIX PART A & PART B in equal parts.**  
**CREATES CHLORINE DIOXIDE.** Reaches minimum Chlorine Dioxide levels within 15 minutes of mixing. Will maintain minimum Chlorine Dioxide levels for up to 24 hours. It is recommended to replace with freshly mixed XTRA-7 after 24 hours.

**General Use**

Use a teat dip cup to apply. Clean dip cup just prior to use.  
 If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.  
 If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.  
 If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.  
 Can also be foamed or sprayed using recommended equipment.

**Pre Dipping Use**

Perform fores tripping and inspect teats for clinical mastitis.  
 Pre-wash teats thoroughly just prior to milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels.  
 Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product.  
 Allow product to stand for 30 seconds.  
 Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

**Post Dipping Use**

Immediately after each milking, use product.  
 Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe.  
 If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

**Extended Use**

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking.  
 Dipping should be resumed at least twice daily one week before calving.



**AgroChem**

26 Freedom Way • Saratoga Springs, NY 12866 • (518) 226-4850

## XTRA-7 (PART B)

chlorine dioxide teat dip solution

### Product Information

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:17307-3650
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM CHLORITE (UNII: G538EBV4VF) (CHLORITE ION - UNII: Z63H374SB6)	SODIUM CHLORITE	0.103 kg in 1 kg

### Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17307-3650-1	19 kg in 1 CONTAINER		
2	NDC:17307-3650-2	57 kg in 1 DRUM		
3	NDC:17307-3650-3	208 kg in 1 DRUM		
4	NDC:17307-3650-4	1040 kg in 1 CONTAINER		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		06/14/2018	

**Labeler** - AgroChem Inc (805427700)**Registrant** - AgroChem Inc (805427700)**Establishment**

Name	Address	ID/FEI	Business Operations
AgroChem Inc		805427700	api manufacture, manufacture

Revised: 6/2018

AgroChem Inc