METHYLPHENIDATE HYDROCHLORIDE- methylphenidate hydrochloride tablet, extended release AvPAK

HIGHLIGHTS OF PRESCRIBING INFORMATION Methylphenidate Hydrochloride Extended-Release Tablets, USP CII Rx Only

These highlights do not include all the information needed to use METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE TABLETS safely and effectively. See full prescribing information for METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE TABLETS.

Initial U.S. Approval: 2000

WARNING: DRUG DEPENDENCE

See full prescribing information for complete boxed warning.

Methylphenidate hydrochloride extended-release tablets should be given cautiously to patients with a history of drug dependence or alcoholism. Chronic abusive use can lead to marked tolerance and psychological dependence, with varying degrees of abnormal behavior.

Methylphenidate hydrochloride extended-release tablets is a CNS stimulant indicated for the treatment of Attention Deficit Hyperactivity Disorder (ADHD) in children 6 years of age and older, adolescents, and adults up to the age of 65. (1)

------ DOSAGE AND ADMINISTRATION

- Methylphenidate hydrochloride extended-release tablets should be taken once daily in the morning and swallowed whole with the aid of liquids. Methylphenidate hydrochloride extended-release tablets should not be chewed or crushed. Methylphenidate hydrochloride extended-release tablets may be taken with or without food. (2.1)
- For children and adolescents new to methylphenidate, the recommended starting dosage is 18 mg once daily. Dosage may be increased by 18 mg/day at weekly intervals and should not exceed 54 mg/day in children and 72 mg/day in adolescents. (2.2)
- For adult patients new to methylphenidate, the recommended starting dose is 18 or 36 mg/day. Dosage may be increased by 18 mg/day at weekly intervals and should not exceed 72 mg/day for adults. (2.2)
- For patients currently using methylphenidate, dosing is based on current dose regimen and clinical judgment. (2.3)

DOSAGE FORMS AND STRENGTHS
Tablets: 18, 27, 36, and 54 mg (3)
CONTRAINDICATIONS
 Known hypersensitivity to the product (4.1)
Marked anxiety, tension, or agitation (4.2)
• Glaucoma (4.3)
 Tics or a family history or diagnosis of Tourette's syndrome (4.4)
 Do not use methylphenidate hydrochloride extended-release tablet s in patients currently using or within 2 weeks of using an MAO inhibitor (4.5)
WARNINGS AND RECAUTIONS

• Serious Cardiovascular Events: Sudden death has been reported in association with CNS stimulant treatment at usual doses in children and adolescents with structural cardiac abnormalities or other serious heart problems. Sudden death, stroke, and myocardial infarction have been reported in adults taking stimulant drugs at usual doses for ADHD. Stimulant products generally should not be used in patients with known structural cardiac abnormalities, cardiomyopathy, serious heart rhythm abnormalities, coronary artery disease, or other serious heart problems. (5.1)

- Increase in Blood Pressure: Monitor patients for changes in heart rate and blood pressure and use with caution in patients for whom an increase in blood pressure or heart rate would be problematic. (5.1)
- Psychiatric Adverse Events: Use of stimulants may cause treatment-emergent psychotic or manic symptoms in patients with no prior history, or exacerbation of symptoms in patients with preexisting psychiatric illness. Clinical evaluation for Bipolar Disorder is recommended prior to stimulant use. Monitor for aggressive behavior. (5.2)
- Seizures: Stimulants may lower the convulsive threshold. Discontinue in the presence of seizures. (5.3)
- Priapism: cases of painful and prolonged penile erections and priapism have been reported with methylphenidate products. Immediate medical attention should be sought if signs or symptoms of painful or prolonged penile erections or priapism are observed. (5.4)
- Peripheral Vasculopathy, including Raynaud's Phenomenon: Stimulants used to treat ADHD are associated with peripheral vasculopathy, including Raynaud's phenomenon. Careful observation for digital changes is necessary during treatment with ADHD stimulants. (5.5)
- Visual Disturbance: difficulties with accommodation and blurring of vision have been reported with stimulant treatment. (5.7)
- Long-Term Suppression of Growth: monitor height and weight at appropriate intervals in pediatric patients. (5.6)
- Gastrointestinal obstruction with preexisting GI narrowing. (5.8)
- Hematologic monitoring: Periodic CBC, differential, and platelet counts are advised during prolonged therapy. (5.9)

ADVERSE REACTIONS The most common adverse reaction in double-blind clinical trials (>5%) in children and adolescents was abdominal pain upper. The most common adverse reactions in double-blind clinical trials (>5%) in adult patients were decreased appetite, headache, dry mouth, nausea, insomnia, anxiety, dizziness, weight decreased, irritability, and hyperhidrosis. (6.1 and 6.2)

The most common adverse reactions associated with discontinuation ($\geq 1\%$) from either pediatric or adult clinical trials were anxiety, irritability, insomnia, and blood pressure increased. (6.3)

To report SUSPECTED ADVERSE REACTIONS, contact AvKARE at 1-855-361-3993 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

- Do not use methylphenidate hydrochloride extended-release tablets in patients currently using or within
- Do not use methylphenidate hydrochloride extended-release tablets in patients currently using or within 2 weeks of using an MAO inhibitor (7.1)
- Methylphenidate hydrochloride extended-release tablets may increase blood pressure; use cautiously with vasopressors (7.2)
- Inhibition of metabolism of coumarin anticoagulants, anticonvulsants, and some antidepressants (7.3)
- USE IN SPECIFIC POPULATIONS
- Caution should be exercised if administered to nursing mothers (8.3)
- Safety and efficacy has not been established in children less than six years old or elderly patients greater than 65 years of age (8.4 and 8.5)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 4/2021

FULL PRESCRIBING INFORMATION: CONTENTS* WARNING: DRUG DEPENDENCE 1 INDICATIONS AND USAGE

- 1.1 Special Diagnostic Considerations
- 1.2 Need for Comprehensive Treatment Program

2 DOSAGE AND ADMINISTRATION

- 2.1 General Dosing Information
- 2.2 Patients New to Methylphenidate
- 2.3 Patients Currently Using Methylphenidate

- 2.4 Dose Titration
- 2.5 Maintenance/Extended Treatment
- 2.6 Dose Reduction and Discontinuation

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

- 4.1 Hypersensitivity to Methylphenidate
- 4.2 Agitation
- 4.3 Glaucoma
- 4.4 Tics
- 4.5 Monoamine Oxidase Inhibitors

5 WARNINGS AND PRECAUTIONS

- 5.1 Serious Cardiovascular Events
- 5.2 Psychiatric Adverse Events
- 5.3 Seizures
- 5.4 Priapism
- 5.5 Peripheral Vasculopathy, including Raynaud's Phenomenon
- 5.6 Long-Term Suppression of Growth
- 5.7 Visual Disturbance
- 5.8 Potential for Gastrointestinal Obstruction
- 5.9 Hematologic Monitoring

6 ADVERSE REACTIONS

6.1 Commonly Observed Adverse Reactions in Double-Blind, Placebo-Controlled Clinical Trials

6.2 Other Adverse Reactions Observed in Methylphenidate hydrochloride extendedrelease tablets Clinical Trials

6.3 Discontinuation Due to Adverse Reactions

6.4 Tics

- 6.5 Blood Pressure and Heart Rate Increases
- 6.6 Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 MAO Inhibitors
- 7.2 Vasopressor Agents

7.3 Coumarin Anticoagulants, Antidepressants, and Selective Serotonin Reuptake Inhibitors

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Labor and Delivery
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use

9 DRUG ABUSE AND DEPENDENCE

- 9.1 Controlled Substance
- 9.2 Abuse
- 9.3 Dependence

10 OVERDOSAGE

- 10.1 Signs and Symptoms
- 10.2 Recommended Treatment
- 10.3 Poison Control Center

11 DESCRIPTION

11.1 System Components and Performance

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

- 14.1 Children
- 14.2 Adolescents
- 14.3 Adults
- **15 REFERENCES**

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: DRUG DEPENDENCE

Methylphenidate hydrochloride extended-release tablets should be given cautiously to patients with a history of drug dependence or alcoholism. Chronic abusive use can lead to marked tolerance and psychological dependence with varying degrees of abnormal behavior. Frank psychotic episodes can occur, especially with parenteral abuse. Careful supervision is required during withdrawal from abusive use since severe depression may occur. Withdrawal following chronic therapeutic use may unmask symptoms of the underlying disorder that may require followup.

1 INDICATIONS AND USAGE

Methylphenidate hydrochloride extended-release tablets is indicated for the treatment of Attention Deficit Hyperactivity Disorder (ADHD) in children 6 years of age and older, adolescents, and adults up to the age of 65 [see Clinical Studies (14)].

A diagnosis of Attention Deficit Hyperactivity Disorder (ADHD; DSM-IV) implies the presence of hyperactive-impulsive or inattentive symptoms that caused impairment and were present before age 7 years. The symptoms must cause clinically significant impairment, e.g., in social, academic, or occupational functioning, and be present in two or more settings, e.g., school (or work) and at home. The symptoms must not be better accounted for by another mental disorder. For the Inattentive Type, at least six of the following symptoms must have persisted for at least 6 months: lack of attention to details/careless mistakes; lack of sustained attention; poor listener; failure to follow through on tasks; poor organization; avoids tasks requiring sustained mental effort; loses things; easily distracted; forgetful. For the Hyperactive-Impulsive Type, at least six of the following symptoms must have persisted for at least 6 months:

fidgeting/squirming; leaving seat; inappropriate running/climbing; difficulty with quiet activities; "on the go;" excessive talking; blurting answers; can't wait turn; intrusive. The Combined Type requires both inattentive and hyperactive-impulsive criteria to be met.

1.1 Special Diagnostic Considerations

Specific etiology of this syndrome is unknown, and there is no single diagnostic test. Adequate diagnosis requires the use of medical and special psychological, educational, and social resources. Learning may or may not be impaired. The diagnosis must be based upon a complete history and evaluation of the patient and not solely on the presence of the required number of DSM-IV characteristics.

1.2 Need for Comprehensive Treatment Program

Methylphenidate hydrochloride extended-release tablets is indicated as an integral part of a total treatment program for ADHD that may include other measures (psychological, educational, social). Drug treatment may not be indicated for all patients with ADHD. Stimulants are not intended for use in patients who exhibit symptoms secondary to environmental factors and/or other primary psychiatric disorders, including psychosis. Appropriate educational placement is essential and psychosocial intervention is often helpful. When remedial measures alone are insufficient, the decision to prescribe stimulant medication will depend upon the physician's assessment of the chronicity and severity of the patient's symptoms.

2 DOSAGE AND ADMINISTRATION

2.1 General Dosing Information

Methylphenidate hydrochloride extended-release tablets should be administered orally once daily in the morning with or without food.

Methylphenidate hydrochloride extended-release tablets must be swallowed whole with the aid of liquids, and must not be chewed, divided, or crushed [see Patient Counseling Information (17)].

2.2 Patients New to Methylphenidate

The recommended starting dose of methylphenidate hydrochloride extended-release tablets for patients who are not currently taking methylphenidate or stimulants other than methylphenidate is 18 mg once daily for children and adolescents and 18 or 36 mg once daily for adults (see Table 1).

Table 1.	Methylphenidate Hydrochloride Extended-Release Tablets Recommended Starting Doses
and Dose	Ranges

Patient Age	Recommended Starting Dose	Dose Range
Children 6-12 years of age	18 mg/day	18 mg - 54 mg/day
Adolescents 13-17 years of age	18 mg/day	18 mg - 72 mg/day not to exceed 2 mg/kg/day
Adults 18-65 years of age	18 or 36 mg/day	18 mg - 72 mg/day
Adults 18-60 years of age	18 of 36 mg/day	18 mg - 72 mg/day

2.3 Patients Currently Using Methylphenidate

The recommended dose of methylphenidate hydrochloride extended-release tablets for patients who are currently taking methylphenidate twice daily or three times daily at doses of 10 to 60 mg/day is provided in Table 2. Dosing recommendations are based on current dose regimen and clinical judgment. Conversion dosage should not exceed 72 mg daily.

Hydrochloride Extended-Release Tablets	
Descrives Mathedala and data Daila Data	Recommended Methylphenidate
Previous Methylphenidate Daily Dose	hydrochloride extended-release tablets
이 아이지 이 가지 않는 것이 아이지 않는 것이 아이지 않는 것이 있는 것이 아이지 않는 것이 있는 것이 있다. 이 가지 않는 것이 있는 것이 있다. 이 가지 않는 것이 있는 것이 있다. 가지 않는 가지 않는 것이 있다. 가지 않는 것이 없다. 가지 이 같이 않는 것이 없다. 가지 않는 것이 없다. 것이 없다. 것이 않는 것이 없다. 것이 없다. 것이 없다. 가지 않는 것이 없다. 것이 않다. 것이 없다. 것이 않다. 것이 없다. 것이 없다. 것이 없다. 것이 없다. 것이 없다. 않다. 것이 없다. 하 것이 않아, 것이 않아, 것이 않아, 것이 않아, 것이 없다. 것이 않아, 않아, 것이 없다. 것이 없다. 것이 없다. 것이 없다. 않아, 것이 없다. 것	Starting Dose
5 mg Methylphenidate twice daily or three times daily	18 mg every morning
10 mg Methylphenidate twice daily or three times daily	36 mg every morning
15 mg Methylphenidate twice daily or three times daily	54 mg every morning
20 mg Methylphenidate twice daily or three times daily	72 mg every morning

Other methylphenidate regimens: Clinical judgment should be used when selecting the starting dose.

2.4 Dose Titration

Doses may be increased in 18 mg increments at weekly intervals for patients who have not achieved an optimal response at a lower dose. Daily dosages above 54 mg in children and 72 mg in adolescents have not been studied and are not recommended. Daily dosages above 72 mg in adults are not recommended.

A 27 mg dosage strength is available for physicians who wish to prescribe between the 18 mg and 36 mg dosages.

2.5 Maintenance/Extended Treatment

There is no body of evidence available from controlled trials to indicate how long the patient with ADHD should be treated with methylphenidate hydrochloride extended-release tablets. It is generally agreed, however, that pharmacological treatment of ADHD may be needed for extended periods.

The effectiveness of methylphenidate hydrochloride extended-release tablets for longterm use, i.e., for more than 7 weeks, has not been systematically evaluated in controlled trials. The physician who elects to use methylphenidate hydrochloride extended-release tablets for extended periods in patients with ADHD should periodically re-evaluate the long-term usefulness of the drug for the individual patient with trials off medication to assess the patient's functioning without pharmacotherapy. Improvement may be sustained when the drug is either temporarily or permanently discontinued.

2.6 Dose Reduction and Discontinuation

If paradoxical aggravation of symptoms or other adverse events occur, the dosage should be reduced, or, if necessary, the drug should be discontinued.

If improvement is not observed after appropriate dosage adjustment over a one-month period, the drug should be discontinued.

3 DOSAGE FORMS AND STRENGTHS

Methylphenidate Hydrochloride Extended-Release Tablets, USP are available in 18 mg, 27 mg, 36 mg and 54 mg of methylphenidate hydrochloride, USP.

- The 18 mg tablets are light yellow to yellow film coated round cylindrical biconvex tablets printed with 212 in black ink.
- The 27 mg tablets are light pink to pink film coated round cylindrical biconvex tablets printed with 213 in black ink.
- The 36 mg tablets are White to off white film coated round cylindrical biconvex tablets printed with 214 in black ink.
- The 54 mg tablets are light to dark brown film coated round cylindrical biconvex tablets printed with 215 in black ink.

4 CONTRAINDICATIONS

4.1 Hypersensitivity to Methylphenidate

Hypersensitivity reactions, such as angioedema and anaphylactic reactions, have been observed in patients treated with methylphenidate hydrochloride extended-release tablets. Therefore, methylphenidate hydrochloride extended-release tablets are contraindicated in patients known to be hypersensitive to methylphenidate or other components of the product [see Adverse Reactions (6.6)].

4.2 Agitation

Methylphenidate hydrochloride extended-release tablets are contraindicated in patients with marked anxiety, tension, and agitation, since the drug may aggravate these symptoms.

4.3 Glaucoma

Methylphenidate hydrochloride extended-release tablets are contraindicated in patients with glaucoma.

4.4 Tics

Methylphenidate hydrochloride extended-release tablets are contraindicated in patients with motor tics or with a family history or diagnosis of Tourette's syndrome [see Adverse Reactions (6.4)].

4.5 Monoamine Oxidase Inhibitors

Methylphenidate hydrochloride extended-release tablets are contraindicated during treatment with monoamine oxidase (MAO) inhibitors, and also within a minimum of 14 days following discontinuation of a MAO inhibitor (hypertensive crises may result) [see Drug Interactions (7.1)].

5 WARNINGS AND PRECAUTIONS

5.1 Serious Cardiovascular Events

Sudden Death and Preexisting Structural Cardiac Abnormalities or Other Serious Heart Problems

Children and Adolescents

Sudden death has been reported in association with CNS stimulant treatment at usual doses in children and adolescents with structural cardiac abnormalities or other serious heart problems. Although some serious heart problems alone carry an increased risk of sudden death, stimulant products generally should not be used in children or adolescents with known serious structural cardiac abnormalities, cardiomyopathy, serious heart rhythm abnormalities, or other serious cardiac problems that may place them at increased vulnerability to the sympathomimetic effects of a stimulant drug.

Adults

Sudden deaths, stroke, and myocardial infarction have been reported in adults taking stimulant drugs at usual doses for ADHD. Although the role of stimulants in these adult cases is also unknown, adults have a greater likelihood than children of having serious structural cardiac abnormalities, cardiomyopathy, serious heart rhythm abnormalities, coronary artery disease, or other serious cardiac problems. Adults with such abnormalities should also generally not be treated with stimulant drugs.

Hypertension and Other Cardiovascular Conditions

Stimulant medications cause a modest increase in average blood pressure (about 2 to 4 mm Hg) and average heart rate (about 3 to 6 bpm) [see Adverse Reactions (6.5)], and individuals may have larger increases. While the mean changes alone would not be expected to have short-term consequences, all patients should be monitored for larger changes in heart rate and blood pressure. Caution is indicated in treating patients whose underlying medical conditions might be compromised by increases in blood pressure or heart rate, e.g., those with preexisting hypertension, heart failure, recent myocardial infarction, or ventricular arrhythmia.

Assessing Cardiovascular Status in Patients Being Treated with Stimulant Medications

Children, adolescents, or adults who are being considered for treatment with stimulant medications should have a careful history (including assessment for a family history of sudden death or ventricular arrhythmia) and physical exam to assess for the presence of cardiac disease, and should receive further cardiac evaluation if findings suggest such disease (e.g., electrocardiogram and echocardiogram). Patients who develop symptoms such as exertional chest pain, unexplained syncope, or other symptoms suggestive of cardiac disease during stimulant treatment should undergo a prompt cardiac evaluation.

5.2 Psychiatric Adverse Events

Preexisting Psychosis

Administration of stimulants may exacerbate symptoms of behavior disturbance and thought disorder in patients with a preexisting psychotic disorder.

Bipolar Illness

Particular care should be taken in using stimulants to treat ADHD in patients with comorbid bipolar disorder because of concern for possible induction of a mixed/manic episode in such patients. Prior to initiating treatment with a stimulant, patients with comorbid depressive symptoms should be adequately screened to determine if they are at risk for bipolar disorder; such screening should include a detailed psychiatric history, including a family history of suicide, bipolar disorder, and depression.

Emergence of New Psychotic or Manic Symptoms

Treatment-emergent psychotic or manic symptoms, e.g., hallucinations, delusional thinking, or mania in patients without a prior history of psychotic illness or mania can be caused by stimulants at usual doses. If such symptoms occur, consideration should be given to a possible causal role of the stimulant, and discontinuation of treatment may be appropriate. In a pooled analysis of multiple short-term, placebo-controlled studies, such symptoms occurred in about 0.1% (4 patients with events out of 3,482 exposed to methylphenidate or amphetamine for several weeks at usual doses) of stimulant-treated patients compared to 0 in placebo-treated patients.

Aggression

Aggressive behavior or hostility is often observed in patients with ADHD, and has been reported in clinical trials and the postmarketing experience of some medications indicated for the treatment of ADHD. Although there is no systematic evidence that stimulants cause aggressive behavior or hostility, patients beginning treatment for ADHD should be monitored for the appearance of or worsening of aggressive behavior or hostility.

5.3 Seizures

There is some clinical evidence that stimulants may lower the convulsive threshold in patients with prior history of seizures, in patients with prior EEG abnormalities in absence of seizures, and, very rarely, in patients without a history of seizures and no prior EEG evidence of seizures. In the presence of seizures, the drug should be discontinued.

5.4 Priapism

Prolonged and painful erections, sometimes requiring surgical intervention, have been reported with methylphenidate products, including Methylphenidate hydrochloride extended-release tablets, in both pediatric and adult patients [see Adverse Reactions (6.6)]. Priapism was not reported with drug initiation but developed after some time on the drug, often subsequent to an increase in dose. Priapism has also appeared during a period of drug withdrawal (drug holidays or during discontinuation). Patients who develop abnormally sustained or frequent and painful erections should seek immediate medical attention.

5.5 Peripheral Vasculopathy, including Raynaud's Phenomenon

Stimulants, including Methylphenidate hydrochloride extended-release tablets, used to treat ADHD are associated with peripheral vasculopathy, including Raynaud's phenomenon. Signs and symptoms are usually intermittent and mild; however, very rare sequelae include digital ulceration and/or soft tissue breakdown. Effects of peripheral vasculopathy, including Raynaud's phenomenon, were observed in post-marketing reports at different times and at therapeutic doses in all age groups throughout the course of treatment. Signs and symptoms generally improve after reduction in dose or discontinuation of drug. Careful observation for digital changes is necessary during treatment with ADHD stimulants. Further clinical evaluation (e.g., rheumatology referral)

may be appropriate for certain patients.

5.6 Long-Term Suppression of Growth

Careful follow-up of weight and height in children ages 7 to 10 years who were randomized to either methylphenidate or nonmedication treatment groups over 14 months, as well as in naturalistic subgroups of newly methylphenidate-treated and nonmedication-treated children over 36 months (to the ages of 10 to 13 years), suggests that consistently medicated children (i.e., treatment for 7 days per week throughout the year) have a temporary slowing in growth rate (on average, a total of about 2 cm less growth in height and 2.7 kg less growth in weight over 3 years), without evidence of growth rebound during this period of development. Published data are inadequate to determine whether chronic use of amphetamines may cause similar suppression of growth; however, it is anticipated that they likely have this effect as well. Therefore, growth should be monitored during treatment with stimulants, and patients who are not growing or gaining height or weight as expected may need to have their treatment interrupted.

5.7 Visual Disturbance

Difficulties with accommodation and blurring of vision have been reported with stimulant treatment.

5.8 Potential for Gastrointestinal Obstruction

Because the methylphenidate hydrochloride extended-release tablet is nondeformable and does not appreciably change in shape in the GI tract, Methylphenidate hydrochloride extended-release tablets should not ordinarily be administered to patients with preexisting severe gastrointestinal narrowing (pathologic or iatrogenic, for example: esophageal motility disorders, small bowel inflammatory disease, "short gut" syndrome due to adhesions or decreased transit time, past history of peritonitis, cystic fibrosis, chronic intestinal pseudo-obstruction, or Meckel's diverticulum). There have been rare reports of obstructive symptoms in patients with known strictures in association with the ingestion of drugs in nondeformable controlled-release formulations. Due to the controlled-release design of the tablet, methylphenidate hydrochloride extended-release tablets should be used only in patients who are able to swallow the tablet whole [see Patient Counseling Information (17)].

5.9 Hematologic Monitoring

Periodic CBC, differential, and platelet counts are advised during prolonged therapy.

6 ADVERSE REACTIONS

The following are discussed in more detail in other sections of the labeling:

- Drug Dependence [see Box Warning]
- Hypersensitivity to Methylphenidate [see Contraindications (4.1)]
- Agitation [see Contraindications (4.2)]
- Glaucoma [see Contraindications (4.3)]
- Tics [see Contraindications (4.4)]
- Monoamine Oxidase Inhibitors [see Contraindications (4.5) and Drug Interactions

(7.1)]

- Serious Cardiovascular Events [see Warnings and Precautions (5.1)]
- Psychiatric Adverse Events [see Warnings and Precautions (5.2)]
- Seizures [see Warnings and Precautions (5.3)]
- Priapism [see Warnings and Precautions (5.4)]
- Long-Term Suppression of Growth [see Warnings and Precautions (5.6)]
- Visual Disturbance [see Warnings and Precautions (5.7)]
- Potential for Gastrointestinal Obstruction [see Warnings and Precautions (5.8)]
- Hematologic Monitoring [see Warnings and Precautions (5.9)]

The most common adverse reaction in double-blind clinical trials (>5%) in pediatric patients (children and adolescents) was abdominal pain upper. The most common adverse reactions in double-blind clinical trials (>5%) in adult patients were decreased appetite, headache, dry mouth, nausea, insomnia, anxiety, dizziness, weight decreased, irritability, and hyperhidrosis [see Adverse Reactions (6.1)].

The most common adverse reactions associated with discontinuation (\geq 1%) from either pediatric or adult clinical trials were anxiety, irritability, insomnia, and blood pressure increased [see Adverse Reactions (6.3)].

The development program for methylphenidate hydrochloride extended-release tablets included exposures in a total of 3,906 participants in clinical trials. Children, adolescents, and adults with ADHD were evaluated in 6 controlled clinical studies and 11 open-label clinical studies (see Table 3). Safety was assessed by collecting adverse events, vital signs, weights, and ECGs, and by performing physical examinations and laboratory analyses.

Patient Population	N	Dose Range
Children	2,216	18 to 54 mg once daily
Adolescents	502	18 to 72 mg once daily
Adults	1,188	18 to 108 mg once daily

Table 3. Methylphenidate hydrochloride extended-release tablets Exposure inDouble-Blind and Open-Label Clinical Studies

Adverse events during exposure were obtained primarily by general inquiry and recorded by clinical investigators using their own terminology. Consequently, to provide a meaningful estimate of the proportion of individuals experiencing adverse events, events were grouped in standardized categories using MedDRA terminology.

The stated frequencies of adverse events represent the proportion of individuals who experienced, at least once, a treatment-emergent adverse event of the type listed. An event was considered treatment-emergent if it occurred for the first time or worsened while receiving therapy following baseline evaluation.

Throughout this section, adverse reactions are reported. Adverse reactions are adverse events that were considered to be reasonably associated with the use of methylphenidate hydrochloride extended-release tablets based on the comprehensive assessment of the available adverse event information. A causal association for methylphenidate hydrochloride extended-release tablets often cannot be reliably established in individual cases. Further, because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in clinical trials of another drug and may not reflect the rates observed in clinical practice.

The majority of adverse reactions were mild to moderate in severity.

6.1 Commonly Observed Adverse Reactions in Double-Blind, Placebo-Controlled Clinical Trials

Adverse reactions in either the pediatric or adult double-blind adverse reactions tables may be relevant for both patient populations.

Children and Adolescents

Table 4 lists the adverse reactions reported in 1% or more of methylphenidate hydrochloride extended-release tablets treated children and adolescent subjects in 4 placebo-controlled, double-blind clinical trials.

System/Organ Class Adverse Reaction	Methylphenidate hydrochloride extended-release tablets	Placebo	
	(n=321) %	(n=318) %	
Gastrointestinal Disorders			
Abdominal pain upper	6.2	3.8	
Vomiting	2.8	1.6	
General Disorders and Administration			
Site Conditions			
Pyrexia	2.2	0.9	
Infections and Infestations			
Nasopharyngitis	2.8	2.2	
Nervous System Disorders			
Dizziness	1.9	0	
Psychiatric Disorders			
Insomnia*	2.8	0.3	
Respiratory, Thoracic and Mediastinal			
Disorders			
Cough	1.9	0.9	
Oropharyngeal pain	1.2	0.9	

*Terms of Initial insomnia (Methylphenidate hydrochloride extended-release tablets =0.6%) and Insomnia (Methylphenidate hydrochloride extended-release tablets =2.2%) are combined into Insomnia

The majority of adverse reactions were mild to moderate in severity.

Adults

Table 5 lists the adverse reactions reported in 1% or more of methylphenidate hydrochloride extended-release tablets-treated adults in 2 placebo-controlled, double-blind clinical trials.

Table5. Adverse Reactions Reported by ≥1% of Methylphenidate Hydrochloride Extended-Release Tablets-Treated Adult Subjects in 2 Placebo-Controlled, Double-Blind Clinical Trials*

System/Organ Class Adverse Reaction	Methylphenidate hydrochloride extended-release tablets (n=415) %	Placeb (n=212 %
Cardiac Disorders		1.0
Tachycardia	4.8	0
Palpitations	3.1	0.9
Ear and Labyrinth Disorders		
Vertigo	1.7	0
Eye Disorders		
Vision blurred	1.7	0.5
GastrointestinalDisorders		
Dry mouth	14.0	3.8
Nausea	12.8	3.3
Dyspepsia	2.2	0.9
Vomiting	1.7	0.5
Constipation	1.4	0.9
General Disorders and Administration Site		
Conditions	0.002	1992
Irritability	5.8	1.4
Infections and Infestations		
Upper respiratory tract infection	2.2	0.9
Investigations		
Weight decreased	6.5	3.3
Visite ballism and Nutatilian Discussion		
Metabolism and Nutrition Disorders Decreased appetite	25.3	6.6
Anorexia	1.7	0
Allorena		-
Musculoskeletal and Connective Tissue Disorders		
Muscle tightness	1.9	0
Nervous System Disorders		
Headache	22.2	15.6
Dizziness	6.7	5.2
Tremor	2.7	0.5
Paresthesia	1.2	0
Sedation	1.2	0
Tension headache	1.2	0.5
Psychiatric Disorders		
Insomnia	12.3	6.1
Anxiety	8.2	2.4
Initial insomnia	4.3	2.8
Depressed mood	3.9	1.4
Nervousness	3.1	0.5
Restlessness	3.1	0
Agitation	2.2	0.5
Aggression	1.7	0.5
Bruxism	1.7	0.5
Depression	1.7	0.9
Libido decreased Affectlability	1.7	0.9
Confusional state	1.4	0.5
Tension	12	0.5
Respiratory, Thoracic and Mediastinal Disorders		
Oropharyngeal pain	1.7	1.4
Skin and Subcutaneous Tissue Disorders	51	0.0
Hyperhidrosis	5.1	0.9

* Included doses up to 108 mg.

The majority of ADRs were mild to moderate in severity.

6.2 Other Adverse Reactions Observed in Methylphenidate hydrochloride extended-release tablets Clinical Trials

This section includes adverse reactions reported by methylphenidate hydrochloride extended-release tablets-treated subjects in double-blind trials that do not meet the criteria specified for Table 4 or Table 5 and all adverse reactions reported by methylphenidate hydrochloride extended-release tablets-treated subjects who participated in open-label and postmarketing clinical trials. Blood and Lymphatic System Disorders: Leukopenia

Eye Disorders: Accommodation disorder, Dry eye

Vascular Disorders: Hot flush

Gastrointestinal Disorders: Abdominal discomfort, Abdominal pain, Diarrhea

General Disorders and Administrative Site Conditions: Asthenia, Fatigue, Feeling jittery, Thirst

Infections and Infestations: Sinusitis

Investigations: Alanine aminotransferase increased, Blood pressure increased, Cardiac murmur, Heart rate increased

Musculoskeletal and Connective Tissue Disorders: Muscle spasms

Nervous System Disorders: Lethargy, Psychomotor hyperactivity, Somnolence

Psychiatric Disorders: Anger, Hypervigilance, Mood altered, Mood swings, Panic attack, Sleep disorder, Tearfulness, Tic

Reproductive System and Breast Disorders: Erectile dysfunction

Respiratory, Thoracic and Mediastinal Disorders: Dyspnea

Skin and Subcutaneous Tissue Disorders: Rash, Rash macular

Vascular Disorders: Hypertension

6.3 Discontinuation Due to Adverse Reactions

Adverse reactions in the 4 placebo-controlled studies of children and adolescents leading to discontinuation occurred in 2 methylphenidate hydrochloride extended-release tablets patients (0.6%) including depressed mood (1, 0.3%) and headache and insomnia (1, 0.3%), and 6 placebo patients (1.9%) including headache and insomnia (1, 0.3%), irritability (2, 0.6%), headache (1, 0.3%), psychomotor hyperactivity (1, 0.3%), and tic (1, 0.3%).

In the 2 placebo-controlled studies of adults, 25 methylphenidate hydrochloride extended-release tablets patients (6.0%) and 6 placebo patients (2.8%) discontinued due to an adverse reaction. Those events with an incidence of >0.5% in the Methylphenidate hydrochloride extended-release tablets patients included anxiety (1.7%), irritability (1.4%), blood pressure increased (1.0%), and nervousness (0.7%). In placebo patients, blood pressure increased and depressed mood had an incidence of >0.5% (0.9%).

In the 11 open-label studies of children, adolescents, and adults, 266 methylphenidate hydrochloride extended-release tablets patients (7.0%) discontinued due to an adverse reaction. Those events with an incidence of >0.5% included insomnia (1.2%), irritability (0.8%), anxiety (0.7%), decreased appetite (0.7%), and tic (0.6%).

6.4 Tics

In a long-term uncontrolled study (n=432 children), the cumulative incidence of new onset of tics was 9% after 27 months of treatment with methylphenidate hydrochloride extended-release tablets.

In a second uncontrolled study (n=682 children) the cumulative incidence of new-onset tics was 1% (9/682 children). The treatment period was up to 9 months with mean treatment duration of 7.2 months.

6.5 Blood Pressure and Heart Rate Increases

In the laboratory classroom clinical trials in children (Studies 1 and 2), both methylphenidate hydrochloride extended-release tablets once daily and methylphenidate three times daily increased resting pulse by an average of 2 to 6 bpm and produced average increases of systolic and diastolic blood pressure of roughly 1 to 4 mm Hg during the day, relative to placebo. In the placebo-controlled adolescent trial (Study 4), mean increases from baseline in resting pulse rate were observed with methylphenidate hydrochloride extended-release tablets and placebo at the end of the double-blind phase (5 and 3 beats/minute, respectively). Mean increases from baseline in blood pressure at the end of the double-blind phase for methylphenidate hydrochloride extended-release tablets and placebo-treated patients were 0.7 and 0.7 mm Hg (systolic) and 2.6 and 1.4 mm Hg (diastolic), respectively. In one placebo-controlled study in adults (Study 6), dose-dependent mean increases of 3.9 to 9.8 bpm from baseline in standing pulse rate were observed with methylphenidate hydrochloride extended-release tablets at the end of the double-blind treatment vs. an increase of 2.7 beats/minute with placebo. Mean changes from baseline in standing blood pressure at the end of double-blind treatment ranged from 0.1 to 2.2 mm Hg (systolic) and 0.7 to 2.2 mm Hg (diastolic) for methylphenidate hydrochloride extended-release tablets and was 1.1 mm Hg (systolic) and -1.8 mm Hg (diastolic) for placebo. In a second placebo-controlled study in adults (Study 5), mean changes from baseline in resting pulse rate were observed for methylphenidate hydrochloride extended-release tablets and placebo at the end of the double-blind treatment (3.6 and -1.6 beats/minute, respectively). Mean changes from baseline in blood pressure at the end of the double-blind treatment for methylphenidate hydrochloride extended-release tablets and placebo-treated patients were -1.2 and -0.5 mm Hg (systolic) and 1.1 and 0.4 mm Hg (diastolic), respectively [see Warnings and Precautions (5.1)].

6.6 Postmarketing Experience

The following additional adverse reactions have been identified during postapproval use of methylphenidate hydrochloride extended-release tablets. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency:

Blood and Lymphatic System Disorders: Pancytopenia, Thrombocytopenia, Thrombocytopenic purpura

Cardiac Disorders: Angina pectoris, Bradycardia, Extrasystoles, Supraventricular tachycardia, Ventricular extrasystoles

Eye Disorders: Diplopia, Mydriasis, Visual impairment

General Disorders: Chest pain, Chest discomfort, Drug effect decreased, Hyperpyrexia, Therapeutic response decreased

Hepatobiliary disorders: Hepatocellular injury, Acute hepatic failure

Immune System Disorders: Hypersensitivity reactions such as Angioedema, Anaphylactic reactions, Auricular swelling, Bullous conditions, Exfoliative conditions, Urticarias, Pruritus NEC, Rashes, Eruptions, and Exanthemas NEC

Investigations: Blood alkaline phosphatase increased, Blood bilirubin increased, Hepatic enzyme increased, Platelet count decreased, White blood cell count abnormal

Musculoskeletal, Connective Tissue and Bone Disorders: Arthralgia, Myalgia, Muscle twitching, Rhabdomyolysis

Nervous System Disorders: Convulsion, Grand mal convulsion, Dyskinesia, Serotonin syndrome in combination with serotonergic drugs

Psychiatric Disorders: Disorientation, Hallucination, Hallucination auditory, Hallucination visual, Mania, Logorrhea, Libido changes

Reproductive System and Breast Disorders: Priapism

Skin and Subcutaneous Tissue Disorders: Alopecia, Erythema

Vascular Disorders: Raynaud's phenomenon

To report SUSPECTED ADVERSE REACTIONS contact AvKARE at 1-855-361-3993; email drugsafety@avkare.com; or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

7 DRUG INTERACTIONS

7.1 MAO Inhibitors

Methylphenidate hydrochloride extended-release tablets should not be used in patients being treated (currently or within the preceding 2 weeks) with MAO inhibitors [see Contraindications (4.5)].

7.2 Vasopressor Agents

Because of possible increases in blood pressure, methylphenidate hydrochloride extended-release tablets should be used cautiously with vasopressor agents [see Warnings and Precautions (5.1)].

7.3 Coumarin Anticoagulants, Antidepressants, and Selective Serotonin Reuptake Inhibitors

Human pharmacologic studies have shown that methylphenidate may inhibit the metabolism of coumarin anticoagulants, anticonvulsants (eg, phenobarbital, phenytoin, primidone), and some antidepressants (tricyclics and selective serotonin reuptake inhibitors). Downward dose adjustment of these drugs may be required when given concomitantly with methylphenidate. It may be necessary to adjust the dosage and monitor plasma drug concentrations (or, in the case of coumarin, coagulation times), when initiating or discontinuing concomitant methylphenidate.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category C

Methylphenidate has been shown to have teratogenic effects in rabbits when given in doses of 200 mg/kg/day, which is approximately 100 times and 40 times the maximum recommended human dose on a mg/kg and mg/m² basis, respectively.

A reproduction study in rats revealed no evidence of harm to the fetus at oral doses up to 30 mg/kg/day, approximately 15-fold and 3-fold the maximum recommended human dose of methylphenidate hydrochloride extended-release tablets on a mg/kg and mg/m ² basis, respectively. The approximate plasma exposure to methylphenidate plus its main metabolite PPAA in pregnant rats was 1-2 times that seen in trials in volunteers and patients with the maximum recommended dose of methylphenidate hydrochloride extended-release tablets based on the AUC.

The safety of methylphenidate for use during human pregnancy has not been established. There are no adequate and well-controlled studies in pregnant women. Methylphenidate hydrochloride extended-release tablets should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

8.2 Labor and Delivery

The effect of methylphenidate hydrochloride extended-release tablets on labor and delivery in humans is unknown.

8.3 Nursing Mothers

It is not known whether methylphenidate is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised if methylphenidate hydrochloride extended-release tablets is administered to a nursing woman.

In lactating female rats treated with a single oral dose of 5 mg/kg radiolabeled methylphenidate, radioactivity (representing methylphenidate and/or its metabolites) was observed in milk and levels were generally similar to those in plasma.

8.4 Pediatric Use

Methylphenidate hydrochloride extended-release tablets should not be used in children under six years, since safety and efficacy in this age group have not been established. Long-term effects of methylphenidate in children have not been well established.

8.5 Geriatric Use

Methylphenidate hydrochloride extended-release tablets has not been studied in patients greater than 65 years of age.

9 DRUG ABUSE AND DEPENDENCE

9.1 Controlled Substance

Methylphenidate is a Schedule II controlled substance under the Controlled Substances Act.

9.2 Abuse

As noted in the Box Warning, methylphenidate hydrochloride extended-release tablets

should be given cautiously to patients with a history of drug dependence or alcoholism. Chronic abusive use can lead to marked tolerance and psychological dependence with varying degrees of abnormal behavior. Frank psychotic episodes can occur, especially with parenteral abuse.

In two placebo-controlled human abuse potential studies, single oral doses of methylphenidate hydrochloride extended-release tablets were compared to single oral doses of immediate-release methylphenidate (IR MPH) and placebo in subjects with a history of recreational stimulant use to assess relative abuse potential. For the purpose of this assessment, the response for each of the subjective measures was defined as the maximum effect within the first 8 hours after dose administration.

In one study (n=40), both methylphenidate hydrochloride extended-release tablets (108 mg) and 60 mg IR MPH compared to placebo produced statistically significantly greater responses on the five subjective measures suggestive of abuse potential. In comparisons between the two active treatments, however, methylphenidate hydrochloride extended-release tablets (108 mg) produced variable responses on positive subjective measures that were either statistically indistinguishable from (Abuse Potential, Drug Liking, Amphetamine, and Morphine Benzedrine Group [Euphoria]) or statistically less than (Stimulation – Euphoria) responses produced by 60 mg IR MPH.

In another study (n=49), both doses of methylphenidate hydrochloride extendedrelease tablets (54 mg and 108 mg) and both doses of IR MPH (50 mg and 90 mg) produced statistically significantly greater responses compared to placebo on the two primary scales used in the study (Drug Liking, Euphoria). When doses of Methylphenidate hydrochloride extended-release tablets (54 mg and 108 mg) were compared to IR MPH (50 mg and 90 mg), respectively, methylphenidate hydrochloride extended-release tablets produced statistically significantly lower subjective responses on these two scales than IR MPH. Methylphenidate hydrochloride extended-release tablets (108 mg) produced responses that were statistically indistinguishable from the responses on these two scales produced by IR MPH (50 mg). Differences in subjective responses to the respective doses should be considered in the context that only 22% of the total amount of methylphenidate in methylphenidate hydrochloride extended-release tablets is available for immediate release from the drug overcoat [see System *Components and Performance (11.1)*].

Although these findings reveal a relatively lower response to methylphenidate hydrochloride extended-release tablets on subjective measures suggestive of abuse potential compared to IR MPH at roughly equivalent total MPH doses, the relevance of these findings to the abuse potential of methylphenidate hydrochloride extended-release tablets in the community is unknown.

9.3 Dependence

As noted in the Box Warning, careful supervision is required during withdrawal from abusive use since severe depression may occur. Withdrawal following chronic therapeutic use may unmask symptoms of the underlying disorder that may require follow-up.

10 OVERDOSAGE

10.1 Signs and Symptoms

Signs and symptoms of methylphenidate hydrochloride extended-release tablets overdosage, resulting principally from overstimulation of the CNS and from excessive sympathomimetic effects, may include the following: vomiting, agitation, muscle twitching, convulsion, grand mal convulsion, confusional state, hallucinations (auditory and/or visual), hyperhidrosis, headache, pyrexia, tachycardia, palpitations, heart rate increased, sinus arrhythmia, hypertension, rhabdomyolysis, mydriasis, and dry mouth.

10.2 Recommended Treatment

Treatment consists of appropriate supportive measures. The patient must be protected against self-injury and against external stimuli that would aggravate overstimulation already present. Gastric contents may be evacuated by gastric lavage as indicated. Before performing gastric lavage, control agitation and seizures if present and protect the airway. Other measures to detoxify the gut include administration of activated charcoal and a cathartic. Intensive care must be provided to maintain adequate circulation and respiratory exchange; external cooling procedures may be required for pyrexia.

Efficacy of peritoneal dialysis or extracorporeal hemodialysis for methylphenidate hydrochloride extended-release tablets overdosage has not been established.


The prolonged release of methylphenidate from methylphenidate hydrochloride extended-release tablets should be considered when treating patients with overdose.

10.3 Poison Control Center

As with the management of all overdosage, the possibility of multiple-drug ingestion should be considered. The physician may wish to consider contacting a poison control center for up-to-date information on the management of overdosage with methylphenidate.

11 DESCRIPTION

Methylphenidate hydrochloride extended-release tablets, USP is a central nervous system (CNS) stimulant. Methylphenidate hydrochloride extended-release tablets are available in four tablet strengths. Each extended-release tablet for once-a-day oral administration contains 18, 27, 36, or 54 mg of methylphenidate HCI USP and is designed to have a 12-hour duration of effect. Chemically, methylphenidate HCI is d, I (racemic) methyl α -phenyl-2-piperidineacetate hydrochloride. Its empirical formula is C ${}_{14}H {}_{19}NO {}_{2}$ ·HCI. Its structural formula is:



Methylphenidate HCI USP is a white to off-white crystalline powder. Its solutions are acid to litmus. It is freely soluble in water and in methanol, soluble in alcohol, and slightly soluble in chloroform and in acetone. Its molecular weight is 269.77.

Methylphenidate hydrochloride extended-release tablets, USP also contains the following inert ingredients and are common to all strengths: butylated hydroxytoluene, cellulose acetate, hypromellose, phosphoric acid, polyethylene glycol, polyethylene oxides, povidone, propylene glycol, sodium chloride, stearic acid, succinic acid, ferric oxide yellow, FD&C Red No 40 and titanium dioxide. The 18 mg tablet strength also contains iron oxide yellow and Polysorbate 80. The 27 mg tablet strength also contains iron oxide red. The 36 mg tablet strength also contains talc. The 54 mg tablet strength also contains contains iron oxide yellow, iron oxide red and talc.

Each tablet strength also contains black iron oxide, hypromellose and propylene glycol as imprinting ink.

USP Dissolution Test Pending.

11.1 System Components and Performance

Methylphenidate hydrochloride extended-release tablets uses osmotic pressure to deliver methylphenidate HCl at a controlled rate. The system, which resembles a conventional tablet in appearance, comprises an osmotically active trilayer core surrounded by a semipermeable membrane with an immediate-release drug overcoat. The trilayer core is composed of two drug layers containing the drug and excipients, and a push laver containing osmotically active components. There is a precision-laser drilled orifice on the drug-layer end of the tablet. In an aqueous environment, such as the gastrointestinal tract, the drug overcoat dissolves within one hour, providing an initial dose of methylphenidate. Water permeates through the membrane into the tablet core. As the osmotically active polymer excipients expand, methylphenidate is released through the orifice. The membrane controls the rate at which water enters the tablet core, which in turn controls drug delivery. Furthermore, the drug release rate from the system increases with time over a period of 6 to 7 hours due to the drug-concentration gradient incorporated into the two drug layers of methylphenidate hydrochloride extended-release tablets. The biologically inert components of the tablet remain intact during gastrointestinal transit and are eliminated in the stool as a tablet shell along with insoluble core components. It is possible that methylphenidate hydrochloride extendedrelease tablets may be visible on abdominal x-rays under certain circumstances, especially when digital enhancing techniques are utilized.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Methylphenidate HCl is a central nervous system (CNS) stimulant. The mode of therapeutic action in Attention Deficit Hyperactivity Disorder (ADHD) is not known. Methylphenidate is thought to block the reuptake of norepinephrine and dopamine into the presynaptic neuron and increase the release of these monoamines into the extraneuronal space.

12.2 Pharmacodynamics

Methylphenidate is a racemic mixture comprised of the d- and l-isomers. The d-isomer is more pharmacologically active than the l-isomer.


12.3 Pharmacokinetics

Absorption

Methylphenidate is readily absorbed. Following oral administration of methylphenidate hydrochloride extended-release tablets, plasma methylphenidate concentrations increase rapidly, reaching an initial maximum at about 1 hour, followed by gradual ascending concentrations over the next 5 to 9 hours, after which a gradual decrease begins. Mean times to reach peak plasma concentrations across all doses of methylphenidate hydrochloride extended-release tablets occurred between 6 and 10 hours.

Methylphenidate hydrochloride extended-release tablets once daily minimizes the fluctuations between peak and trough concentrations associated with immediate-release methylphenidate three times daily (see Figure 1). The relative bioavailability of methylphenidate hydrochloride extended-release tablets once daily and methylphenidate three times daily in adults is comparable.

Figure 1. Mean methylphenidate plasma concentrations in 36 adults, following a single dose of methylphenidate hydrochloride extended-release tablets 18 mg once daily and immediate-release methylphenidate 5 mg three times daily administered every 4 hours.



The mean single-dose pharmacokinetic parameters in 36 healthy adults following the administration of methylphenidate hydrochloride extended-release tablets 18 mg once daily and methylphenidate 5 mg three times daily are summarized in Table 6.

	Methylphenidate hydrochloride extended-release tablets (18 mg once daily)	Methylphenidate (5 mg three times daily) (n=35)		
Parameters	(n=36)			
C _{max} (ng/mL)	3.7 ± 1.0	4.2 ± 1.0		
$T_{max}(h)$	6.8 ± 1.8	6.5 ± 1.8		
AUC _{inf} (ng•h/mL)	41.8 ± 13.9	38.0 ± 11.0		
t _{1/2} (h)	3.5 ± 0.4	3.0 ± 0.5		

Table 6. Pharmacokinetic Parameters (Mean \pm SD) After Single Dose in Healthy Adults

The pharmacokinetics of methylphenidate hydrochloride extended-release tablets were evaluated in healthy adults following single- and multiple-dose administration (steady state) of doses up to 144 mg/day. The mean half-life was about 3.6 hours. No differences in the pharmacokinetics of methylphenidate hydrochloride extended-release tablets were noted following single and repeated once-daily dosing, indicating no significant drug accumulation. The AUC and t $_{1/2}$ following repeated once-daily dosing are similar to those following the first dose of methylphenidate hydrochloride extended-release tablets in a dose range of 18 to 144 mg.

Dose Proportionality

Following administration of methylphenidate hydrochloride extended-release tablets in single doses of 18, 36, and 54 mg/day to healthy adults, C $_{max}$ and AUC $_{(0-inf)}$ of d-methylphenidate were proportional to dose, whereas I-methylphenidate C $_{max}$ and AUC $_{(0-inf)}$ increased disproportionately with respect to dose. Following administration of methylphenidate hydrochloride extended-release tablets, plasma concentrations of the l-isomer were approximately 1/40 the plasma concentrations of the d-isomer.

In healthy adults, single and multiple dosing of once-daily methylphenidate hydrochloride extended-release tablets doses from 54 to 144 mg/day resulted in linear and dose-proportional increases in C _{max} and AUC _{inf} for total methylphenidate (MPH) and its major metabolite, α -phenyl-piperidine acetic acid (PPAA). There was no time dependency in the pharmacokinetics of methylphenidate. The ratio of metabolite (PPAA) to parent drug (MPH) was constant across doses from 54 to 144 mg/day, both after single dose and upon multiple dosing.

In a multiple-dose study in adolescent ADHD patients aged 13 to 16 administered their prescribed dose (18 to 72 mg/day) of methylphenidate hydrochloride extended-release tablets, mean C $_{\rm max}$ and AUC $_{\rm TAU}$ of d- and total methylphenidate increased proportionally with respect to dose.

Distribution

Plasma methylphenidate concentrations in adults and adolescents decline biexponentially following oral administration. The half-life of methylphenidate in adults and adolescents following oral administration of methylphenidate hydrochloride extended-release tablets was approximately 3.5 hours.

Metabolism and Excretion

In humans, methylphenidate is metabolized primarily by de-esterification to PPAA, which has little or no pharmacologic activity. In adults the metabolism of methylphenidate hydrochloride extended-release tablets once daily as evaluated by metabolism to PPAA is similar to that of methylphenidate three times daily. The metabolism of single and repeated once-daily doses of methylphenidate hydrochloride extended-release tablets is similar.

After oral dosing of radiolabeled methylphenidate in humans, about 90% of the radioactivity was recovered in urine. The main urinary metabolite was PPAA, accounting for approximately 80% of the dose.

Food Effects

In patients, there were no differences in either the pharmacokinetics or the pharmacodynamic performance of methylphenidate hydrochloride extended-release tablets when administered after a high-fat breakfast. There is no evidence of dose dumping in the presence or absence of food.

Alcohol Effect

An *in vitro* study was conducted to explore the effect of alcohol on the release

characteristics of methylphenidate from the methylphenidate hydrochloride extendedrelease tablets 18 mg tablet dosage form. At an alcohol concentration up to 40% there was no increased release of methylphenidate in the first hour. The results with the 18 mg tablet strength are considered representative of the other available tablet strengths.

Special Populations

Gender

In healthy adults, the mean dose-adjusted AUC $_{(0-inf)}$ values for methylphenidate hydrochloride extended-release tablets were 36.7 ng·h/mL in men and 37.1 ng·h/mL in women, with no differences noted between the two groups.

Race

In adults receiving methylphenidate hydrochloride extended-release tablets, doseadjusted AUC _(0-inf) was consistent across ethnic groups; however, the sample size may have been insufficient to detect ethnic variations in pharmacokinetics.

Age

Increase in age resulted in increased apparent oral clearance (CL/F) (58% increase in adolescents compared to children). Some of these differences could be explained by body weight differences among these populations. This suggests that subjects with higher body weight may have lower exposures of total methylphenidate at similar doses.

The pharmacokinetics of methylphenidate hydrochloride extended-release tablets have not been studied in children less than 6 years of age.

Renal Insufficiency

There is no experience with the use of methylphenidate hydrochloride extended-release tablets in patients with renal insufficiency. After oral administration of radiolabeled methylphenidate in humans, methylphenidate was extensively metabolized and approximately 80% of the radioactivity was excreted in the urine in the form of PPAA. Since renal clearance is not an important route of methylphenidate clearance, renal insufficiency is expected to have little effect on the pharmacokinetics of methylphenidate hydrochloride extended-release tablets.

Hepatic Insufficiency

There is no experience with the use of methylphenidate hydrochloride extended-release tablets in patients with hepatic insufficiency.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

In a lifetime carcinogenicity study carried out in B6C3F1 mice, methylphenidate caused an increase in hepatocellular adenomas and, in males only, an increase in hepatoblastomas at a daily dose of approximately 60 mg/kg/day. This dose is approximately 30 times and 4 times the maximum recommended human dose of methylphenidate hydrochloride extended-release tablets on a mg/kg and mg/m² basis, respectively. Hepatoblastoma is a relatively rare rodent malignant tumor type. There was no increase in total malignant hepatic tumors. The mouse strain used is sensitive to the development of hepatic tumors, and the significance of these results to humans is unknown.

Methylphenidate did not cause any increases in tumors in a lifetime carcinogenicity study carried out in F344 rats; the highest dose used was approximately 45 mg/kg/day, which is approximately 22 times and 5 times the maximum recommended human dose of methylphenidate hydrochloride extended-release tablets on a mg/kg and mg/m² basis, respectively.

In a 24-week carcinogenicity study in the transgenic mouse strain p53+/-, which is sensitive to genotoxic carcinogens, there was no evidence of carcinogenicity. Male and female mice were fed diets containing the same concentration of methylphenidate as in the lifetime carcinogenicity study; the high-dose groups were exposed to 60 to 74 mg/kg/day of methylphenidate.

Mutagenesis

Methylphenidate was not mutagenic in the *in vitro* Ames reverse mutation assay or the *in vitro* mouse lymphoma cell forward mutation assay. Sister chromatid exchanges and chromosome aberrations were increased, indicative of a weak clastogenic response, in an *in vitro* assay in cultured Chinese Hamster Ovary cells. Methylphenidate was negative *in vivo* in males and females in the mouse bone marrow micronucleus assay.

Impairment of Fertility

Methylphenidate did not impair fertility in male or female mice that were fed diets containing the drug in an 18-week Continuous Breeding study. The study was conducted at doses up to 160 mg/kg/day, approximately 80-fold and 8-fold the highest recommended human dose of Methylphenidate hydrochloride extended-release tablets on a mg/kg and mg/m² basis, respectively.

14 CLINICAL STUDIES

Methylphenidate hydrochloride extended-release tablets was demonstrated to be effective in the treatment of Attention Deficit Hyperactivity Disorder (ADHD) in 4 randomized, double-blind, placebo-controlled studies in children and adolescents and 2 double-blind placebo-controlled studies in adults who met the Diagnostic and Statistical Manual 4 th edition (DSM-IV) criteria for ADHD.


14.1 Children

Three double-blind, active- and placebo-controlled studies were conducted in 416 children aged 6 to 12 years. The controlled studies compared Methylphenidate hydrochloride extended-release tablets given once daily (18, 36, or 54 mg), methylphenidate given three times daily over 12 hours (15, 30, or 45 mg total daily dose), and placebo in two single-center, 3-week crossover studies (Studies 1 and 2) and in a multicenter, 4-week, parallel-group comparison (Study 3). The primary comparison of interest in all three trials was methylphenidate hydrochloride extended-release tablets versus placebo.

Symptoms of ADHD were evaluated by community schoolteachers using the Inattention/Overactivity with Aggression (IOWA) Conners scale. Statistically significant

reduction in the Inattention/Overactivity subscale versus placebo was shown consistently across all three controlled studies for methylphenidate hydrochloride extended-release tablets. The scores for methylphenidate hydrochloride extendedrelease tablets and placebo for the three studies are presented in Figure 2.


Figure2. Mean Community School Teacher IOWA Conners Inattention/Overactivity Scores with methylphenidate hydrochloride extended-release tablets once daily (18, 36, or 54 mg) and placebo. Studies 1 and 2 involved a 3-way crossover of 1 week per treatment arm. Study 3 involved 4 weeks of parallel-group treatments with a Last Observation Carried Forward analysis at week 4. Error bars represent the mean plus standard error of the mean.



In Studies 1 and 2, symptoms of ADHD were evaluated by laboratory schoolteachers using the SKAMP* laboratory school rating scale. The combined results from these two studies demonstrated statistically significant improvements in attention and behavior in patients treated with methylphenidate hydrochloride extended-release tablets versus placebo that were maintained through 12 hours after dosing. Figure 3 presents the laboratory schoolteacher SKAMP ratings for methylphenidate hydrochloride extendedrelease tablets and placebo.

*Swanson, Kotkin, Agler, M-Fynn, and Pelham

Figure 3. Laboratory School Teacher SKAMP Ratings: Mean (SEM) of Combined Attention (Studies 1 and 2)



14.2 Adolescents

In a randomized, double-blind, multicenter, placebo-controlled trial (Study 4) involving 177 patients, methylphenidate hydrochloride extended-release tablets was demonstrated to be effective in the treatment of ADHD in adolescents aged 13 to 18 years at doses up to 72 mg/day (1.4 mg/kg/day). Of 220 patients who entered an open 4-week titration phase, 177 were titrated to an individualized dose (maximum of 72 mg/day) based on meeting specific improvement criteria on the ADHD Rating Scale and the Global Assessment of Effectiveness with acceptable tolerability. Patients who met these criteria were then randomized to receive either their individualized dose of methylphenidate hydrochloride extended-release tablets (18 – 72 mg/day, n=87) or placebo (n=90) during a two-week double-blind phase. At the end of this phase, mean scores for the investigator rating on the ADHD Rating Scale demonstrated that methylphenidate hydrochloride extended-release tablets was statistically significantly superior to placebo.

14.3 Adults

Two double-blind, placebo-controlled studies were conducted in 627 adults aged 18 to 65 years. The controlled studies compared methylphenidate hydrochloride extendedrelease tablets administered once daily and placebo in a multicenter, parallel-group, 7week dose-titration study (Study 5) (36 to 108 mg/day) and in a multicenter, parallelgroup, 5-week, fixed-dose study (Study 6) (18, 36, and 72 mg/day).

Study 5 demonstrated the effectiveness of methylphenidate hydrochloride extendedrelease tablets in the treatment of ADHD in adults aged 18 to 65 years at doses from 36 mg/day to 108 mg/day based on the change from baseline to final study visit on the Adult ADHD Investigator Rating Scale (AISRS). Of 226 patients who entered the 7-week trial, 110 were randomized to methylphenidate hydrochloride extended-release tablets and 116 were randomized to placebo. Treatment was initiated at 36 mg/day and patients continued with incremental increases of 18 mg/day (36 to 108 mg/day) based on meeting specific improvement criteria with acceptable tolerability. At the final study visit, mean change scores (LS Mean, SEM) for the investigator rating on the AISRS demonstrated that methylphenidate hydrochloride extended-release tablets was statistically significantly superior to placebo.

Study 6 was a multicenter, double-blind, randomized, placebo-controlled, parallel-group, dose-response study (5-week duration) with 3 fixed-dose groups (18, 36, and 72 mg). Patients were randomized to receive methylphenidate hydrochloride extended-release tablets administered at doses of 18 mg (n=101), 36 mg (n=102), 72 mg/day (n=102), or placebo (n=96). All three doses of methylphenidate hydrochloride extended-release tablets were statistically significantly more effective than placebo in improving CAARS (Conners' Adult ADHD Rating Scale) total scores at double-blind end point in adult subjects with ADHD.

15 REFERENCES

American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. 4th ed. Washington, DC: American Psychiatric Association, 1994.

16 HOW SUPPLIED/STORAGE AND HANDLING

Methylphenidate Hydrochloride Extended-Release Tablets, USP are available in 18 mg, 27 mg, 36 mg, and 54 mg dosage strengths.

The 18 mg tablets are light yellow to yellow film coated round cylindrical biconvex tablets printed with 212 in black ink.

NDC 50268-545-12 (5 tablets per card, 4 cards per tablet).

The 27 mg tablets are light pink to pink film coated round cylindrical biconvex tablets printed with 213 in black ink.

NDC 50268-546-12 (5 tablets per card, 4 cards per tablet).

The 36 mg tablets are White to off white film coated round cylindrical biconvex tablets printed with 214 in black ink.

NDC 50268-547-12 (5 tablets per card, 4 cards per tablet).

The 54 mg tablets are light to dark brown film coated round cylindrical biconvex tablets printed with 215 in black ink.

NDC 50268-548-12 (5 tablets per card, 4 cards per tablet).

For Institutional Use Only. Dispensed in Unit Dose Package.

Storage and Handling

Store at 20° to 25°C (68° to 77°F). [see USP Controlled Room Temperature]. Protect from humidity.

17 PATIENT COUNSELING INFORMATION

See Medication Guide

Priapism

Advise patients, caregivers, and family members of the possibility of painful or prolonged penile erections (priapism). **Instruct the patient to seek immediate medical attention in the event of priapism** [see Warnings and Precautions (5.4)].

Circulation Problems in Fingers and Toes [Peripheral Vasculopathy, including Raynaud's Phenomenon]

Instruct patients beginning treatment with methylphenidate hydrochloride extendedrelease tablets about the risk of peripheral vasculopathy, including Raynaud's phenomenon, and associated signs and symptoms: fingers or toes may feel numb, cool, painful, and/or may change color from pale, to blue, to red.

Instruct patients to report to their physician any new numbness, pain, skin color change, or sensitivity to temperature in fingers or toes.

Instruct patients to call their physician immediately with any signs of unexplained wounds appearing on fingers or toes while taking methylphenidate hydrochloride extended-release tablets.

Further clinical evaluation (e.g., rheumatology referral) may be appropriate for certain patients.

General Considerations

Prescribers or other health professionals should inform patients, their families, and their caregivers about the benefits and risks associated with treatment with methylphenidate and should counsel them in its appropriate use. A patient Medication Guide is available for methylphenidate hydrochloride extended-release tablets. The prescriber or health professional should instruct patients, their families, and their caregivers to read the Medication Guide and should assist them in understanding its contents. Patients should be given the opportunity to discuss the contents of the Medication Guide and to obtain answers to any questions they may have. The complete text of the Medication Guide is reprinted at the end of this document.

Administration Instructions

Patients should be informed that methylphenidate hydrochloride extended-release tablets should be swallowed whole with the aid of liquids. Tablets should not be chewed, divided, or crushed. The medication is contained within a nonabsorbable shell designed to release the drug at a controlled rate. The tablet shell, along with insoluble core components, is eliminated from the body; patients should not be concerned if they occasionally notice in their stool something that looks like a tablet.

Driving or Operating Heavy Machinery

Stimulants may impair the ability of the patient to operate potentially hazardous machinery or vehicles. Patients should be cautioned accordingly until they are reasonably certain that methylphenidate hydrochloride extended-release tablets does not adversely affect their ability to engage in such activities.

For more information call 1-855-361-3993.

Manufactured for:

AvKARE

Pulaski, TN 38478

Mfg. Rev. 02/19 AV 04/21 (P) AvPAK

Methylphenidate hydrochloride Extended-release Tablets, USP CII

(meth ' il fen ' i date hye " droe klor ' ide)

Read the Medication Guide that comes with methylphenidate hydrochloride extendedrelease tablets before you or your child starts taking it and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking to your doctor about your or your child's treatment with methylphenidate hydrochloride extended-release tablets. What is the most important information I should know about methylphenidate hydrochloride extended-release tablets?

The following have been reported with use of methylphenidate HCI and other stimulant medicines:

- 1. Heart-related problems:
- sudden death in patients who have heart problems or heart defects
- stroke and heart attack in adults

increased blood pressure and heart rate

Tell your doctor if you or your child has any heart problems, heart defects, high blood pressure, or a family history of these problems.

Your doctor should check you or your child carefully for heart problems before starting methylphenidate hydrochloride extended-release tablets.

Your doctor should check your or your child's blood pressure and heart rate regularly during treatment with methylphenidate hydrochloride extended-release tablets.

Call your doctor right away if you or your child has any signs of heart problems such as chest pain, shortness of breath, or fainting while taking methylphenidate hydrochloride extended-release tablets.

2. Mental (Psychiatric) problems:

All Patients

- new or worse behavior and thought problems
- new or worse bipolar illness
- new or worse aggressive behavior or hostility

Children and Teenagers

 new psychotic symptoms (such as hearing voices, believing things that are not true, are suspicious) or new manic symptoms

Tell your doctor about any mental problems you or your child have, or about a family history of suicide, bipolar illness, or depression.

Call your doctor right away if you or your child has any new or worsening mental symptoms or problems while taking methylphenidate hydrochloride extended-release tablets, especially seeing or hearing things that are not real, believing things that are not real, or are suspicious.

3. Painful and prolonged erections (priapism)

Painful and prolonged erections (priapism) have occurred with methylphenidate. If you or your child develop priapism, seek medical help right away.

Because of the potential for lasting damage, priapism should be evaluated by

a doctor immediately.

4. Circulation problems in fingers and toes

[Peripheral vasculopathy, including Raynaud's phenomenon]:

- fingers or toes may feel numb, cool, painful
- fingers or toes may change color from pale, to blue, to red

Tell your doctor if you have or your child has numbness, pain, skin color change, or sensitivity to temperature in your fingers or toes.

Call your doctor right away if you have or your child has any signs of unexplained wounds appearing on fingers or toes while taking methylphenidate hydrochloride extended-release tablets.

What is methylphenidate hydrochloride extended-release tablets?

Methylphenidate hydrochloride extended-release tablets is a central nervous system stimulant prescription medicine. **It is used for the treatment of attention deficit and hyperactivity disorder(ADHD).** Methylphenidate hydrochloride extended-release tablets may help increase attention and decrease impulsiveness and hyperactivity in patients with ADHD.

Methylphenidate hydrochloride extended-release tablets should be used as a part of a total treatment program for ADHD that may include counseling or other therapies.

Methylphenidate hydrochloride extended-release tablets is a federally controlled substance (CII) because it can be abused or lead to dependence. Keep methylphenidate hydrochloride extended-release tablets in a safe place to prevent misuse and abuse. Selling or giving away Methylphenidate hydrochloride extended-release tablets may harm others, and is against the law.

Tell your doctor if you or your child has (or has a family history of) ever abused or been dependent on alcohol, prescription medicines, or street drugs.

Who should not take methylphenidate hydrochloride extended-release tablets?

Methylphenidate hydrochloride extended-release tablets should not be taken if you or your child:

- is very anxious, tense, or agitated
- has an eye problem called glaucoma
- has tics or Tourette's syndrome, or a family history of Tourette's syndrome. Tics are hard-to-control repeated movements or sounds.
- is taking or has taken within the past 14 days an antidepression medicine called a monoamine oxidase inhibitor or MAOI.

• is allergic to anything in methylphenidate hydrochloride extended-release tablets. See the end of this Medication Guide for a complete list of ingredients.

Methylphenidate hydrochloride extended-release tablets should not be used in children less than 6 years old because it has not been studied in this age group.

Methylphenidate hydrochloride extended-release tablets may not be right for you or your child. Before starting methylphenidate hydrochloride extendedrelease tablets, tell your or your child's doctor about all health conditions (or a family history of) including:

- heart problems, heart defects, or high blood pressure
- mental problems including psychosis, mania, bipolar illness, or depression
- tics or Tourette's syndrome
- seizures or have had an abnormal brain wave test (EEG)
- circulation problems in fingers and toes
- esophagus, stomach, or small or large intestine problems

Tell your doctor if you or your child is pregnant, planning to become pregnant, or breastfeeding.

Can methylphenidate hydrochloride extended-release tablets be taken with other medicines?

Tell your doctor about all of the medicines that you or your child takes including prescription and nonprescription medicines, vitamins, and herbal supplements. Methylphenidate hydrochloride extended-release tablets and some medicines may interact with each other and cause serious side effects. Sometimes the doses of other medicines will need to be adjusted while taking methylphenidate hydrochloride extended-release tablets.

Your doctor will decide whether methylphenidate hydrochloride extended-release tablets can be taken with other medicines.

Especially tell your doctor if you or your child takes:

- antidepression medicines including MAOIs
- seizure medicines
- blood thinner medicines
- blood pressure medicines
- cold or allergy medicines that contain decongestants

Know the medicines that you or your child takes.

Keep a list of your medicines with you to show your doctor and pharmacist.

Do not start any new medicine while taking methylphenidate hydrochloride extended-release tablets without talking to your doctor first.

How should methylphenidate hydrochloride extended-release tablets be taken?

• Take methylphenidate hydrochloride extended-release tablets exactly as prescribed. Your doctor may adjust the dose until it is right for you or your child.

• Do not chew, crush, or divide the tablets.

Swallow methylphenidate hydrochloride extended-release tablets whole with water or other liquids. Tell your doctor if you or your child cannot swallow methylphenidate hydrochloride extended-release tablets whole. A different medicine may need to be prescribed.

• methylphenidate hydrochloride extended-release tablets can be taken with or without food.

• Take methylphenidate hydrochloride extended-release tablets once each day in the morning.

methylphenidate hydrochloride extended-release tablets is an extended-release tablet. It releases medication into your or your child's body throughout the day.

• The methylphenidate hydrochloride extended-release tablets does not dissolve completely in the body after all the medicine has been released. You or your child may sometimes notice the empty tablet in a bowel movement. This is normal.

• From time to time, your doctor may stop methylphenidate hydrochloride extendedrelease tablets treatment for a while to check ADHD symptoms.

• Your doctor may do regular checks of the blood, heart, and blood pressure while taking methylphenidate hydrochloride extended-release tablets. Children should have their height and weight checked often while taking methylphenidate hydrochloride extended-release tablets. Methylphenidate hydrochloride extended-release tablets treatment may be stopped if a problem is found during these checkups.

If you or your child takes too much methylphenidate hydrochloride extended-release tablets or overdoses, call your doctor or poison control center right away, or get emergency treatment.

What are possible side effects of methylphenidate hydrochloride extendedrelease tablets?

See "What is the most important information I should know about methylphenidate hydrochloride extended-release tablets?" for information on reported heart and mental problems.

Other serious side effects include:

- slowing of growth (height and weight) in children
- seizures, mainly in patients with a history of seizures
- eyesight changes or blurred vision

• blockage of the esophagus, stomach, small or large intestine in patients who already have a narrowing in any of these organs

Common side effects include:

- decreased appetite headache
- dry mouth nausea
- trouble sleeping anxiety

- dizziness weight loss
- stomach ache irritability
- increased sweating

Stimulants may impair the ability of you or your child to operate potentially hazardous machinery or vehicles. You or your child should exercise caution until you or your child is reasonably certain that methylphenidate hydrochloride extended-release tablets does not adversely affect your or your child's ability to engage in such activities.

Talk to your doctor if you or your child has side effects that are bothersome or do not go away.

This is not a complete list of possible side effects.

Ask your doctor or pharmacist for more information.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

You may also report side effects to AvKARE at 1-855-361-3993.

How should I store methylphenidate hydrochloride extended-release tablets?

• Store methylphenidate hydrochloride extended-release tablets in a safe place at room temperature, Store at 20° to 25°C (68° to 77°F). Protect from moisture.

• Keep methylphenidate hydrochloride extended-release tabletsand all medicines out of the reach of children.

General information about methylphenidate hydrochloride extended-release tablets

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use methylphenidate hydrochloride extended-release tablets for a condition for which it was not prescribed. Do not give methylphenidate hydrochloride extended-release tablets to other people, even if they have the same condition. It may harm them and it is against the law.

This Medication Guide summarizes the most important information about methylphenidate hydrochloride extended-release tablets. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about methylphenidate hydrochloride extended-release tablets that was written for healthcare professionals. For more information about methylphenidate hydrochloride extended-release tablets call 1-855-361-3993.

What are the ingredients in methylphenidate hydrochloride extended-release tablets?

Active Ingredient: methylphenidate HCl

Inactive Ingredients: Methylphenidate hydrochloride extended-release tablets contains the following inert ingredients and are common to all strengths: butylated hydroxytoluene, cellulose acetate, hypromellose, phosphoric acid, polyethylene glycol, polyethylene oxides, povidone, propylene glycol, sodium chloride, stearic acid, succinic acid, ferric oxide yellow, FD&C Red No 40 and titanium dioxide. The 18 mg tablet strength also contains iron oxide yellow and Polysorbate 80. The 27 mg tablet strength

also contains iron oxide red. The 36 mg tablet strength also contains talc. The 54 mg tablet strength also contains iron oxide yellow, iron oxide red and talc.

Each tablet strength also contains black iron oxide, hypromellose and propylene glycol as imprinting ink.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Manufactured for:


AvKARE

Pulaski, TN 38478

Mfg. Rev. 02/19

AV 04/21 (P)

AvPAK



PHARMACIST: Dispense the enclosed Medication Guide to each patient.

268545

Each tablet contains 18 mg methylphenidate hydrochloride in a controlled-release formulation.

Usual Dosage: Once daily. See package insert for dosage information.

Store at 20° to 25°C (68° to 77°F); [see USP Controlled Room Temperature]. Protect from humidity.

Keep this and all medication out of the reach of children.

Manufactured for: AvKARE Pulaski, TN 38478



Mfg. Rev. 02/19

AV 04/21 (P)

٦





268546

PHARMACIST: Dispense the enclosed Medication Guide to each patient.

Each tablet contains 27 mg methylphenidate hydrochloride in a controlled-release formulation.

Usual Dosage: Once daily. See package insert for dosage information.

Store at 20° to 25°C (68° to 77°F); [see USP Controlled Room Temperature]. Protect from humidity.

Keep this and all medication out of the reach of children.

Manufactured for: AvKARE Pulaski, TN 38478



Mfg. Rev. 02/19

AV 04/21 (P)

1

٦



٦

Hydrochloride Extended-Release Tablets, USP

> 36 mg Rx Only 20 Tablets (4 x 5) Unit Dose



PHARMACIST: Dispense the enclosed Medication Guide to each patient.

Each tablet contains 36 mg methylphenidate hydrochloride in a controlled-release formulation.

Usual Dosage: Once daily. See package insert for dosage information.

Store at 20° to 25°C (68° to 77°F); [see USP Controlled Room Temperature]. Protect from humidity.

Keep this and all medication out of the reach of children.


Manufactured for: AvKARE Pulaski, TN 38478



Mfg. Rev. 02/19

AV 04/21 (P)

.



METHYLPHENIDATE HYDROCHLORIDE

methylphenidate hydrochloride tablet, extended release

Product Information

Draduat Tuna

HUMAN

am Cada (Sauraa)

NDC:50268-

Route of Administration	ctive Moiety Ingredient Name ROCHLORIDE (UNII: 4B350	DEA Sched		CII	
- METHYLPHENIDATE HYDF	Ingredient Name ROCHLORIDE (UNII: 4B3SC	C438HI)	Pasis of		
METHYLPHENIDATE HYDF	Ingredient Name ROCHLORIDE (UNII: 4B3SC	C438HI)	Pasis of		
- METHYLPHENIDATE HYDF	Ingredient Name ROCHLORIDE (UNII: 4B3SC	C438HI)	Rasis of		
METHYLPHENIDATE HYDP	ROCHLORIDE (UNII: 4B3SC	C438HI)	Basis of		
		C438HI)	Dasis Vi	Strength	Strengt
(METHYLPHENIDATE - UNII:2)	07ZZ9QZ49)		METHYLPHEN		18 mg
			HYDROCHLOI	RIDE	
Inactive Ingredients	S				
	Ingredient N	Name		S	trength
BUTYLATED HYDROXYTO	•				
CELLULOSE ACETATE (UN	· ·	,			
HYPROMELLOSES (UNII: 3	-				
PHOSPHORIC ACID (UNII: I					
POLYETHYLENE GLYCOL,	UNSPECIFIED (UNII: 3W)C	QOSDW1A)			
POVIDONE (UNII: FZ989GH	194E)				
PROPYLENE GLYCOL (UNI	I: 6DC9Q167V3)				
SODIUM CHLORIDE (UNII:	451W47IQ8X)				
STEARIC ACID (UNII: 4ELV7	7Z65AP)				
SUCCINIC ACID (UNII: AB6	MNQ6J6L)				
ERROSOFERRIC OXIDE ((UNII: XM0M87F357)				
FERRIC OXIDE YELLOW (U	· /				
TITANIUM DIOXIDE (UNII: 1	UNII: EX438O2MRT)				
	UNII: EX438O2MRT) 15FIX9V2JP)				
POLYSORBATE 80 (UNII: 6	UNII: EX438O2MRT) 15FIX9V2JP) 6OZP39ZG8H)				
POLYSORBATE 80 (UNII: 6	UNII: EX438O2MRT) 15FIX9V2JP) 6OZP39ZG8H)				
POLYSORBATE 80 (UNII: 6 FD&C RED NO. 40 (UNII: W	UNII: EX438O2MRT) 15FIX9V2JP) 6OZ P39Z G8H) NZ B9127XOA)				
POLYSORBATE 80 (UNII: 6 FD&C RED NO. 40 (UNII: W Product Characteris	UNII: EX43802MRT) 15FIX9V2JP) 50ZP39ZG8H) WZB9127XOA) Stics				
POLYSORBATE 80 (UNII: 6 FD&C RED NO. 40 (UNII: W Product Characteris Color yel	UNII: EX438O2MRT) 15FIX9V2JP) 5OZ P39Z G8H) VZ B9127XOA) stics Ilow	Score	5	no sc	
POLYSORBATE 80 (UNII: 6 FD&C RED NO. 40 (UNII: W Product Characteris Color yel Shape RO	UNII: EX43802MRT) 15FIX9V2JP) 50ZP39ZG8H) WZB9127XOA) Stics	Size		12mm	
POLYSORBATE 80 (UNII: 6 FD&C RED NO. 40 (UNII: W Product Characteris Color yel Shape RO Flavor	UNII: EX438O2MRT) 15FIX9V2JP) 5OZ P39Z G8H) VZ B9127XOA) stics Ilow	Size	, nt Code		
POLYSORBATE 80 (UNII: 6 FD&C RED NO. 40 (UNII: W Product Characteris Color yel Shape RO Flavor	UNII: EX438O2MRT) 15FIX9V2JP) 5OZ P39Z G8H) VZ B9127XOA) stics Ilow	Size		12mm	
POLYSORBATE 80 (UNII: 6 FD&C RED NO. 40 (UNII: W Product Characteris Color yel Shape RO Flavor	UNII: EX438O2MRT) 15FIX9V2JP) 5OZ P39Z G8H) VZ B9127XOA) stics Ilow	Size		12mm	
POLYSORBATE 80 (UNII: 6 FD&C RED NO. 40 (UNII: W Product Characteris Color yel Shape RO Flavor Contains	UNII: EX438O2MRT) 15FIX9V2JP) 5OZ P39Z G8H) VZ B9127XOA) stics Ilow	Size		12mm	
POLYSORBATE 80 (UNII: 6 FD&C RED NO. 40 (UNII: W Product Characteris Color yel Shape RO Flavor Contains	UNII: EX438O2MRT) 15FIX9V2JP) 5OZ P39Z G8H) VZ B9127XOA) stics Ilow	Size	nt Code	12mm 212	1
POLYSORBATE 80 (UNII: 6 FD&C RED NO. 40 (UNII: W Product Characteris Color yel Shape RO Flavor Contains S	UNII: EX438O2MRT) 15FIX9V2JP) 5OZ P39Z G8H) VZ B9127XOA) stics Ilow	Size Impri		12mm 212 t Marke	
POLYSORBATE 80 (UNII: 6 FD&C RED NO. 40 (UNII: W Product Characteris Color yel Shape RO Flavor Contains RO Packaging # Item Code	UNII: EX43802MRT) 15FIX9V2JP) 50ZP39ZG8H) VZB9127XOA) stics Ilow DUND (cylindrical) Package Descript	Size Impri	nt Code Marketing Stai	12mm 212 t Marke	ting End
POLYSORBATE 80 (UNII: 6 FD&C RED NO. 40 (UNII: W Product Characteris Color yel Shape RO Flavor Contains RO Packaging # Item Code 1 NDC:50268-545- 20 in 1 NDC:50268-545 1 in 1 R	UNII: EX43802MRT) 15FIX9V2JP) 60ZP39ZG8H) VZB9127XOA) Stics Ilow DUND (cylindrical) Package Descript BOX	Size Impri	nt Code Marketing Star Date	12mm 212 t Marke	ting End
POLYSORBATE 80 (UNII: 6 FD&C RED NO. 40 (UNII: W Product Characteris Color yel Shape RO Flavor Contains RO Packaging # Item Code 1 NDC:50268-545- 20 in 1 NDC:50268-545 1 in 1 R	UNII: EX43802MRT) 15FIX9V2JP) 50ZP39ZG8H) VZB9127XOA) stics Ilow DUND (cylindrical) Package Descript BOX BLISTER PACK; Type 0: Not	Size Impri	nt Code Marketing Star Date	12mm 212 t Marke	ting End
POLYSORBATE 80 (UNII: 6 FD&C RED NO. 40 (UNII: W Product Characteris Color yel Shape RO Flavor RO Contains Item Code I NDC:50268-545- 12 20 in 1 NDC:50268-545- 1 in 1 B	UNII: EX43802MRT) 15FIX9V2JP) 50ZP39ZG8H) VZB9127XOA) stics Ilow DUND (cylindrical) Package Descript BOX BLISTER PACK; Type 0: Not	Size Impri	nt Code Marketing Star Date	12mm 212 t Marke	ting End
PULYSORBATE 80 (UNII: 6 FD&C RED NO. 40 (UNII: W PUDET Characteris Color yel Shape RO Flavor RO Contains Image: Shape W Item Code 1 NDC:50268-545- 12 20 in 1 1 NDC:50268-545- 11 1 in 1 B	UNII: EX43802MRT) 15FIX9V2JP) 60ZP39ZG8H) WZB9127XOA) Stics Ilow DUND (cylindrical) Package Descript BOX BLISTER PACK; Type 0: Not	Size Impri	nt Code Marketing Star Date	12mm 212 t Marke	ting End
POLYSORBATE 80 (UNII: 6 FD&C RED NO. 40 (UNII: W Product Characteris Color yel Shape RO Flavor RO Contains Image: Shape W Item Code 1 NDC:50268-545- 11 20 in 1 NDC:50268-545- 11 1 in 1 B	UNII: EX43802MRT) 15FIX9V2JP) 60ZP39ZG8H) WZB9127XOA) Stics Ilow DUND (cylindrical) Package Descript BOX BLISTER PACK; Type 0: Not	Size Impri	nt Code Marketing Star Date	12mm 212 t Marke	ting End
Shape RO Flavor RO Contains Item Code # Item Code 1 NDC:50268-545- 1 In 1 B NDC:50268-545- In 1 B	UNII: EX43802MRT) 15FIX9V2JP) 60ZP39ZG8H) WZB9127XOA) Stics Ilow DUND (cylindrical) Package Descript BOX BLISTER PACK; Type 0: Not	Size Impri	nt Code Marketing Star Date	12mm 212	ting End

Μ	ETHYLPH	ENIDATE	HYDROCHLO	RIDE				
me	ethylphenidate	hydrochlorid	le tablet, extended	release				
Ρ	roduct Infor	mation						
Р	roduct Type		HUMAN PRESCRIPTION DRUG	Item Coc	de (Soui	ce)	NDC:5026 546(NDC:3	8- 31722-953)
R	oute of Admini	stration	ORAL	DEA Sch	edule		CII	
A	ctive Ingredi	ent/Active	Moiety					
		Ingree	dient Name			Basis of S	Strength	Strength
			ORIDE (UNII: 4B3SC438	3HI)		METHYLPHENID		27 mg
(M	ETHYLPHENIDATE	- UNII:207ZZ90	QZ49)			HYDROCHLORIE	DE	27 mg
L m	activo Ingro	dianta						
In	active Ingre	aients	In our all and Maria					C
			Ingredient Nan	ne				Strength
			(UNII: 1P9D0Z171K)					
	ELLULOSE ACETA							
	IOSPHORIC ACI							
			CIFIED (UNII: 3WQ0SI					
	VIDONE (UNII: F			5112, 19				
	OPYLENE GLYC		Q167V3)					
sc	DIUM CHLORID	E (UNII: 451W47	7IQ8X)					
ST	EARIC ACID (UN	II: 4ELV7Z65AP))					
รบ	ICCINIC ACID (U	NII: AB6MNQ6J6	L)					
FE	RROSOFERRIC	OXIDE (UNII: XM	10M87F357)					
FE	RRIC OXIDE YEL	LOW (UNII: EX	43802MRT)					
	RRIC OXIDE REI							
	TANIUM DIOXIDI		•					
FC	0&C RED NO. 40	(UNII: WZ B912)	7XOA)					
D .	roduct Chara							
				C -				
	olor	pink ROUND (cy	dindrical)		ore		no se	
	аре	ROUND (C)	/iindrical)	Siz		-	12mi	m
	avor			Im	print Co	ae	213	
C	ontains							
P	ackaging							
					Mar	ceting Start	Mark	eting End
#	Item Code	Pa	ckage Description	1	Mari	Date		Date
1	NDC:50268-546- 12	20 in 1 BOX			04/22/2	021		
1	NDC:50268-546-	1 in 1 BLISTER	PACK; Type 0: Not a C	ombination				

Marketing Information							
Application Number or Monograph Citation	Marketing Start Date	Marketing End Date					
ANDA211009	04/22/2021						
	Application Number or Monograph Citation	Application Number or MonographMarketing StartCitationDate					

METHYLPHENIDATE HYDROCHLORIDE

methylphenidate hydrochloride tablet, extended release

Product Informati	ion					
Product Type		HUMAN PRESCRIPTION DRUG	Item Code (Source)			- L722-954)
Route of Administrat	ion	ORAL	DEA Schedule		CII	
Active Ingredient//	Active	Moiety				
	Ingrea	dient Name		Basis of S	Strength	Strengt
METHYLPHENIDATE HYL (METHYLPHENIDATE - UNII:			3HI)	METHYLPHENID HYDROCHLORII		36 mg
Inactive Ingredien	ts					
		Ingredient Nan	ne		S	trength
BUTYLATED HYDROXYT	OLUENE	(UNII: 1P9D0Z171K)				
CELLULOSE ACETATE (U	JNII: 3J2PC)7GVB6)				
HYPROMELLOSES (UNII:	3NXW29V	3WO)				
PHOSPHORIC ACID (UNII	: E4GA88	84NN)				
POLYETHYLENE GLYCO	L, UNSPE	CIFIED (UNII: 3WJQ051	OW1A)			
POVIDONE (UNII: FZ989G	•					
PROPYLENE GLYCOL (UN						
SODIUM CHLORIDE (UNI						
STEARIC ACID (UNII: 4EL)						
SUCCINIC ACID (UNII: AB	-					
FERROSOFERRIC OXIDE						
FERRIC OXIDE YELLOW	(UNII: EX4	13802MRT)				
TALC (UNII: 7SEV7J4R1U)	. 155000					
		-				
FD&C RED NO. 40 (UNII:	MY R917					
Product Character						
	vhite		Score		no sco	
-	OUND (cy	lindrical)	Size		15mm	l
Flavor			Imprint Co	de	214	
Contains						

Da								
ГС	ackaging							
#	Item Code		ckage Description	ı	Marl	ceting Start Date	Mar	keting End Date
1	NDC:50268-547-	20 in 1 BOX			04/22/2	021		
1		L in 1 BLISTER Product	PACK; Type 0: Not a C	ombination				
Μ	arketing I	nformat	ion					
	Marketing Category	Applicat	tion Number or Mo Citation	nograph	Ma	rketing Start Date	Ма	rketing End Date
AN	DA	ANDA211009	9		04/22/	2021		
ne	thylphenidate	hydrochlorid	e tablet, extended	release				
Pr	roduct Inform	nation						
Pr	oduct Type		HUMAN PRESCRIPTION DRUG	ltem Code	e (Soui	ce)	268- 2:31722-955)	
Rc	oute of Adminis	tration	ORAL	DEA Scheo	dule CII			
Ac	tive Ingredie	ent/Active	Moiety					
		Ingree	dient Name			Basis of S	strengt	h Strengt
	THYLPHENIDATE THYLPHENIDATE -		DRIDE (UNII: 4B3SC438 (Z49)		METHYLPHENID HYDROCHLORIE	54 mg		
In	active Ingred	dients						
			Ingredient Nan	ne				Strength
			(UNII: 1P9D0Z171K)					
СЕ	LLULOSE ACETA							
	PROMELLOSES (
		(UNII: E4GA88	84NN)					
PH								
PH PO	LYETHYLENE GL		CIFIED (UNII: 3WJQ0SI	DW1A)				
PH PO PO	LYETHYLENE GL VIDONE (UNII: FZ	989GH94E)		DWIA)				
PH PO PO PR	LYETHYLENE GL VIDONE (UNII: FZ OPYLENE GLYCO	989GH94E) • UNII: 6DC90	Q167V3)	DW1A)				
PH PO PO PR SO	LYETHYLENE GL VIDONE (UNII: FZ OPYLENE GLYCO DIUM CHLORIDE	989GH94E) DL (UNII: 6DC90 (UNII: 451W47	Q167V3) IQ8X)	DWIA)				
PH PO PR SO ST	LYETHYLENE GL VIDONE (UNII: FZ OPYLENE GLYCO DIUM CHLORIDE EARIC ACID (UNII	989GH94E) L (UNII: 6DC90 (UNII: 451W47 : 4ELV7Z65AP)	Q167V3) IQ8X)	DWIA)				
PH PO PR SO ST SU	LYETHYLENE GL VIDONE (UNII: FZ OPYLENE GLYCO DIUM CHLORIDE EARIC ACID (UNII CCINIC ACID (UN	989GH94E) L (UNII: 6DC90 (UNII: 451W47 : 4ELV7Z65AP) II: AB6MNQ6J6	Q167V3) (Q8X) L)	DWIA)				
PH PO PR SO ST SU FE	LYETHYLENE GL VIDONE (UNII: FZ OPYLENE GLYCO DIUM CHLORIDE EARIC ACID (UNII CCINIC ACID (UN RROSOFERRIC O	989GH94E) L (UNII: 6DC90 (UNII: 451W47 : 4ELV7Z65AP) II: AB6MNQ6J6 XIDE (UNII: XM	Q167V3) IQ8X) L) IOM87F357)	DWIA)				
PH PO PR SO ST SU FE	LYETHYLENE GL VIDONE (UNII: FZ OPYLENE GLYCO DIUM CHLORIDE EARIC ACID (UNII CCINIC ACID (UN	989GH94E) L (UNII: 6DC90 (UNII: 451W47 : 4ELV7Z 65AP) II: AB6MNQ6J6 XIDE (UNII: XM LOW (UNII: EX4	2167V3) (Q8X) L) 10M87F357) 438O2MRT)	DWIA)				

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

		•	• •									
FC	0&C RED NO. 40	(UNII	: WZB9127XOA)									
-												
Product Characteristics												
Co	olor		brown S	cor	e		no score					
Shape			ROUND (cylindrical)		e.		15mm					
Fla	avor		In	mpr	int Code		215					
Сс	ontains											
Packaging												
	ackaying											
#	Item Code		Package Description		Marketing Start Date	M	larketing End Date					
#		20 in				Μ						
# 1	Item Code NDC:50268-548- 12 NDC:50268-548-		1 BOX 1 BLISTER PACK; Type 0: Not a Combinatio		Date	M						
#	Item Code NDC:50268-548- 12 NDC:50268-548-	1 in 1	1 BOX 1 BLISTER PACK; Type 0: Not a Combinatio		Date	M						
#	Item Code NDC:50268-548- 12 NDC:50268-548-	1 in 1	1 BOX 1 BLISTER PACK; Type 0: Not a Combinatio		Date	M						
# 1 1	Item Code NDC:50268-548- 12 NDC:50268-548-	1 in Prod	1 BOX 1 BLISTER PACK; Type 0: Not a Combinatio uct		Date	M						
# 1 1	Item Code NDC:50268-548- 12 NDC:50268-548- 11	1 in Prod	1 BOX 1 BLISTER PACK; Type 0: Not a Combinatio uct	on	Date							
# 1 1	Item Code NDC:50268-548- 12 NDC:50268-548- 11	1 in Prod	a 1 BOX 1 BLISTER PACK; Type 0: Not a Combinatio uct Ormation Application Number or Monograpi	on	Date 04/22/2021 Marketing Start		Date Marketing End					

Labeler - AvPAK (832926666)

Revised: 1/2024

AvPAK