

# GADOBUTROL- gadobutrol injection

## Fresenius Kabi USA, LLC

### HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use GADOBUTROL INJECTION safely and effectively. See full prescribing information for GADOBUTROL INJECTION.

GADOBUTROL injection, for intravenous use  
Initial U.S. Approval: 2011

IMAGING BULK PACKAGE  
NOT FOR DIRECT INFUSION

#### **WARNING: RISK ASSOCIATED WITH INTRATHECAL USE and NEPHROGENIC SYSTEMIC FIBROSIS**

*See full prescribing information for complete boxed warning*

- Intrathecal administration of gadolinium-based contrast agents (GBCAs) can cause serious adverse reactions including death, coma, encephalopathy, and seizures. Gadobutrol injection is not approved for intrathecal use (5.1)
- GBCAs increase the risk for nephrogenic systemic fibrosis (NSF) among patients with impaired elimination of the drugs. Avoid use of gadobutrol injection in these patients unless the diagnostic information is essential and not available with non-contrasted MRI or other modalities.

The risk for NSF appears highest among patients with:

- Chronic, severe kidney disease ( $\text{GFR} < 30 \text{ mL/min/1.73m}^2$ ), or
- Acute kidney injury.

Screen patients for acute kidney injury and other conditions that may reduce renal function. For patients at risk for chronically reduced renal function (for example, age  $> 60$  years, hypertension or diabetes), estimate the glomerular filtration rate (GFR) through laboratory testing (5.2).

#### **RECENT MAJOR CHANGES**

Boxed Warning

1/2024

Warnings and Precautions, Risk Associated with Intrathecal Use (5.1)

1/2024

#### **INDICATIONS AND USAGE**

Gadobutrol injection is a gadolinium-based contrast agent indicated for use with magnetic resonance imaging (MRI):

- To detect and visualize areas with disrupted blood brain barrier and/or abnormal vascularity of the central nervous system in adult and pediatric patients (including term neonates) (1.1)
- To assess the presence and extent of malignant breast disease in adult patients (1.2)
- To evaluate known or suspected supra-aortic or renal artery disease in adult and pediatric patients (including term neonates) (1.3)
- To assess myocardial perfusion (stress, rest) and late gadolinium enhancement in adult patients with known or suspected coronary artery disease (CAD). (1.4).

#### **DOSAGE AND ADMINISTRATION**

- Recommended dose for adults and pediatric patients (including term neonates) is 0.1 mL/kg body weight (2.1)
- Administer as an intravenous bolus injection (2.2)
- Follow injection with a normal saline flush (2.2)

#### **DOSAGE FORMS AND STRENGTHS**

Gadobutrol injection contains 604.72 mg gadobutrol/mL (equivalent to 1 mmol gadobutrol/mL) (3)

#### **CONTRAINDICATIONS**

History of severe hypersensitivity reaction to gadobutrol (4)

#### **WARNINGS AND PRECAUTIONS**

- Anaphylactic and other hypersensitivity reactions with cardiovascular, respiratory or cutaneous manifestations, ranging from mild to severe, including death, have occurred. Monitor patients closely during and after administration of gadobutrol. (5.3)
- Gadolinium is retained for months or years in brain, bone, and other organs. (5.4)

#### **ADVERSE REACTIONS**

Most common adverse reactions (incidence  $\geq 0.5\%$ ) are headache, nausea, and dizziness (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Fresenius Kabi USA, LLC at 1-800-551-7176 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch)**

#### **USE IN SPECIFIC POPULATIONS**

Pregnancy: Use only if imaging is essential during pregnancy and cannot be delayed. (8.1)

**See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.**

**Revised: 9/2024**

---

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

### **WARNING: RISK ASSOCIATED WITH INTRATHECAL USE and NEPHROGENIC SYSTEMIC FIBROSIS**

#### **1 INDICATIONS AND USAGE**

- 1.1 Magnetic Resonance Imaging (MRI) of the Central Nervous System (CNS)
- 1.2 MRI of the Breast
- 1.3 Magnetic Resonance Angiography (MRA)
- 1.4 Cardiac MRI

#### **2 DOSAGE AND ADMINISTRATION**

- 2.1 Recommended Dose
- 2.2 Administration Guidelines
- 2.3 Drug Handling
- 2.4 Imaging Bulk Package Preparation Instructions

#### **3 DOSAGE FORMS AND STRENGTHS**

#### **4 CONTRAINDICATIONS**

#### **5 WARNINGS AND PRECAUTIONS**

- 5.1 Risk Associated with Intrathecal Use
- 5.2 Nephrogenic Systemic Fibrosis
- 5.3 Hypersensitivity Reactions
- 5.4 Gadolinium Retention
- 5.5 Acute Kidney Injury
- 5.6 Extravasation and Injection Site Reactions
- 5.7 Overestimation of Extent of Malignant Disease in MRI of the Breast
- 5.8 Low Sensitivity for Significant Arterial Stenosis

#### **6 ADVERSE REACTIONS**

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

#### **8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment

**10 OVERDOSAGE****11 DESCRIPTION****12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

**13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

13.2 Animal Toxicology and/or Pharmacology

**14 CLINICAL STUDIES**

14.1 MRI of the CNS

14.2 MRI of the Breast

14.3 MRA

14.4 Cardiac MRI

**16 HOW SUPPLIED/STORAGE AND HANDLING**

16.1 How Supplied

16.2 Storage and Handling

**17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

**FULL PRESCRIBING INFORMATION**

## **WARNING: RISK ASSOCIATED WITH INTRATHECAL USE and NEPHROGENIC SYSTEMIC FIBROSIS**

### **Risk Associated with Intrathecal Use**

Intrathecal administration of gadolinium-based contrast agents (GBCAs) can cause serious adverse reactions including death, coma, encephalopathy, and seizures. Gadobutrol injection is not approved for intrathecal use [see Warnings and Precautions (5.1)].

### **Nephrogenic Systemic Fibrosis**

GBCAs increase the risk for nephrogenic systemic fibrosis (NSF) among patients with impaired elimination of drugs. Avoid use of gadobutrol injection in these patients unless the diagnostic information is essential and not available with non-contrasted MRI or other modalities. NSF may result in fatal or debilitating fibrosis affecting the skin, muscle and internal organs.

- The risk for NSF appears highest among patients with:
  - Chronic, severe kidney disease (GFR < 30 mL/min/1.73m<sup>2</sup>), or
  - Acute kidney injury.
- Screen patients for acute kidney injury and other conditions that may reduce renal function. For patients at risk for chronically reduced renal function (for example, age > 60 years, hypertension or diabetes), estimate the glomerular filtration rate (GFR) through laboratory testing.
- For patients at highest risk for NSF, do not exceed the recommended gadobutrol dose and allow a sufficient period of time for elimination of the drug from the body prior to any re-administration [see Warnings and Precautions (5.2)].

## **1 INDICATIONS AND USAGE**

### **1.1 Magnetic Resonance Imaging (MRI) of the Central Nervous System (CNS)**

Gadobutrol injection is indicated for use with magnetic resonance imaging (MRI) in adult and pediatric patients, including term neonates, to detect and visualize areas with disrupted blood brain barrier and/or abnormal vascularity of the central nervous system.

### **1.2 MRI of the Breast**

Gadobutrol injection is indicated for use with MRI in adult patients to assess the presence and extent of malignant breast disease.

### **1.3 Magnetic Resonance Angiography (MRA)**

Gadobutrol injection is indicated for use in magnetic resonance angiography (MRA) in adult and pediatric patients, including term neonates, to evaluate known or suspected supra-aortic or renal artery disease.

### **1.4 Cardiac MRI**

Gadobutrol injection is indicated for use in cardiac MRI (CMRI) to assess myocardial perfusion (stress, rest) and late gadolinium enhancement in adult patients with known or suspected coronary artery disease (CAD).

## 2 DOSAGE AND ADMINISTRATION

### 2.1 Recommended Dose

The recommended dose of gadobutrol injection for adult and pediatric patients (including term neonates) is 0.1 mL/kg body weight (0.1 mmol/kg). Refer to Table 1 to determine the volume to be administered.

**Table 1: Volume of Gadobutrol Injection by Body Weight**

Body Weight (kg)	Volume to be Administered (mL)
2.5	0.25
5	0.5
10	1
15	1.5
20	2
25	2.5
30	3
35	3.5
40	4
45	4.5
50	5
60	6
70	7
80	8
90	9
100	10
110	11
120	12
130	13
140	14

\*For *Cardiac MRI*, the dose is divided into 2 separate, equal injections

### 2.2 Administration Guidelines

Gadobutrol injection is formulated at a higher concentration (1 mmol/mL) compared to certain other gadolinium based contrast agents, resulting in a lower volume of administration. Use Table 1 to determine the volume to be administered.

Use sterile technique when preparing and administering gadobutrol injection.

#### *MRI of the Central Nervous System*

- Administer gadobutrol injection as an intravenous injection, manually or by power injector, at a flow rate of approximately 2 mL/second.

- Follow gadobutrol injection with flush of 0.9% Sodium Chloride Injection, USP to ensure complete administration of the contrast.
- Post contrast MRI can commence immediately following contrast administration.

#### *MRI of the Breast*

- Administer gadobutrol injection as an intravenous bolus by power injector, followed by a flush of 0.9% Sodium Chloride Injection, USP to ensure complete administration of the contrast.
- Start image acquisition following contrast administration and then repeat sequentially to determine peak intensity and wash-out.

#### *MR Angiography*

Image acquisition should coincide with peak arterial concentration, which varies among patients.

#### Adults

- Administer gadobutrol injection by power injector, at a flow rate of approximately 1.5 mL/second, followed by a 30 mL flush of 0.9% Sodium Chloride Injection, USP at the same rate to ensure complete administration of the contrast.

#### Pediatric patients

- Administer gadobutrol injection by power injector or manually, followed by a flush of 0.9% Sodium Chloride Injection, USP to ensure complete administration of the contrast.

#### *Cardiac MRI*

- Administer gadobutrol injection through a separate intravenous line in the contralateral arm if concomitantly providing a continuous infusion of a pharmacologic stress agent.
- Administer gadobutrol injection as two (2) separate bolus injections: 0.05 mL/kg (0.05 mmol/kg) body weight at peak pharmacologic stress followed by 0.05 mL/kg (0.05 mmol/kg) body weight at rest.
- Administer gadobutrol injection via a power injector at a flow rate of approximately 4 mL/second and follow each injection with a flush of 20 mL of 0.9% Sodium Chloride Injection, USP at the same flow rate.

### **2.3 Drug Handling**

- Visually inspect gadobutrol injection for particulate matter and discoloration prior to administration. Do not use the solution if it is discolored, if particulate matter is present or if the container appears damaged.
- Do not mix gadobutrol injection with other medications and do not administer gadobutrol injection in the same intravenous line simultaneously with other medications because of the potential for chemical incompatibility.
- Instructions of the device manufacturer must be followed.

### **2.4 Imaging Bulk Package Preparation Instructions**

Gadobutrol injection Imaging Bulk Package (IBP) is a container of a sterile preparation for parenteral use that contains many single doses of gadobutrol for use with a medical imaging device. Gadobutrol injection Imaging Bulk Package is for intravenous use and not for direct infusion. Gadobutrol injection Imaging Bulk Package is for use only with an

automated contrast injection system, contrast management system, or contrast media transfer set approved or cleared for use with this contrast agent in this Imaging Bulk Package. Please see drug and device labeling for information on devices indicated for use with this Imaging Bulk Package and techniques to help assure safe use.

1. The Gadobutrol injection Imaging Bulk Package is to be used only in a room designated for radiological procedures that involve intravascular administration of a contrast agent.
2. Utilize aseptic technique for penetrating the container closure of the Gadobutrol injection Imaging Bulk Package and transferring Gadobutrol injection.
3. The container closure must be penetrated only one time with a suitable sterile component of the automated contrast injection system, contrast management system, or contrast media transfer set (e.g., transfer spike) approved or cleared for use with this contrast agent in this Imaging Bulk Package.
4. Once the Gadobutrol injection Imaging Bulk Package is punctured, do not remove it from the work area during the entire period of use. Storage temperature of Gadobutrol injection Imaging Bulk Package after the closure has been entered is 20°C to 25°C (68°F to 77°F).
5. A maximum use time of 24 hours from initial puncture is permitted to complete fluid transfer. Discard any unused Gadobutrol injection 24 hours after initial puncture of the Imaging Bulk Package.
6. After the container closure is punctured, if the integrity of the Imaging Bulk Package and the delivery system cannot be assured through direct continuous supervision, the Imaging Bulk Package and all associated disposables for the automated contrast injection system, contrast management system, or contrast media transfer set (e.g., transfer spike) should be discarded.

### **3 DOSAGE FORMS AND STRENGTHS**

Gadobutrol injection is a sterile, clear, and colorless to pale yellow solution for injection containing 604.72 mg gadobutrol per mL (equivalent to 1 mmol gadobutrol/mL).

### **4 CONTRAINDICATIONS**

Gadobutrol injection is contraindicated in patients with history of severe hypersensitivity reactions to gadobutrol.

### **5 WARNINGS AND PRECAUTIONS**

#### **5.1 Risk Associated with Intrathecal Use**

Intrathecal administration of GBCAs can cause serious adverse reactions including death, coma, encephalopathy, and seizures. The safety and effectiveness of Gadobutrol injection have not been established with intrathecal use. Gadobutrol injection is not approved for intrathecal use [*see Dosage and Administration (2.2)*].

#### **5.2 Nephrogenic Systemic Fibrosis**

GBCAs increase the risk for nephrogenic systemic fibrosis (NSF) among patients with impaired elimination of the drugs. Avoid use of gadobutrol injection among these

patients unless the diagnostic information is essential and not available with non-contrast MRI or other modalities. The GBCA-associated NSF risk appears highest for patients with chronic, severe kidney disease (GFR < 30 mL/min/1.73m<sup>2</sup>) as well as patients with acute kidney injury. The risk appears lower for patients with chronic, moderate kidney disease (GFR 30 to 59 mL/min/1.73m<sup>2</sup>) and little, if any, for patients with chronic, mild kidney disease (GFR 60 to 89 mL/min/1.73m<sup>2</sup>). NSF may result in fatal or debilitating fibrosis affecting the skin, muscle and internal organs. Report any diagnosis of NSF following gadobutrol administration to Fresenius Kabi USA, LLC at 1-800-551-7176 or FDA (1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch)).

Screen patients for acute kidney injury and other conditions that may reduce renal function. Features of acute kidney injury consist of rapid (over hours to days) and usually reversible decrease in kidney function, commonly in the setting of surgery, severe infection, injury or drug-induced kidney toxicity. Serum creatinine levels and estimated GFR may not reliably assess renal function in the setting of acute kidney injury. For patients at risk for chronically reduced renal function (for example, age > 60 years, diabetes mellitus or chronic hypertension), estimate the GFR through laboratory testing.

Among the factors that may increase the risk for NSF are repeated or higher than recommended doses of a GBCA and degree of renal impairment at the time of exposure. Record the specific GBCA and the dose administered to a patient. For patients at highest risk for NSF, do not exceed the recommended gadobutrol dose and allow a sufficient period of time for elimination of the drug prior to re-administration. For patients receiving hemodialysis, consider the prompt initiation of hemodialysis following the administration of a GBCA in order to enhance the contrast agent's elimination [see *Use in Specific Populations (8.6)* and *Clinical Pharmacology (12.3)*]. The usefulness of hemodialysis in the prevention of NSF is unknown [see *Clinical Pharmacology (12.3)*].

### **5.3 Hypersensitivity Reactions**

Anaphylactic and other hypersensitivity reactions with cardiovascular, respiratory or cutaneous manifestations, ranging from mild to severe, including death, have uncommonly occurred following gadobutrol administration [see *Adverse Reactions (6)*].

- Before gadobutrol administration, assess all patients for any history of a reaction to contrast media, bronchial asthma and/or allergic disorders. These patients may have an increased risk for a hypersensitivity reaction to gadobutrol.
- Administer gadobutrol only in situations where trained personnel and therapies are promptly available for the treatment of hypersensitivity reactions, including personnel trained in resuscitation.

Most hypersensitivity reactions to gadobutrol have occurred within half an hour after administration. Delayed reactions can occur up to several days after administration. Observe patients for signs and symptoms of hypersensitivity reactions during and following gadobutrol administration.

### **5.4 Gadolinium Retention**

Gadolinium is retained for months or years in several organs. The highest concentrations (nanomoles per gram of tissue) have been identified in the bone, followed by other organs (for example, brain, skin, kidney, liver, and spleen). The duration of retention also varies by tissue and is longest in bone. Linear GBCAs cause



more retention than macrocyclic GBCAs. At equivalent doses, gadolinium retention varies among the linear agents with Omniscan (gadodiamide) and Optimark (gadoversetamide) causing greater retention than other linear agents [Eovist (gadoxetate disodium), Magnevist (gadopentetate dimeglumine), MultiHance (gadobenate dimeglumine)]. Retention is lowest and similar among the macrocyclic GBCAs [Dotarem (gadoterate meglumine), Gadobutrol injection (gadobutrol), ProHance (gadoteridol)].

Consequences of gadolinium retention in the brain have not been established. Pathologic and clinical consequences of GBCA administration and retention in skin and other organs have been established in patients with impaired renal function [see *Warnings and Precautions* (5.2)]. There are rare reports of pathologic skin changes in patients with normal renal function. Adverse events involving multiple organ systems have been reported in patients with normal renal function without an established causal link to gadolinium retention [see *Adverse Reactions* (6.2)].

While clinical consequences of gadolinium retention have not been established in patients with normal renal function, certain patients might be at higher risk. These include patients requiring multiple lifetime doses, pregnant and pediatric patients, and patients with inflammatory conditions. Consider the retention characteristics of the agent when choosing a GBCA for these patients. Minimize repetitive GBCA imaging studies particularly closely spaced studies, when possible.

### **5.5 Acute Kidney Injury**

In patients with chronic renal impairment, acute kidney injury sometimes requiring dialysis has been observed with the use of some GBCAs. Do not exceed the recommended dose; the risk of acute kidney injury may increase with higher than recommended doses.

### **5.6 Extravasation and Injection Site Reactions**

Ensure catheter and venous patency before the injection of gadobutrol. Extravasation into tissues during gadobutrol administration may result in moderate irritation [see *Nonclinical Toxicology* (13.2)].

### **5.7 Overestimation of Extent of Malignant Disease in MRI of the Breast**

Gadobutrol MRI of the breast overestimated the histologically confirmed extent of malignancy in the diseased breast in up to 50% of the patients [see *Clinical Studies* (14.2)].

### **5.8 Low Sensitivity for Significant Arterial Stenosis**

The performance of gadobutrol MRA for detecting arterial segments with significant stenosis (>50% renal, >70% supra-aortic) has not been shown to exceed 55%. Therefore, a negative MRA study alone should not be used to rule out significant stenosis [see *Clinical Studies* (14.3)].

## **6 ADVERSE REACTIONS**

The following serious adverse reactions are discussed elsewhere in labeling:

- Nephrogenic Systemic Fibrosis (NSF) [see *Boxed Warning and Warnings and Precautions* (5.2)].
- Hypersensitivity reactions [see *Contraindications* (4) and *Warnings and Precautions* (5.3)].

## 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

The adverse reactions described in this section reflect gadobutrol exposure in 7713 subjects (including 184 pediatric patients, ages 0 to 17 years) with the majority receiving the recommended dose. Approximately 52% of the subjects were male and the ethnic distribution was 62% Caucasian, 28% Asian, 5% Hispanic, 2.5% Black, and 2.5% patients of other ethnic groups. The average age was 56 years (range from 1 week to 93 years).

Overall, approximately 4% of subjects reported one or more adverse reactions during a follow-up period that ranged from 24 hours to 7 days after gadobutrol administration.

Adverse reactions associated with the use of gadobutrol were usually mild to moderate in severity and transient in nature.

Table 2 lists adverse reactions that occurred in  $\geq 0.1\%$  subjects who received gadobutrol.

**Table 2: Adverse Reactions**

<b>Reaction</b>	<b>Rate (%) n=7713</b>
Headache	1.7
Nausea	1.2
Dizziness	0.5
Dysgeusia	0.4
Feeling Hot	0.4
Injection site reactions	0.4
Vomiting	0.4
Rash (includes generalized, macular, papular, pruritic)	0.3
Erythema	0.2
Paresthesia	0.2
Pruritus (includes generalized)	0.2
Dyspnea	0.1
Urticaria	0.1

Adverse reactions that occurred with a frequency of  $< 0.1\%$  in subjects who received gadobutrol include: hypersensitivity/anaphylactic reaction, loss of consciousness, convulsion, parosmia, tachycardia, palpitation, dry mouth, malaise and feeling cold.

## 6.2 Postmarketing Experience

The following additional adverse reactions have been reported during postmarketing use of gadobutrol. Because these reactions are reported voluntarily from a population of uncertain size, it is not possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

- Cardiac arrest
- Nephrogenic Systemic Fibrosis (NSF)
- Hypersensitivity reactions (anaphylactic shock, circulatory collapse, respiratory arrest, pulmonary edema, bronchospasm, cyanosis, oropharyngeal swelling, laryngeal edema, blood pressure increased, chest pain, angioedema, conjunctivitis, hyperhidrosis, cough, sneezing, burning sensation, and pallor) [*see Warnings and Precautions (5.3)*].
- General Disorders and Administration Site Conditions: Adverse events with variable onset and duration have been reported after GBCA administration [*see Warnings and Precautions (5.4)*]. These include fatigue, asthenia, pain syndromes, and heterogeneous clusters of symptoms in the neurological, cutaneous, and musculoskeletal systems.
- Skin: Gadolinium associated plaques
- Gastrointestinal Disorders: Acute pancreatitis with onset within 48 hours after GBCA administration

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### *Risk Summary*

GBCAs cross the placenta and result in fetal exposure and gadolinium retention. The human data on the association between GBCAs and adverse fetal outcomes are limited and inconclusive (*see Data*). In animal reproduction studies, although teratogenicity was not observed, embryoletality was observed in monkeys, rabbits and rats receiving intravenous gadobutrol during organogenesis at doses 8 times and above the recommended human dose. Retardation of embryonal development was observed in rabbits and rats receiving intravenous gadobutrol during organogenesis at doses 8 and 12 times, respectively, the recommended human dose (*see Data*). Because of the potential risks of gadolinium to the fetus, use gadobutrol only if imaging is essential during pregnancy and cannot be delayed.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and is 15% to 20%, respectively.

#### *Data*

##### Human Data.

Contrast enhancement is visualized in the placenta and fetal tissues after maternal GBCA administration.

Cohort studies and case reports on exposure to GBCAs during pregnancy have not reported a clear association between GBCAs and adverse effects in the exposed neonates. However, a retrospective cohort study, comparing pregnant women who had

a GBCA MRI to pregnant women who did not have an MRI, reported a higher occurrence of stillbirths and neonatal deaths in the group receiving GBCA MRI. Limitations of this study include a lack of comparison with non-contrast MRI and lack of information about the maternal indication for MRI. Overall, these data preclude a reliable evaluation of the potential risk of adverse fetal outcomes with the use of GBCAs in pregnancy.

### Animal Data

#### Gadolinium Retention

GBCAs administered to pregnant non-human primates (0.1 mmol/kg on gestational days 85 and 135) result in measurable gadolinium concentration in the offspring in bone, brain, skin, liver, kidney, and spleen for at least 7 months. GBCAs administered to pregnant mice (2 mmol/kg daily on gestational days 16 through 19) result in measurable gadolinium concentrations in the pups in bone, brain, kidney, liver, blood, muscle, and spleen at one month postnatal age.

#### Reproductive Toxicology

Embryo lethality was observed when gadobutrol was administered intravenously to monkeys during organogenesis at doses 8 times the recommended single human dose (based on body surface area); gadobutrol was not maternally toxic or teratogenic at this dose. Embryo lethality and retardation of embryonal development also occurred in pregnant rats receiving maternally toxic doses of gadobutrol ( $\geq 7.5$  mmol/kg body weight; equivalent to 12 times the human dose based on body surface area) and in pregnant rabbits ( $\geq 2.5$  mmol/kg body weight; equivalent to 8 times the recommended human dose based on body surface area). In rabbits, this finding occurred without evidence of pronounced maternal toxicity and with minimal placental transfer (0.01% of the administered dose detected in the fetuses).

Because pregnant animals received repeated daily doses of gadobutrol, their overall exposure was significantly higher than that achieved with the standard single dose administered to humans.

## **8.2 Lactation**

### *Risk Summary*

There are no data on the presence of gadobutrol in human milk, the effects on the breastfed infant, or the effects on milk production. However, published lactation data on other GBCAs indicate that 0.01% to 0.04% of the maternal gadolinium dose is present in breast milk and there is limited GBCA gastrointestinal absorption in the breast-fed infant. Gadobutrol is present in rat milk (*see Data*). The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for gadobutrol and any potential adverse effects on the breastfed infant from gadobutrol or from the underlying maternal condition.

### *Data*

In lactating rats receiving 0.5 mmol/kg of intravenous [ $^{153}\text{Gd}$ ]-gadobutrol, 0.01% of the total administered radioactivity was transferred to the pup via maternal milk within 3 hours after administration, and the gastrointestinal absorption is poor (approximately 5% of the dose orally administered was excreted in the urine).

## **8.4 Pediatric Use**

The safety and effectiveness of gadobutrol have been established in pediatric patients, including term neonates, for use with MRI to detect and visualize areas with disrupted blood brain barrier and/or abnormal vascularity of the central nervous system and for use in MRA to evaluate known or suspected supra-aortic or renal artery disease. Use of gadobutrol in these indications is supported by adequate and well-controlled studies in adults and supportive imaging data in two studies in 135 patients 2 to less than 18 years of age and 44 patients less than 2 years of age with CNS and non-CNS lesions, and pharmacokinetic data in 130 patients 2 to less than 18 years of age and 43 patients less than 2 years of age, including term neonates [see *Clinical Pharmacology* (12.3) and *Clinical Studies* (14.1)]. The frequency, type, and severity of adverse reactions in pediatric patients were similar to adverse reactions in adults [see *Adverse Reactions* (6.1)]. No dose adjustment according to age is necessary in pediatric patients [see *Dosage and Administration* (2.1), *Clinical Pharmacology* (12.3), and *Clinical Studies* (14.1)]. The safety and effectiveness of gadobutrol have not been established in preterm neonates for any indication or in pediatric patients of any age for use with MRI to assess the presence and extent of malignant breast disease, or for use in CMRI to assess myocardial perfusion (stress, rest) and late gadolinium enhancement in patients with known or suspected coronary artery disease (CAD).

#### *NSF Risk*

No case of NSF associated with gadobutrol or any other GBCA has been identified in pediatric patients ages 6 years and younger. Pharmacokinetic studies suggest that clearance of gadobutrol is similar in pediatric patients and adults, including pediatric patients age younger than 2 years. No increased risk factor for NSF has been identified in juvenile animal studies of gadobutrol. Normal estimated GFR (eGFR) is around 30 mL/min/1.73m<sup>2</sup> at birth and increases to mature levels around 1 year of age, reflecting growth in both glomerular function and relative body surface area. Clinical studies in pediatric patients younger than 1 year of age have been conducted in patients with the following minimum eGFR: 31 mL/min/1.73m<sup>2</sup> (age 2 to 7 days), 38 mL/min/1.73m<sup>2</sup> (age 8 to 28 days), 62 mL/min/1.73m<sup>2</sup> (age 1 to 6 months), and 83 mL/min/1.73m<sup>2</sup> (age 6 to 12 months).

#### *Juvenile Animal Data*

Single and repeat-dose toxicity studies in neonatal and juvenile rats did not reveal findings suggestive of a specific risk for use in pediatric patients including term neonates and infants.

### **8.5 Geriatric Use**

In clinical studies of gadobutrol, 1,377 patients were 65 years of age and over, while 104 patients were 80 years of age and over. No overall differences in safety or effectiveness were observed between these subjects and younger subjects, and other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, use of gadobutrol in elderly patients should be cautious, reflecting the greater frequency of impaired renal function and concomitant disease or other drug therapy. No dose adjustment according to age is necessary in this population.

### **8.6 Renal Impairment**

Prior to administration of gadobutrol, screen all patients for renal dysfunction by

obtaining a history and/or laboratory tests [see *Warnings and Precautions* (5.2)]. No dosage adjustment is recommended for patients with renal impairment.


Gadobutrol can be removed from the body by hemodialysis [see *Warnings and Precautions* (5.2) and *Clinical Pharmacology* (12.3)].

## 10 OVERDOSAGE

The maximum dose of gadobutrol tested in healthy volunteers, 1.5 mL/kg body weight (1.5 mmol/kg; 15 times the recommended dose), was tolerated in a manner similar to lower doses. Gadobutrol can be removed by hemodialysis [see *Use in Specific Populations* (8.6) and *Clinical Pharmacology* (12.3)].

## 11 DESCRIPTION

Gadobutrol injection is a paramagnetic macrocyclic contrast agent administered intravenously for magnetic resonance imaging. The chemical name for gadobutrol is 10-[(1S,2R)-2,3-dihydroxy-1-hydroxymethylpropyl]-1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid, gadolinium complex. Gadobutrol has a molecular formula of  $C_{18}H_{31}GdN_4O_9$  and a molecular weight of 604.72.



Gadobutrol injection is a sterile, clear, colorless to pale yellow solution containing 604.72 mg (1.0 mmol) of gadobutrol per mL as the active ingredient with 0.513 mg of calcobutrol sodium, 1.211 mg of trometamol, hydrochloric acid (for pH adjustment) and water for injection.

Gadobutrol injection contains no preservatives.

The main physicochemical properties of gadobutrol injection (1 mmol/mL solution for injection) are listed below:

Density (g/mL at 37°C)	1.3
Osmolarity at 37°C (mOsm/L solution)	1117

Osmolality at 37°C (mOsm/kg H <sub>2</sub> O)	1603
Viscosity at 37°C (mPa·s)	4.96
pH	6.6 to 8

The thermodynamic stability constants for gadobutrol (log K<sub>therm</sub> and log K<sub>cond</sub> at pH 7.4) are 21.8 and 15.3, respectively.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

In MRI, visualization of normal and pathological tissue depends in part on variations in the radiofrequency signal intensity that occurs with:

- Differences in proton density
- Differences of the spin-lattice or longitudinal relaxation times (T<sub>1</sub>)
- Differences in the spin-spin or transverse relaxation time (T<sub>2</sub>)

When placed in a magnetic field, gadobutrol shortens the T<sub>1</sub> and T<sub>2</sub> relaxation times. The extent of decrease of T<sub>1</sub> and T<sub>2</sub> relaxation times, and therefore the amount of signal enhancement obtained from gadobutrol, is based upon several factors including the concentration of gadobutrol in the tissue, the field strength of the MRI system, and the relative ratio of the longitudinal and transverse relaxation times. At the recommended dose, the T<sub>1</sub> shortening effect is observed with greatest sensitivity in T<sub>1</sub>-weighted magnetic resonance sequences. In T<sub>2</sub>\*-weighted sequences the induction of local magnetic field inhomogeneities by the large magnetic moment of gadolinium and at high concentrations (during bolus injection) leads to a signal decrease.

### 12.2 Pharmacodynamics

Gadobutrol leads to distinct shortening of the relaxation times even in low concentrations. At pH 7, 37°C and 1.5 T, the relaxivity (r<sub>1</sub>) - determined from the influence on the relaxation times (T<sub>1</sub>) of protons in plasma - is 5.2 L/(mmol·sec) and the relaxivity (r<sub>2</sub>) - determined from the influence on the relaxation times (T<sub>2</sub>) - is 6.1 L/(mmol·sec). These relaxivities display only slight dependence on the strength of the magnetic field. The T<sub>1</sub> shortening effect of paramagnetic contrast agents is dependent on concentration and r<sub>1</sub> relaxivity (see Table 3). This may improve tissue visualization.

**Table 3: Relaxivity (r<sub>1</sub>) of Gadolinium Chelates at 1.5 T**

Gadolinium-Chelate	r <sub>1</sub> (L·mmol <sup>-1</sup> ·s <sup>-1</sup> )
Gadobenate	6.3
Gadobutrol	5.2
Gadodiamide	4.3
Gadofosveset	16
Gadopentetate	4.1
Gadoterate	3.6
Gadoteridol	4.1
Gadoversetamide	4.7
Gadoxetate	6.9

Compared to 0.5 molar gadolinium-based contrast agents, the higher concentration of gadobutrol results in half the volume of administration and a more compact contrast bolus injection. At the site of imaging, the relative height and width of the time intensity curve for gadobutrol varies as a function of imaging location and multiple patient, injection, and device-specific factors.

Gadobutrol is a water-soluble, hydrophilic compound with a partition coefficient between n-butanol and buffer at pH 7.6 of about 0.006.

## **12.3 Pharmacokinetics**

### *Distribution*

After intravenous administration, gadobutrol is rapidly distributed in the extracellular space. After a gadobutrol dose of 0.1 mmol/kg body weight, an average level of 0.59 mmol gadobutrol/L was measured in plasma 2 minutes after the injection and 0.3 mmol gadobutrol/L 60 minutes after the injection. Gadobutrol does not display any particular protein binding. Following GBCA administration, gadolinium is present for months or years in brain, bone, skin, and other organs [see *Warnings and Precautions* (5.4)].

### *Metabolism*

Gadobutrol is not metabolized.

### *Elimination*

Values for AUC, body weight normalized plasma clearance and half-life are given in Table 4, below.

Gadobutrol is excreted in an unchanged form via the kidneys. In healthy subjects, renal clearance of gadobutrol is 1.1 to 1.7 mL/(min•kg) and thus comparable to the renal clearance of inulin, confirming that gadobutrol is eliminated by glomerular filtration.

Within two hours after intravenous administration more than 50% and within 12 hours more than 90% of the given dose is eliminated via the urine. Extra-renal elimination is negligible.

### *Specific Populations*

#### Gender

Gender has no clinically relevant effect on the pharmacokinetics of gadobutrol.

#### Geriatric

A single intravenous dose of 0.1 mmol/kg gadobutrol was administered to 15 elderly and 16 non-elderly subjects. AUC was slightly higher and clearance slightly lower in elderly subjects as compared to non-elderly subjects [see *Use in Specific Populations* (8.5)].

#### Pediatric

The pharmacokinetics of gadobutrol were evaluated in two studies in a total of 130 patients age 2 to less than 18 years and in 43 patients less than 2 years of age (including term neonates). Patients received a single intravenous dose of 0.1 mmol/kg of gadobutrol. The pharmacokinetic profile of gadobutrol in pediatric patients is similar to



that in adults, resulting in similar values for AUC, body weight normalized plasma clearance, as well as elimination half-life. Approximately 99% (median value) of the dose was recovered in urine within 6 hours (this information was derived from the 2 to less than 18 year old age group).

**Table 4: Pharmacokinetics by Age Group (Median [Range])**

	<b>0 to &lt; 2 years N=43</b>	<b>2 to 6 years N=45</b>	<b>7 to 11 years N=39</b>	<b>12 to &lt; 18 years N=46</b>	<b>Adults N=93</b>
AUC (μmol·h/L)	781 [513, 1891]	846 [412, 1331]	1025 [623, 2285]	1237 [946, 2211]	1072 [667, 1992]
CL (L/h/kg)	0.128 [0.053, 0.195]	0.119 [0.08, 0.215]	0.099 [0.043, 0.165]	0.081 [0.046, 0.103]	0.094 [0.051, 0.15]
t <sub>1/2</sub> (h)	2.91 [1.6, 12.4]	1.91 [1.04, 2.7]	1.66 [0.91, 2.71]	1.68 [1.31, 2.48]	1.8 [1.2, 6.55]
C <sub>20</sub> (μmol/L)	367 [280, 427]	421 [369, 673]	462 [392, 760]	511 [387, 1077]	441 [281, 829]

### Renal Impairment

In patients with impaired renal function, the serum half-life of gadobutrol is prolonged and correlated with the reduction in creatinine clearance.

After intravenous injection of 0.1 mmol gadobutrol/kg body weight, the elimination half-life was  $5.8 \pm 2.4$  hours in mild to moderately impaired patients ( $80 > \text{CLCR} > 30$  mL/min) and  $17.6 \pm 6.2$  hours in severely impaired patients not on dialysis ( $\text{CLCR} < 30$  mL/min). The mean AUC of gadobutrol in patients with normal renal function was  $1.1 \pm 0.1$  mmol·h/L, compared to  $4 \pm 1.8$  mmol·h/L in patients with mild to moderate renal impairment and  $11.5 \pm 4.3$  mmol·h/L in patients with severe renal impairment.

Complete recovery in the urine was seen in patients with mild or moderate renal impairment within 72 hours. In patients with severely impaired renal function about 80% of the administered dose was recovered in the urine within 5 days. For patients receiving hemodialysis, physicians may consider the prompt initiation of hemodialysis following the administration of gadobutrol in order to enhance the contrast agent's elimination. Sixty-eight percent (68%) of gadobutrol is removed from the body after the first dialysis, 94% after the second dialysis, and 98% after the third dialysis session. *[See Warnings and Precautions (5.2) and Use in Specific Populations (8.6).]*

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

No carcinogenicity studies of gadobutrol have been conducted.

Gadobutrol was not mutagenic in *in vitro* reverse mutation tests in bacteria, in the HGPRT (hypoxanthine-guanine phosphoribosyl transferase) test using cultured Chinese hamster V79 cells, or in chromosome aberration tests in human peripheral blood lymphocytes, and was negative in an *in vivo* micronucleus test in mice after intravenous

injection of 0.5 mmol/kg.

Gadobutrol had no effect on fertility and general reproductive performance of male and female rats when given in doses 12.2 times the human equivalent dose (based on body surface area).

### 13.2 Animal Toxicology and/or Pharmacology

Local intolerance reactions, including moderate irritation associated with infiltration of inflammatory cells was observed after paravenous administration to rabbits, suggesting the possibility of occurrence of local irritation if the contrast medium leaks around veins in a clinical setting [see *Warnings and Precautions* (5.6)].

## 14 CLINICAL STUDIES

### 14.1 MRI of the CNS

Patients referred for MRI of the central nervous system with contrast were enrolled in two clinical trials that evaluated the visualization characteristics of lesions. In both studies, patients underwent a baseline, pre-contrast MRI prior to administration of gadobutrol at a dose of 0.1 mmol/kg, followed by a post-contrast MRI. In Study A, patients also underwent an MRI before and after the administration of gadoteridol. The studies were designed to demonstrate superiority of gadobutrol MRI to non-contrast MRI for lesion visualization. For both studies, pre-contrast and pre-plus-post contrast images (paired images) were independently evaluated by three readers for contrast enhancement and border delineation using a scale of 1 to 4, and for internal morphology using a scale of 1 to 3 (Table 5). Lesion counting was also performed to demonstrate non-inferiority of paired gadobutrol image sets to pre-contrast MRI. Readers were blinded to clinical information.

**Table 5: Primary Endpoint Visualization Scoring System**

Score	Visualization Characteristics		
	Contrast Enhancement	Border Delineation	Internal Morphology
1	None	None	Poorly visible
2	Weak	Moderate	Moderately visible
3	Clear	Clear but incomplete	Sufficiently visible
4	Clear and bright	Clear and complete	N/A

Efficacy was determined in 657 subjects. The average age was 49 years (range 18 to 85 years) and 42% were male. The ethnic representations were 39% Caucasian, 4% Black, 16% Hispanic, 38% Asian, and 3% of other ethnic groups.

Table 6 shows a comparison of visualization results between paired images and pre-contrast images. Gadobutrol provided a statistically significant improvement for each of the three lesion visualization parameters when averaged across three independent readers for each study.

**Table 6: Visualization Endpoint Results of Central Nervous System Adult MRI Studies with 0.1 mmol/kg Gadobutrol**

Endpoint	Study A N=336		Difference <sup>1</sup>	Study B N=321		
	Pre-contrast	Paired		Pre-contrast	Paired	Difference
Contrast Enhancement	0.97	2.26	1.29 <sup>2</sup>	0.93	2.86	1.94 <sup>2</sup>
Border Delineation	1.98	2.58	0.6 <sup>2</sup>	1.92	2.94	1.02 <sup>2</sup>
Internal Morphology	1.32	1.93	0.6 <sup>2</sup>	1.57	2.35	0.78 <sup>2</sup>
Average # Lesions Detected	8.08	8.25	0.17 <sup>4</sup>	2.65	2.97	0.32 <sup>3</sup>

<sup>1</sup> Difference of means = (paired mean) - (pre-contrast mean)

<sup>2</sup> p<0.001

<sup>3</sup> Met noninferiority margin of -0.35

<sup>4</sup> Did not meet noninferiority margin of -0.35

Performances of gadobutrol and gadoteridol for visualization parameters were similar. Regarding the number of lesions detected, Study B met the prespecified noninferiority margin of -0.35 for paired read versus pre-contrast read while in Study A, gadobutrol and gadoteridol did not.

For the visualization endpoints contrast enhancement, border delineation, and internal morphology, the percentage of patients scoring higher for paired images compared to pre-contrast images ranged from 93% to 99% for Study A, and 95% to 97% for Study B. For both studies, the mean number of lesions detected on paired images exceeded that of the pre-contrast images; 37% for Study A and 24% for Study B. There were 29% and 11% of subjects in which the pre-contrast images detected more lesions for Study A and Study B, respectively.

The percentage of patients whose average reader mean score changed by  $\leq 0$ , up to 1, up to 2, and  $\geq 2$  scoring categories presented in Table 5 is shown in Table 7. The categorical improvement of ( $\leq 0$ ) represents higher ( $< 0$ ) or identical ( $= 0$ ) scores for the pre-contrast read, the categories with scores  $> 0$  represent the magnitude of improvement seen for the paired read.

**Table 7: Primary Endpoint Visualization Categorical Improvement for Average Reader**

Endpoint	Study A N=336				Study B N=321			
	Categorical Improvement (Paired - Pre-Contrast) %				Categorical Improvement (Paired - Pre-Contrast) %			
	$\leq 0$	$> 0$ to $< 1$	1 to $< 2$	$\geq 2$	$\leq 0$	$> 0$ to $< 1$	1 to $< 2$	$\geq 2$
Contrast Enhancement	1	30	55	13	3	6	34	57
Border Delineation	7	73	18	1	5	38	51	5
Internal Morphology	4	79	17	0	5	61	33	1

For both studies, the improvement of visualization endpoints in paired gadobutrol

images compared to pre-contrast images resulted in improved assessment of normal and abnormal CNS anatomy.

### *Pediatric Patients*

Two studies in 44 pediatric patients age younger than 2 years and 135 pediatric patients age 2 to less than 18 years with CNS and non-CNS lesions supported extrapolation of adult CNS efficacy findings. For example, comparing pre vs paired pre- and post-contrast images, investigators selected the best of four descriptors under the heading, “Visualization of lesion-internal morphology (lesion characterization) or homogeneity of vessel enhancement” for 27/44 (62% = pre) vs 43/44 (98% = paired) MR images from patients age 0 to less than 2 years and 106/135 (78% = pre) vs 108/135 (80% = paired) MR images from patients age 2 to less than 18 years.

## **14.2 MRI of the Breast**

Patients with recently diagnosed breast cancer were enrolled in two identical clinical trials to evaluate the ability of gadobutrol to assess the presence and extent of malignant breast disease prior to surgery. Patients underwent non-contrast breast MRI (BMR) prior to gadobutrol (0.1 mmol/kg) breast MRI. BMR images and gadobutrol BMR (combined contrast plus non-contrast) images were independently evaluated in each study by three readers blinded to clinical information. In separate reading sessions the BMR images and gadobutrol BMR images were also interpreted together with X-ray mammography images (XRM).

The studies evaluated 787 patients: Study 1 enrolled 390 women with an average age of 56 years, 74% were white, 25% Asian, 0.5% black, and 0.5% other; Study 2 enrolled 396 women and 1 man with an average age of 57 years, 71% were white, 24% Asian, 3% black, and 2% other.

The readers assessed 5 regions per breast for the presence of malignancy using each reading modality. The readings were compared to an independent standard of truth (SoT) consisting of histopathology for all regions where excisions were made and tissue evaluated. XRM plus ultrasound was used for all other regions.

The assessment of malignant disease was performed using a region based within-subject sensitivity. Sensitivity for each reading modality was defined as the mean of the percentage of malignant breast regions correctly interpreted for each subject. The within-subject sensitivity of gadobutrol BMR was superior to that of BMR. The lower bound of the 95% Confidence Interval (CI) for the difference in within-subject sensitivity ranged from 19% to 42% for Study 1 and from 12% to 27% for Study 2. The within-subject sensitivity for gadobutrol BMR and BMR as well as for gadobutrol BMR plus XRM and BMR plus XRM is presented in Table 8.

**Table 8: Sensitivity of Gadobutrol BMR for Detection of Malignant Breast Disease**

<b>Study 1</b>					<b>Study 2</b>				
<b>Sensitivity (%)</b>					<b>Sensitivity (%)</b>				
<b>N=388 Patients</b>					<b>N=390 Patients</b>				
Reader	BMR	BMR + XRM	Gadobutrol BMR	Gadobutrol BMR	Reader	BMR	BMR + XRM	Gadobutrol BMR	Gadobutrol BMR

				+XRM					+XRM
1	37	71	83	84	4	73	83	87	90
2	49	76	80	83	5	57	81	89	90
3	63	75	87	87	6	55	80	86	88

Specificity was defined as the percentage of non-malignant breasts correctly identified as non-malignant. The lower limit of the 95% confidence interval for specificity of gadobutrol BMR was greater than 80% for 5 of 6 readers. (Table 9)

**Table 9: Specificity of Gadobutrol BMR in Non-Malignant Breasts**

Study 1			Study 2		
Specificity (%) N=372 Patients			Specificity (%) N=367 Patients		
Reader	Gadobutrol BMR	Lower Limit 95% CI	Reader	Gadobutrol BMR	Lower Limit 95% CI
1	86	82	4	92	89
2	95	93	5	84	80
3	89	85	6	83	79

Three additional readers in each study read XRM alone. For these readers over both studies, sensitivity ranged from 68% to 73% and specificity in non-malignant breasts ranged from 86% to 94%.

In breasts with malignancy, a false positive detection rate was calculated as the percentage of subjects for which the readers assessed a region as malignant which could not be verified by SoT. The false positive detection rates for gadobutrol BMR ranged from 39% to 53% (95% CI Upper Bounds ranged from 44% to 58%).

### 14.3 MRA

Patients with known or suspected disease of the supra-aortic arteries (for evaluation up to but excluding the basilar artery) were enrolled in Study C, and patients with known or suspected disease of the renal arteries were enrolled in Study D. In both studies, non-contrast, 2D time-of-flight (ToF) magnetic resonance angiography (MRA) was performed prior to gadobutrol MRA using a single intravenous injection of 0.1 mmol/kg. The injection rate of 1.5 mL/second was selected to extend the injection duration to at least half of the imaging duration. Imaging was performed with parallel-channel, 1.5T MRI devices and an automatic bolus tracking technique to trigger the image acquisition following gadobutrol administration using elliptically encoded, T1-weighted, 3D gradient-echo image acquisition and single breath hold. Three central readers blinded to clinical information interpreted the ToF and gadobutrol MRA images. Three additional central readers interpreted separately acquired computed tomographic angiography (CTA) images, which were used as the standard of reference (SoR) in each study.

The studies included 749 subjects: 457 were evaluated in Study C, with an average age of 68 (range 25 to 93); 64% were male; 80% white, 28% black, and 16% Asian. An additional 292 subjects were evaluated in Study D, with an average age of 55 (range 18 to 88); 54% were male; 68% white, 7% black, and 22% Asian.

Efficacy was evaluated based on anatomical visualization and performance for distinguishing between normal and abnormal anatomy. The visualization metric depended on whether readers selected, “Yes, it can be visualized along its entire length...” when responding to the question, “Is this segment assessable?”. Twenty-one segments in Study C and six segments in Study D were presented per subject to each reader. The performance metrics, sensitivity and specificity, depended on digital caliper-based quantitation of arterial narrowing in visualized, non-occluded, abnormal-appearing segments. Significant stenosis was defined as at least 70% in Study C and 50% in Study D. Performance of gadobutrol MRA compared to ToF MRA was calculated using an imputation method for non-visualized segments by assigning them as a 50% match with SoR and a 50% mismatch. Performance of gadobutrol MRA compared to a pre-specified threshold of 50% was calculated after excluding non-visualized segments. Measurement variability and visualization of accessory renal arteries was also evaluated.

Results were analyzed for each of the three central readers.

**Table 10: Visualization, Sensitivity, Specificity**

<b>STUDY C: SUPRA-AORTIC ARTERIES (457 patients)</b>									
<b>Performance at the segment level</b>									
<b>9597<sup>1</sup> segments of which 158<sup>1</sup> were positive for stenosis by SoR<sup>2</sup></b>									
	<b>VISUALIZATION (%)</b>			<b>SENSITIVITY (%)</b>			<b>SPECIFICITY (%)</b>		
<b>READER</b>	<b>GAD MRA</b>	<b>ToF MRA</b>	<b>GAD – ToF (CI<sup>3</sup>)</b>	<b>GAD MRA</b>	<b>ToF MRA</b>	<b>GAD – ToF (CI<sup>4</sup>)</b>	<b>GAD MRA</b>	<b>ToF MRA</b>	<b>GAD – ToF (CI<sup>4</sup>)</b>
1	88	24	64 (61, 67)	60	54	6 (-4, 14)	92	62	30 (29, 32)
2	95	75	20 (18, 21)	60	54	6 (-3, 14)	95	85	10 (9, 11)
3	97	82	15 (13, 17)	58	55	3 (-4, 11)	97	89	8 (7, 9)
<b>STUDY D: RENAL ARTERIES (292 patients)</b>									
<b>Performance at the segment level</b>									
<b>1752<sup>1</sup> segments of which 133<sup>1</sup> were positive for stenosis by SoR<sup>2</sup></b>									
4	98	82	16 (13, 20)	52	51	1 (-9, 11)	94	83	11 (9, 14)
5	96	72	24 (21, 28)	54	39	15 (6, 24)	95	85	10 (8, 12)
6	96	78	17 (14, 21)	53	50	3 (-6, 12)	94	81	13 (11, 16)

<sup>1</sup>Number of segments varied between readers; number for majority-reader shown.

<sup>2</sup>Standard of Reference based on aggregate interpretation of three central CTA readers.

<sup>3</sup>95.1/95% (Study C/D) confidence interval for two-sided comparison.

<sup>4</sup>90.1/90% (Study C/D) confidence interval for one-sided comparison against non-inferiority margin of -7.5.

GAD MRA = Post-contrast Gadobutrol Magnetic Resonance Angiography, ToF = Non-

contrast 2D-Time of Flight.

For all three supra-aortic artery readers in Study C, the lower bound of confidence for the sensitivity of gadobutrol MRA did not exceed 54%. For all three renal artery readers in Study D, the lower bound of confidence for the sensitivity of gadobutrol MRA did not exceed 46%.

#### *Measurement Variability*

For both MRA and CTA, readers varied in the quantity of narrowing they assigned to the same arterial segments. Table 11 shows the percentage of patients in whom the measurement range was 30% or greater for the left or right internal carotid and proximal renal artery segments. There were approximately four measurements per patient segment, one from the site and three from the central readers. Measurement variability was high for both CTA and MRA, but numerically lower for gadobutrol compared to non-contrast ToF MRA.

**Table 11: Percent of Patients with Range  $\geq 30\%$ ,  $\geq 50\%$ ,  $\geq 70\%$  for Measurement of Stenoses and Normal Vessel Diameters**

	Internal Carotid				Proximal Main Renal			
	N	$\geq 30\%$	$\geq 50\%$	$\geq 70\%$	N	$\geq 30\%$	$\geq 50\%$	$\geq 70\%$
CTA	456	40	11	4	292	59	33	9
ToF MRA	443	55	22	9	270	44	22	9
Gadobutrol MRA	454	47	13	4	286	34	14	4

#### *Visualization of Accessory Renal Arteries for Surgical Planning and Renal Donor Evaluation (Study D only)*

Of 1752 main arteries visualized by the central CTA readers, 266 (15%) were also associated with positive visualization of at least one accessory (duplicate) artery. With the central MRA readers, the comparable rates were 232 of 1752 (13%) for gadobutrol MRA compared to 53 of 1752 (3%) for ToF MRA.

### **14.4 Cardiac MRI**

Two studies similar in design, Study E and Study F, evaluated the sensitivity and specificity of gadobutrol cardiac MRI (CMRI) for detection of coronary artery disease (CAD) in adult patients with known or suspected CAD. Patients were excluded from study if they had a history of coronary artery bypass grafting, or if it was known in advance that they were unable to hold their breath, or had atrial fibrillation or other arrhythmia likely to prevent electrocardiogram-gated CMRI. The studies were multi-center, open-label, and evaluated 764 subjects for efficacy: 376 in Study E, with an average age of 59 (range 20 to 84); 69% male; 74% white, 1% black, and 25% Asian; and 388 subjects in Study F, with an average age of 59 (range 23 to 82); 61% male; 67% white, 17% black, and 12% Asian.

All subjects underwent dynamic first-pass gadobutrol imaging during vasodilator stress, followed ~10 minutes later by dynamic first-pass gadobutrol imaging at rest, followed ~5 minutes later with imaging during a period of gradual gadobutrol washout from the myocardium (late gadolinium enhancement, LGE). Imaging was performed on 1.5 T or 3.0 T MRI devices equipped with multichannel surface coils to support accelerated

acquisitions with parallel imaging, T1-weighted, 2D gradient-echo, dynamic acquisition of perfusion with at least 3 slices per heartbeat. Gadobutrol was administered intravenously at a rate of ~4 mL/second as two separate bolus injections (0.05 mmol/kg each), the first at peak pharmacologic stress (~3 minutes after start of ongoing adenosine infusion, or immediately after completion of regadenoson administration, at approved doses). No additional gadobutrol was administered for LGE imaging.

Images were read by three independent readers blinded to clinical information. Reader detection of CAD depended on visually detecting defective perfusion or scar on gadobutrol CMRI (stress, rest, LGE) imaging. Quantitative coronary angiography (QCA) was used to measure intraluminal narrowing and served as the standard of reference (SoR).

Computed tomographic angiography (CTA) was used as the SoR if disease could be unequivocally excluded, and no coronary angiography (CA) was available. The left ventricular myocardium was divided into six regions. Readers provided per-region (CMRI, CTA) and per-artery (QCA) interpretations for each subject. Subject-level endpoints reflected each subject's most abnormal localized finding.

The sensitivity results for gadobutrol CMRI to detect CAD defined as either maximum stenosis  $\geq 50\%$  or  $\geq 70\%$  by QCA are presented in Table 12. For each reader, sensitivity of gadobutrol CMRI larger than 60% can be concluded if the lower 95% confidence limit of the sensitivity estimate exceeds the pre-specified threshold of 60%.

**Table 12: Sensitivity (%) of Gadobutrol CMRI for Detection of CAD in Patients with Maximum Stenosis\* of  $\geq 50\%$  and  $\geq 70\%$**

	<b>Study E</b>		<b>Study F</b>	
	<b><math>\geq 50\%</math> N=141</b>	<b><math>\geq 70\%</math> N=108</b>	<b><math>\geq 50\%</math> N=150</b>	<b><math>\geq 70\%</math> N=105</b>
<b>Reader 1**</b>	77 ( <b>69</b> , 83)***	90 ( <b>83</b> , 95)	65 ( <b>57</b> , 72)	77 ( <b>68</b> , 85)
<b>Reader 2**</b>	65 ( <b>57</b> , 73)	80 ( <b>71</b> , 87)	56 ( <b>48</b> , 64)	71 ( <b>62</b> , 80)
<b>Reader 3**</b>	65 ( <b>56</b> , 72)	79 ( <b>70</b> , 86)	61 ( <b>53</b> , 69)	76 ( <b>67</b> , 84)

\* Stenosis determined by Quantitative Coronary Angiography (QCA)

\*\* CMRI images were assessed by six independent blinded readers, three in each study.

\*\*\* The bolded value represents the lower limit of the 95% confidence interval, which is compared to a pre-specified threshold of 60% for evaluation of sensitivity.

The specificity results for gadobutrol CMRI to detect CAD defined as either maximum stenosis  $\geq 50\%$  or  $\geq 70\%$  by QCA are presented in Table 13. For each reader, specificity of gadobutrol CMRI larger than 55% can be concluded if the lower 95% confidence limit of the specificity estimate exceeds the pre-specified threshold of 55%.

**Table 13: Specificity (%) of Gadobutrol CMRI for Exclusion of CAD in Patients with Maximum Stenosis\* of  $\geq 50\%$  and  $\geq 70\%$**

	<b>Study E</b>		<b>Study F</b>	
	<b><math>\geq 50\%</math></b>	<b><math>\geq 70\%</math></b>	<b><math>\geq 50\%</math></b>	<b><math>\geq 70\%</math></b>



	N=235	N=268	N=239	N=283
<b>Reader 1**</b>	85 ( <b>80</b> , 89)***	83 ( <b>78</b> , 87)	85 ( <b>80</b> , 90)	82 ( <b>77</b> , 86)
<b>Reader 2**</b>	92 ( <b>88</b> , 95)	91 ( <b>87</b> , 94)	89 ( <b>84</b> , 92)	87 ( <b>83</b> , 91)
<b>Reader 3**</b>	92 ( <b>88</b> , 95)	91 ( <b>87</b> , 94)	90 ( <b>85</b> , 93)	87 ( <b>82</b> , 91)

\* Stenosis determined by Quantitative Coronary Angiography (QCA)

\*\* CMRI images were assessed by six independent blinded readers, three in each study.

\*\*\* The bolded value represents the lower limit of the 95% confidence interval, which is compared to a pre-specified threshold of 55% for evaluation of specificity.

In Study E, among the 33 patients with maximum stenosis by QCA between 50% and <70%, the proportion of gadobutrol-CMRI positive detections of CAD ranged from 15% to 33%. In Study F, among the 45 patients with maximum stenosis by QCA between 50% and < 70%, the proportion of gadobutrol-CMRI positive detections of CAD ranged from 20% to 35%. The results of gadobutrol-CMRI reads to detect CAD in patients with maximum stenosis between 50% and < 70% are summarized in Table 14.

**Table 14: Gadobutrol-CMRI Detection of CAD in Patients with Maximum Stenosis\* between 50% and < 70%**

	<b>Study E (n=33)</b>	<b>Study F (n=45)</b>
	<b>Gadobutrol-CMRI positive</b>	<b>Gadobutrol-CMRI positive</b>
<b>Reader 1**</b>	11 (33%)	16 (35%)
<b>Reader 2**</b>	5(15%)	9 (20%)
<b>Reader 3**</b>	6(18%)	12 (26%)

\* Stenosis determined by Quantitative Coronary Angiography (QCA).

\*\*CMRI images were assessed by six independent blinded readers, three in each study.

### Left Mainstem Stenosis (LMS)

The studies did not include sufficient numbers of subjects to characterize the performance of gadobutrol CMRI for detection of LMS, a subgroup at high risk from false negative reads. In Studies E and F, only three subjects had isolated LMS stenosis >50%. In two of the three cases, the CMRI was interpreted as normal by at least two of the three readers (false negative). Sixteen subjects had LMS stenosis >50% (including subjects with isolated LMS stenosis and subjects with LMS stenosis in addition to stenoses elsewhere). In five of these sixteen cases, the CMR was interpreted as normal by at least two of the three readers (false negative).

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

## 16.1 How Supplied

Gadobutrol injection is a sterile, clear and colorless to pale yellow solution containing 604.72 mg gadobutrol per mL (equivalent to 1 mmol gadobutrol per mL). Gadobutrol injection is supplied in the following Multiple-Dose container sizes:

Product Code	Unit of Sale	Each
287230	NDC 65219-287-30 Packaged in cartons of 10.	NDC 65219-287-10 30 mL Imaging Bulk Package with rubber stopper.
287265	NDC 65219-289-65 Packaged in cartons of 10.	NDC 65219-289-10 65 mL Imaging Bulk Package with rubber stopper.

## 16.2 Storage and Handling

Store at 25°C (77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].

Should freezing occur, gadobutrol injection should be brought to room temperature before use. If allowed to stand at room temperature, gadobutrol injection should return to a clear and colorless to pale yellow solution. Visually inspect gadobutrol injection for particulate matter and discoloration prior to administration. Do not use the solution if it is discolored, if particulate matter is present or if the container appears damaged.

## 17 PATIENT COUNSELING INFORMATION

- Advise the patient to read the FDA-approved patient labeling (Medication Guide).

### Nephrogenic Systemic Fibrosis

Instruct patients to inform their physician if they:

- Have a history of kidney disease and/or liver disease, or
- Have recently received a GBCA

GBCAs increase the risk of NSF among patients with impaired elimination of drugs. To counsel patients at risk of NSF:

- Describe the clinical manifestation of NSF
- Describe procedures to screen for the detection of renal impairment

Instruct the patients to contact their physician if they develop signs or symptoms of NSF following gadobutrol injection administration, such as burning, itching, swelling, scaling, hardening and tightening of the skin; red or dark patches on the skin; stiffness in joints with trouble moving, bending or straightening the arms, hands, legs or feet; pain in the hip bones or ribs; or muscle weakness.

### Common Adverse Reactions

Inform patients that they may experience:

- Reactions along the venous injection site, such as mild and transient burning or pain or feeling of warmth or coldness at the injection site
- Side effects of headache, nausea, abnormal taste and feeling hot

## General Precautions

### Gadolinium Retention

- Advise patients that gadolinium is retained for months or years in brain, bone, skin, and other organs in patients with normal renal function. The clinical consequences of retention are unknown. Retention depends on multiple factors and is greater following administration of linear GBCAs than following administration of macrocyclic GBCAs [see *Warnings and Precautions (5.4)*].

Instruct patients receiving gadobutrol injection to inform their physician if they:

- Are pregnant or breastfeeding
- Have a history of allergic reaction to contrast media, bronchial asthma or allergic respiratory disorder

Manufactured for:



Lake Zurich, IL 60047

[www.fresenius-kabi.com/us](http://www.fresenius-kabi.com/us)

Product of China

Revised: September 2024

451796B

26EUF03

## Medication Guide

<p style="text-align: center;"><b>Gadobutrol Injection</b> <b>Gadobutrol (“gad” oh bue’ trol)</b> <b>Injection for intravenous use</b></p>
<p><b>What is gadobutrol injection?</b></p> <ul style="list-style-type: none"><li>• Gadobutrol injection is a prescription medicine called a gadolinium-based contrast agent (GBCA). Gadobutrol injection, like other GBCAs, is injected into your vein and used with a magnetic resonance imaging (MRI) scanner.</li><li>• An MRI exam with a GBCA, including gadobutrol injection, helps your doctor to see problems better than an MRI exam without a GBCA.</li><li>• Your doctor has reviewed your medical records and has determined that you would benefit from using a GBCA with your MRI exam.</li></ul>
<p><b>What is the most important information I should know about gadobutrol injection?</b></p> <ul style="list-style-type: none"><li>• GBCAs like Gadobutrol injection may cause serious side effects including death, coma, encephalopathy, and seizures when it is given intrathecally (injection given into the spinal canal). It is not known if Gadobutrol injection is safe and effective with intrathecal use. Gadobutrol injection is not approved for this use.</li></ul>

- Gadobutrol injection contains a metal called gadolinium. Small amounts of gadolinium can stay in your body including the brain, bones, skin and other parts of your body for a long time (several months to years).
- It is not known how gadolinium may affect you, but so far, studies have not found harmful effects in patients with normal kidneys.
- Rarely, patients have reported pains, tiredness, and skin, muscle or bone ailments for a long time, but these symptoms have not been directly linked to gadolinium.
- There are different GBCAs that can be used for your MRI exam. The amount of gadolinium that stays in the body is different for different gadolinium medicines. Gadolinium stays in the body more after Omniscan or Optimark than after Eovist, Magnevist, or MultiHance. Gadolinium stays in the body the least after Dotarem, Gadobutrol injection, or ProHance.
- People who get many doses of gadolinium medicines, women who are pregnant and young children may be at increased risk from gadolinium staying in the body.
- Some people with kidney problems who get gadolinium medicines can develop a condition with severe thickening of the skin, muscles and other organs in the body (nephrogenic systemic fibrosis). Your healthcare provider should screen you to see how well your kidneys are working before you receive gadobutrol injection.

**Do not receive gadobutrol injection if you** have had a severe allergic reaction to gadobutrol injection.

**Before receiving gadobutrol injection, tell your healthcare provider about all your medical conditions, including if you:**

- have had any MRI procedures in the past where you received a GBCA. Your healthcare provider may ask you for more information including the dates of these MRI procedures.
- are pregnant or plan to become pregnant. It is not known if gadobutrol injection can harm your unborn baby. Talk to your healthcare provider about the possible risks to an unborn baby if a GBCA such as gadobutrol injection is received during pregnancy.
- have kidney problems, diabetes, or high blood pressure
- have had an allergic reaction to dyes (contrast agents) including GBCAs

**What are the possible side effects of gadobutrol injection?**

- **See “What is the most important information I should know about gadobutrol injection?”**
- **Allergic reactions. Gadobutrol injection can cause allergic reactions that can sometimes be serious. Your healthcare provider will monitor you closely for symptoms of an allergic reaction.**

**The most common side effects of gadobutrol injection include: headache, nausea, and dizziness.**

These are not all the possible side effects of gadobutrol injection.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**General information about the safe and effective use of gadobutrol injection.**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. You can ask your healthcare provider for information about gadobutrol injection that is written for health professionals.

**What are the ingredients in gadobutrol injection?**

Active ingredient: gadobutrol

Inactive ingredients: calcobutrol sodium, trometamol, hydrochloric acid (for pH adjustment) and water for injection

Manufactured for:



Lake Zurich, IL 60047

[www.fresenius-kabi.com/us](http://www.fresenius-kabi.com/us)

For more information, call Fresenius Kabi USA, LLC at 1-800-551-7176.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

26EUF03

9/2024

451796B

**PRINCIPAL DISPLAY PANEL - 30 mmol/30 mL (1 mmol/mL) - Shelf Carton**

**Dose: 0.1 mL/kg**

NDC 65219-287-30

**Gadobutrol Injection**

**30 mmol/30 mL**

(1 mmol/mL)

**Imaging Bulk Package - Not for Direct Infusion**

**For Intravenous Use Only**

**Discard 24 hours after initial puncture**

**Dispense the enclosed Medication Guide to each patient.**



(01)20365219287302



Manufactured for:

www.fresenius-kabi.com/us

Lake Zurich, IL 60047

SP26EUS01

621882

Product of China

For use only with an automated contrast injection system, contrast management system, or contrast media transfer set approved or cleared for use with this contrast agent in this imaging Bulk Package.  
See drug and device labeling for information on devices indicated or use with this imaging Bulk Package and techniques to help assure safe use.  
Each mL contains 604.72 mg of gadobutrol with 0.513 mg of calcobutrol sodium, 1.211 mg of trometamol, hydrochloric acid (for pH adjustment) and water for injection.  
Gadobutrol injection contains no antimicrobial preservative.  
Store at 25°C (77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].  
Recommended Dose: See Prescribing Information.

**Dose: 0.1 mL/kg**

**10 x 30 mL**  
Multiple-Dose Bottles

Rx only  
Sterile Solution

NDC 65219-287-30

## Gadobutrol Injection

**30 mmol/30 mL**  
**(1 mmol/mL)**

**Imaging Bulk Package -  
Not for Direct Infusion**

**For Intravenous Use Only**

**Discard 24 hours after initial puncture**

**Dispense the enclosed Medication Guide to each patient.**



**Dose: 0.1 mL/kg**

**10 x 30 mL**  
Multiple-Dose Bottles

Rx only  
Sterile Solution

NDC 65219-287-30

## Gadobutrol Injection

**30 mmol/30 mL**  
**(1 mmol/mL)**

**Imaging Bulk Package - Not for Direct Infusion**

**For Intravenous Use Only**

**Discard 24 hours after initial puncture**

**Dispense the enclosed Medication Guide to each patient.**





**PRINCIPAL DISPLAY PANEL - 30 mmol/30 mL (1 mmol/mL) - Bottle Label**

**Dose: 0.1 mL/kg**

NDC 65219-287-10

**Gadobutrol**

**Injection**

**30 mmol/30 mL**

(1 mmol/mL)


**Imaging Bulk Package -  
Not for Direct Infusion**

**For Intravenous Use Only**

**Multiple-Dose Bottle**

**Discard 24 hours after initial  
puncture**

Discard after \_\_\_\_ / \_\_\_\_ / \_\_\_\_  
at \_\_\_\_ : \_\_\_\_

Gadobutrol Injection 30 mmol/30 mL (1 mmol/mL)	Lot	Gadobutrol Injection 30 mmol/30 mL (1 mmol/mL)	Lot	- 30 mL - Injection	NDC 65219-287-10	<b>Dose: 0.1 mL/kg</b>	<b>30 mL</b>	<b>Rx only Sterile Solution</b>	For use only with an automated contrast injection system, contrast management system, or contrast media transfer set approved or cleared for use with this contrast agent in this Imaging Bulk Package. See drug and device labeling for information on devices indicated for use with this Imaging Bulk Package and techniques to help assure safe use.	Each mL contains 604.72 mg of gadobutrol with 0.513 mg of calcobutrol sodium, 1.211 mg of trometamol, hydrochloric acid (for pH adjustment) and water for injection.	Manufactured for: <b>FRESENIUS KABI</b> Lake Zurich, IL 60047 Product of China <b>403991</b> 26EUA01	 (01)00365219287100	
Gadobutrol Injection 30 mmol/30 mL (1 mmol/mL)	Lot	Gadobutrol Injection 30 mmol/30 mL (1 mmol/mL)	Lot			<b>Gadobutrol Injection</b>	<b>30 mmol/30 mL (1 mmol/mL)</b>		<b>Imaging Bulk Package - Not for Direct Infusion</b>				
						<b>For Intravenous Use Only Multiple-Dose Bottle Discard 24 hours after initial puncture</b>							
						Discard after ____ / ____ / ____ at ____ : ____							
						<b>Recommended Dose: See Prescribing Information.</b>							
						Store at 25°C (77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].							
				Lot/Exp.									

**PRINCIPAL DISPLAY PANEL - 65 mmol/65 mL (1 mmol/mL) - Shelf Carton**

**Dose: 0.1 mL/kg**

NDC 65219-289-65

**Gadobutrol Injection**

**65 mmol/65 mL**

(1 mmol/mL)

**Imaging Bulk Package - Not for Direct Infusion**

**For Intravenous Use Only**

**Discard 24 hours after initial puncture**

**Dispense the enclosed Medication Guide to each patient.**





(01)20365219289658

Manufactured for:

**FRESENIUS**  
**KABI**

Lake Zurich, IL 60047

www.fresenius-kabi.com/us

621883  
SP26FUS01

Product of China

For use only with an automated contrast injection system, contrast management system, or contrast media transfer set approved or cleared for use with this contrast agent in this imaging Bulk Package.  
See drug and device labeling for information on devices indicated or use with this imaging Bulk Package and techniques to help assure safe use.  
Each mL contains 604.72 mg of gadobutrol with 0.513 mg of calcobutrol sodium, 1.211 mg of trometamol, hydrochloric acid (for pH adjustment) and water for injection.  
Gadobutrol injection contains no antimicrobial preservative.  
Store at 25°C (77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].  
Recommended Dose: See Prescribing Information.

**Dose: 0.1 mL/kg**

NDC 65219-289-65

## Gadobutrol Injection

65 mmol/65 mL  
(1 mmol/mL)

10 x 65 mL  
Multiple-Dose Bottles

Rx only  
Sterile Solution

**Imaging Bulk Package - Not for Direct Infusion**

For Intravenous Use Only

Discard 24 hours after initial puncture

Dispense the enclosed Medication Guide to each patient.

**FRESENIUS**  
**KABI**

**Dose: 0.1 mL/kg**

NDC 65219-289-65

## Gadobutrol Injection

65 mmol/65 mL  
(1 mmol/mL)

10 x 65 mL  
Multiple-Dose Bottles

Rx only  
Sterile Solution

**Imaging Bulk Package - Not for Direct Infusion**

For Intravenous Use Only



### Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
Gadobutrol (UNII: 1BJ477IO2L) (Gadolinium Cation (3+) - UNII:AZV954TZ9N)		Gadobutrol	604.72 mg in 1 mL	
Inactive Ingredients				
Ingredient Name			Strength	
Tromethamine (UNII: 023C2WHX2V)				
Calcobutrol Sodium (UNII: PO5286589K)				
Hydrochloric Acid (UNII: QTT17582CB)				
Water (UNII: 059QF0KO0R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65219-287-30	10 in 1 CARTON	10/01/2023	
1	NDC:65219-287-10	30 mL in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA216081	10/01/2023	

GADOBUTROL			
gadobutrol injection			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:65219-289
Route of Administration	INTRAVENOUS		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
Gadobutrol (UNII: 1BJ477IO2L) (Gadolinium Cation (3+) - UNII:AZV954TZ9N)		Gadobutrol	604.72 mg in 1 mL
Inactive Ingredients			
Ingredient Name			Strength
Tromethamine (UNII: 023C2WHX2V)			
Calcobutrol Sodium (UNII: PO5286589K)			
Hydrochloric Acid (UNII: QTT17582CB)			

<b>Water</b> (UNII: 059QF0KO0R)				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65219-289-65	10 in 1 CARTON	10/01/2023	
1	NDC:65219-289-10	65 mL in 1 BOTTLE; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA216081	10/01/2023	

**Labeler** - Fresenius Kabi USA, LLC (013547657)

**Registrant** - Jiangsu Hengrui Pharmaceuticals Co., Ltd. (654147255)

<b>Establishment</b>				
Name		Address	ID/FEI	Business Operations
Jiangsu Hengrui Pharmaceuticals Co., Ltd. (Dongjin Road Site)			421324417	MANUFACTURE(65219-287, 65219-289)

Revised: 12/2024

Fresenius Kabi USA, LLC