

**TRAVEL EASE- meclizine hcl tablet, chewable**  
**Chain Drug Marketing Association, Inc.**

-----  
**Quality Choice 44-404A**

***Active ingredient (in each chewable tablet)***

Meclizine HCl 25 mg

***Purpose***

Antiemetic

***Uses***

for prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness

***Warnings***

**Do not use**

for children under 12 years of age unless directed by a doctor.

**Ask a doctor before use if you have**

- glaucoma
- difficulty in urination due to enlargement of the prostate gland
- a breathing problem such as emphysema or chronic bronchitis

**Ask a doctor or pharmacist before use if you are**

taking sedatives or tranquilizers.

**When using this product**

- drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- **do not take more than directed**
- dosage should be taken one hour before travel starts
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- adults and children 12 years and over: take 1 to 2 chewable tablets once daily or as directed by a doctor

**Other information**

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from heat and humidity
- see end flap for expiration date and lot number

**Inactive ingredients**

colloidal silicon dioxide, dextrose, FD&C red #40 aluminum lake, flavor, lactose monohydrate, magnesium stearate, pregelatinized starch, propylene glycol, saccharin sodium, silicon dioxide

**Questions or comments?**

**1-800-426-9391**

**Principal Display Panel**

QC  
QUALITY  
CHOICE

NDC 83324-027-16

\*Compare to the  
Active Ingredient in  
Bonine® Chewable Tablets

**Travel Ease****Meclizine HCl 25 mg Chewable Tablets | Antiemetic**

Helps Prevent Nausea, Vomiting and Dizziness  
Due to Motion Sickness for up to 24 Hours

Chew or crush tablets  
completely before swallowing.

Raspberry Flavored

**16** Tablets (25 mg each)

**Actual Size**

Distributed by CDMA, Inc.  
Novi, MI 48375

[www.qualitychoice.com](http://www.qualitychoice.com)

Questions: 800-935-2362

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF  
BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

\*This product is not manufactured or  
distributed by WellSpring Pharmaceutical  
Corporation, owner of the registered  
trademark Bonine® Chewable Tablets.  
50844 ORG082340421

NDC 83324-027-16



# Travel Ease

Ages 12 Years and Over

## Meclizine HCl 25 mg Chewable Tablets | Antiemetic

Helps Prevent Nausea, Vomiting and Dizziness  
Due to Motion Sickness for up to 24 Hours

NDC 83324-027-16



\*Compare to the  
Active Ingredient in  
Bonine® Chewable Tablets

# Travel Ease

## Meclizine HCl 25 mg Chewable Tablets | Antiemetic

Helps Prevent Nausea, Vomiting and Dizziness  
Due to Motion Sickness for up to 24 Hours

Chew or crush tablets  
completely before swallowing.

Raspberry Flavored



**16** Tablets (25 mg each)

Actual Size

LEBG296G4  
(E1042)

\*This product is not manufactured or  
distributed by WellSpring Pharmaceutical  
Corporation, owner of the registered  
trademark Bonine® Chewable Tablets.  
50844 ORG082340421

Distributed by CDMA, Inc.  
Novi, MI 48375  
www.qualitychoice.com  
Questions: 800-935-2362



B-0220-404A-21  
ORG082340421

### Questions or comments? 1-800-426-9391

**Inactive ingredients** colloidal silicon dioxide,  
dextrose, FD&C red #40 aluminum lake, flavor, lactose  
monohydrate, magnesium stearate, pregelatinized  
starch, polyethylene glycol, saccharin sodium, silicon  
dioxide

■ see end flap for expiration date and lot number  
■ protect from heat and humidity  
■ store at 25°C (77°F); excursions permitted between  
15°-30°C (59°-86°F)

**Other information**  
■ TAMPER EVIDENT; DO NOT USE IF OUTER PACKAGE  
IS OPENED OR BUSTER IS TORN OR BROKEN

**Directions** ■ do not take more than directed  
■ do not swallow tablets whole  
■ adults and children 12 years and over: take 1 to 2  
chewable tablets once daily or as directed by a doctor

**Purpose**  
Meclizine HCl 25 mg  
Antiemetic

**Warnings**  
Do not use for children under 12 years of age unless  
directed by a doctor.

■ Ask a doctor before use if you have  
■ difficulty in urination due to enlargement of the  
prostate gland ■ glaucoma  
■ a breathing problem such as emphysema or chronic  
bronchitis

■ Ask a doctor or pharmacist before use if you are  
taking sedatives or tranquilizers.

**When using this product** ■ drowsiness may occur  
■ alcohol, sedatives, and tranquilizers may increase  
■ drowsiness ■ avoid alcoholic beverages  
■ use caution when driving a motor vehicle or  
operating machinery


■ If pregnant or breast-feeding, ask a health  
professional before use.

**Uses** for prevention and treatment of nausea,  
vomiting, or dizziness associated with motion sickness

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF  
BUSTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

No print/No varnish  
Lot & Exp date





Quality choice 44-404A

**TRAVEL EASE**  
meclizine hcl tablet, chewable

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:83324-027
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>MECLIZINE HYDROCHLORIDE</b> (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)	MECLIZINE HYDROCHLORIDE	25 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>DEXTROSE, UNSPECIFIED FORM</b> (UNII: IY9XDZ35W2)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	

**Product Characteristics**

<b>Color</b>	pink	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	9mm
<b>Flavor</b>	RASPBERRY	<b>Imprint Code</b>	44;404
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83324-027-16	2 in 1 CARTON	03/27/2024	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M009	03/27/2024	

**Labeler** - Chain Drug Marketing Association, Inc. (011920774)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(83324-027) , pack(83324-027)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(83324-027)

Revised: 3/2024

Chain Drug Marketing Association, Inc.