


**DIPHENHYDRAMINE HYDROCHLORIDE- diphenhydramine hydrochloride injection, solution**  
**Armas Pharmaceuticals Inc.**

-----

**Diphenhydramine Hydrochloride Injection, USP**

**DESCRIPTION**

Diphenhydramine hydrochloride USP is an antihistamine drug having the chemical name 2-(Diphenylmethoxy)-N,N-dimethylethylamine hydrochloride. It occurs as a white, crystalline powder, is freely soluble in water and alcohol and has a molecular weight of 291.82. The molecular formula is  $C_{17}H_{21}NO \cdot HCl$ . The structural formula is as follows:



Diphenhydramine hydrochloride USP in the parenteral form is a sterile, pyrogen-free solution available in a concentration of 50 mg of diphenhydramine hydrochloride USP per mL. The solutions for parenteral use have been adjusted to a pH between 4 and 6.5 with either sodium hydroxide or hydrochloric acid.

**CLINICAL PHARMACOLOGY**

Diphenhydramine hydrochloride is an antihistamine with anticholinergic (drying) and sedative side effects. Antihistamines appear to compete with histamine for cell receptor sites on effector cells.

Diphenhydramine hydrochloride in the injectable form has a rapid onset of action. Diphenhydramine hydrochloride is widely distributed throughout the body, including the CNS. A portion of the drug is excreted unchanged in the urine, while the rest is metabolized via the liver. Detailed information on the pharmacokinetics of Diphenhydramine Hydrochloride Injection is not available.

**INDICATIONS AND USAGE**

Diphenhydramine hydrochloride in the injectable form is effective in adults and pediatric patients, other than premature infants and neonates, for the following conditions when diphenhydramine hydrochloride in the oral form is impractical.

**Antihistaminic**

For amelioration of allergic reactions to blood or plasma, in anaphylaxis as an adjunct to epinephrine and other standard measures after the acute symptoms have been controlled, and for other uncomplicated allergic conditions of the immediate type when oral therapy is impossible or contraindicated.

### **Motion sickness**

For active treatment of motion sickness.

### **Antiparkinsonism**

For use in parkinsonism, when oral therapy is impossible or contraindicated, as follows: parkinsonism in the elderly who are unable to tolerate more potent agents; mild cases of parkinsonism in other age groups, and in other cases of parkinsonism in combination with centrally acting anticholinergic agents.

## **CONTRAINDICATIONS**

### **Use in Neonates or Premature Infants**

This drug should *not* be used in neonates or premature infants.

### **Use in Nursing Mothers**

Because of the higher risk of antihistamines for infants generally, and for neonates and prematures in particular, antihistamine therapy is contraindicated in nursing mothers.

### **Use as a Local Anesthetic**

Because of the risk of local necrosis, this drug should not be used as a local anesthetic.

### **Antihistamines are also contraindicated in the following conditions**

Hypersensitivity to diphenhydramine hydrochloride and other antihistamines of similar chemical structure.

## **WARNINGS**

Antihistamines should be used with considerable caution in patients with narrow-angle glaucoma, stenosing peptic ulcer, pyloroduodenal obstruction, symptomatic prostatic hypertrophy, or bladder-neck obstruction.

Local necrosis has been associated with the use of subcutaneous or intradermal use of intravenous diphenhydramine hydrochloride injection.

### **Use in Pediatric Patients**

In pediatric patients, especially, antihistamines in *overdosage* may cause hallucinations, convulsions, or death.

As in adults, antihistamines may diminish mental alertness in pediatric patients. In the young pediatric patient, particularly, they may produce excitation.

## **Use in the Elderly (approximately 60 years or older)**

Antihistamines are more likely to cause dizziness, sedation, and hypotension in elderly patients.

## **PRECAUTIONS**

### **General**

Diphenhydramine hydrochloride has an atropine-like action and, therefore, should be used with caution in patients with a history of bronchial asthma, increased intraocular pressure, hyperthyroidism, cardiovascular disease or hypertension. Use with caution in patients with lower respiratory disease including asthma.

### **Information for Patients**

Patients taking diphenhydramine hydrochloride should be advised that this drug may cause drowsiness and has an additive effect with alcohol.

Patients should be warned about engaging in activities requiring mental alertness such as driving a car or operating appliances, machinery, etc.

### **Drug Interactions**

Diphenhydramine hydrochloride has additive effects with alcohol and other CNS depressants (hypnotics, sedatives, tranquilizers, etc).

MAO inhibitors prolong and intensify the anticholinergic (drying) effects of antihistamines.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Long-term studies in animals to determine mutagenic and carcinogenic potential have not been performed.

### **Pregnancy**

Pregnancy Category B. Reproduction studies have been performed in rats and rabbits at doses up to 5 times the human dose and have revealed no evidence of impaired fertility or harm to the fetus due to diphenhydramine hydrochloride. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

### **Pediatric Use**

Diphenhydramine hydrochloride should not be used in neonates and premature infants (see **CONTRAINDICATIONS**).

Diphenhydramine hydrochloride injection may diminish mental alertness, or, in the young pediatric patient, cause excitation. Overdosage may cause hallucinations, convulsions, or death (see **WARNINGS** and **OVERDOSAGE**).

See also **DOSAGE AND ADMINISTRATION** section.

## **ADVERSE REACTIONS**

**The most frequent adverse reactions are underscored.**

1. *General*: Urticaria, drug rash, anaphylactic shock, photosensitivity, excessive perspiration, chills, dryness of mouth, nose, and throat.
2. *Cardiovascular System*: Hypotension, headache, palpitations, tachycardia, extrasystoles.
3. *Hematologic System*: Hemolytic anemia, thrombocytopenia, agranulocytosis.
4. *Nervous System*: Sedation, sleepiness, dizziness, disturbed coordination, fatigue, confusion, restlessness, excitation, nervousness, tremor, irritability, insomnia, euphoria, paresthesia, blurred vision, diplopia, vertigo, tinnitus, acute labyrinthitis, neuritis, convulsions.
5. *GI System*: Epigastric distress, anorexia, nausea, vomiting, diarrhea, constipation.
6. *GU System*: Urinary frequency, difficult urination, urinary retention, early menses.
7. *Respiratory System*: Thickening of bronchial secretions, tightness of chest or throat and wheezing, nasal stuffiness.

## **OVERDOSAGE**

Antihistamine overdosage reactions may vary from central nervous system depression to stimulation. Stimulation is particularly likely in pediatric patients. Atropine-like signs and symptoms; dry mouth; fixed, dilated pupils; flushing; and gastrointestinal symptoms may also occur.

*Stimulants* should not be used.

Vasopressors may be used to treat hypotension.

## **DOSAGE AND ADMINISTRATION**

THIS PRODUCT IS FOR INTRAVENOUS OR INTRAMUSCULAR ADMINISTRATION ONLY.

Diphenhydramine hydrochloride in the injectable form is indicated when the oral form is impractical.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

DOSAGE SHOULD BE INDIVIDUALIZED ACCORDING TO THE NEEDS AND THE RESPONSE OF THE PATIENT.

### **Pediatric Patients, other than premature infants and neonates**

5 mg/kg/24 hr or 150 mg/m<sup>2</sup>/24 hr. Maximum daily dosage is 300 mg.

Divide into four doses, administered intravenously at a rate generally not exceeding 25 mg/min, or deep intramuscularly.

### **Adults**

10 to 50 mg intravenously at a rate generally not exceeding 25 mg/min, or deep intramuscularly, 100 mg if required; maximum daily dosage is 400 mg.

## **HOW SUPPLIED**

**Diphenhydramine Hydrochloride Injection, USP in parenteral form is supplied as:**

Sterile, pyrogen-free solution containing 50 mg diphenhydramine hydrochloride USP in a 1 mL amber color glass vial with yellow bands and white OPC mark, and available in packages of twenty-five packed and five packed in a PVC rondo tray in a single carton.

Clear, colorless solution, free from any visible particles, no visible leaks or damage to the container closure system filled in tubular USP Type 1 glass vials.

1 mL fill in 2 mL Vial	NDC 72485-101-01
Carton of 25 x 1 mL Vial	NDC 72485-101-25
Carton of 5 x 1 mL Vial	NDC 72485-101-05

**Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].**

**Rx only**

**Protect from freezing and light.**

**Keep covered in carton until time of use.**

**Keep this and all drugs out of the reach of children.**

***HIGH POTENCY***

***Latex Free***

**Manufactured by:**

Micro Labs Limited  
Bangalore-560 099. India

**Distributed by:**

Armas Pharmaceuticals, Inc.  
Freehold, NJ 07728 (USA)

M.L.No.: KTK/28/357/2006

**Rev. 10/2022**

**Packaging**

**INNER VIAL LABEL**


Rx Only NDC 72485-101-01  
**DiphenhydrAMINE**  
 HCl Injection, USP  
**50 mg/mL**  
 HIGH POTENCY Latex Free  
 PROTECT FROM LIGHT  
 IV or DEEP IM  
 1 mL Single-Dose Vial



Code: KR/DRUGS/KTK/28/357/2006  
 USL-ML14-030/B

**OUTER CARTON LABELS**

**PACKAGE- 25 VIALS**



USC-030

Rx Only NDC 72485-101-25  
**DiphenhydrAMINE Hydrochloride**  
 Injection, USP  
**50 mg/mL**  
 HIGH POTENCY Latex Free  
 Sterile, Pyrogen-free  
 For Intravenous or Deep Intramuscular Use  
 25 x 1 mL Single-Dose Vials


Each mL contains diphenhydramine hydrochloride 50 mg in water for injection. The pH of the solution may have been adjusted with either sodium hydroxide or hydrochloric acid.  
 Usual Dosage: See package insert.  
 Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].  
 Protect from freezing and light. Keep covered in carton until time of use.  
 Keep this and all drugs out of the reach of children.

Rx Only NDC 72485-101-25  
**DiphenhydrAMINE Hydrochloride**  
 Injection, USP  
**50 mg/mL**  
 HIGH POTENCY Latex Free  
 Sterile, Pyrogen-free  
 For Intravenous or Deep Intramuscular Use  
 25 x 1 mL Single-Dose Vials

Manufactured by:  
 Micro Labs Limited  
 Bangalore-560 099, India.  
 M.L.No.: KTK/28/357/2006

Distributed by:  
 Armas Pharmaceuticals, Inc.  
 Manalapan, NJ 07726

**PACKAGE - 5 VIALS**



USC-ML14-067

Rx Only NDC 72485-101-05  
**DiphenhydrAMINE Hydrochloride**  
 Injection, USP  
**50 mg/mL**  
 HIGH POTENCY Latex Free  
 Sterile, Pyrogen-free  
 For Intravenous or Deep Intramuscular Use  
 5 x 1 mL Single-Dose Vials

Each mL contains diphenhydramine hydrochloride 50 mg in water for injection. The pH of the solution may have been adjusted with either sodium hydroxide or hydrochloric acid.  
 Usual Dosage: See package insert.  
 Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].  
 Protect from freezing and light. Keep covered in carton until time of use.  
 Keep this and all drugs out of the reach of children.

Rx Only NDC 72485-101-05  
**DiphenhydrAMINE Hydrochloride**  
 Injection, USP  
**50 mg/mL**  
 HIGH POTENCY Latex Free  
 Sterile, Pyrogen-free  
 For Intravenous or Deep Intramuscular Use  
 5 x 1 mL Single-Dose Vials

Manufactured by:  
 Micro Labs Limited  
 Bangalore-560 099, India.  
 Distributed by:  
 Armas Pharmaceuticals, Inc.  
 Freehold, NJ 07728 (USA)  
 M.L.No.: KTK/28/357/2006  
 Rev. 10/2022

# DIPHENHYDRAMINE HYDROCHLORIDE

diphenhydramine hydrochloride injection, solution

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:72485-101
<b>Route of Administration</b>	INTRAMUSCULAR, INTRAVENOUS		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DIPHENHYDRAMINE HYDROCHLORIDE</b> (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	50 mg in 1 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>HYDROCHLORIC ACID</b> (UNII: QTT17582CB)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72485-101-25	25 in 1 CARTON	05/01/2019	
1	NDC:72485-101-01	1 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		
2	NDC:72485-101-05	5 in 1 CARTON	05/05/2023	
2	NDC:72485-101-01	1 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA205723	05/01/2019	

**Labeler** - Armas Pharmaceuticals Inc. (098405973)

Revised: 5/2023

Armas Pharmaceuticals Inc.