

MUPIROCIN- mupirocin ointment

DirectRX

MUPIROCIN

SPL UNCLASSIFIED SECTION

For Dermatologic Use

Rx Only

DESCRIPTION SECTION

Each gram of Mupirocin Ointment USP, 2% contains 20 mg mupirocin in a bland water miscible ointment base (polyethylene glycol ointment, NF) consisting of polyethylene glycol 400 and polyethylene glycol 3350. Mupirocin is a naturally occurring antibiotic. The chemical name is (E)-(2S,3R,4R,5S)-5-[(2S,3S,4S,5S)-2,3-Epoxy-5-hydroxy-4-methylhexyl]tetrahydro-3,4-dihydroxy-β-methyl-2H-pyran-2-crotonic acid, ester with 9-hydroxynonanoic acid. The molecular formula of mupirocin is C₂₆H₄₄O₉ and the molecular weight is 500.62.

CLINICAL PHARMACOLOGY SECTION

- Application of ¹⁴C-labeled mupirocin ointment to the lower arm of normal male subjects followed by occlusion for 24 hours showed no measurable systemic absorption (<1.1 nanogram mupirocin per milliliter of whole blood). Measurable radioactivity was present in the stratum corneum of these subjects 72 hours after application.

Following intravenous or oral administration, mupirocin is rapidly metabolized. The principal metabolite, monic acid, is eliminated by renal excretion, and demonstrates no antibacterial activity. In a study conducted in 7 healthy adult male subjects, the elimination half-life after intravenous administration of mupirocin was 20 to 40 minutes for mupirocin and 30 to 80 minutes for monic acid. The pharmacokinetics of mupirocin has not been studied in individuals with renal insufficiency.

Microbiology -

Mupirocin is an antibacterial agent produced by fermentation using the organism *Pseudomonas fluorescens*. It is active against a wide range of gram-positive bacteria including methicillin-resistant *Staphylococcus aureus* (MRSA). It is also active against certain gram-negative bacteria. Mupirocin inhibits bacterial protein synthesis by reversibly and specifically binding to bacterial isoleucyl transfer-RNA synthetase. Due to this unique mode of action, mupirocin demonstrates no in vitro cross-resistance with other classes of antimicrobial agents.

Resistance occurs rarely. However, when mupirocin resistance does occur, it appears to result from the production of a modified isoleucyl-tRNA synthetase. High-level plasmid-mediated resistance (MIC > 10²⁴ mcg/mL) has been reported in some strains of *S. aureus* and coagulase-negative staphylococci.

Mupirocin is bactericidal at concentrations achieved by topical administration. However, the minimum bactericidal concentration (MBC) against relevant pathogens is generally 8-fold to 30-fold higher than the minimum inhibitory concentration (MIC). In addition, mupirocin is highly protein bound (>97%), and the effect of wound secretions on the MICs of mupirocin has not been determined.

Mupirocin has been shown to be active against most strains of *S. aureus* and *Streptococcus pyogenes*, both in vitro and in clinical studies (see INDICATIONS AND USAGE). The following in vitro data are available, BUT THEIR CLINICAL SIGNIFICANCE IS UNKNOWN. Mupirocin is active against most strains of *Staphylococcus epidermidis* and *Staphylococcus saprophyticus*.

INDICATIONS & USAGE SECTION

Mupirocin Ointment USP, 2% is indicated for the topical treatment of impetigo due to: *S. aureus* and *S. pyogenes*.

CONTRAINDICATIONS SECTION

This drug is contraindicated in individuals with a history of sensitivity reactions to any of its components.

WARNINGS SECTION

Mupirocin Ointment USP, 2% is not for ophthalmic use

PRECAUTIONS SECTION

- If a reaction suggesting sensitivity or chemical irritation should occur with the use of Mupirocin Ointment USP, 2%, treatment should be discontinued and appropriate alternative therapy for the infection instituted.

As with other antibacterial products, prolonged use may result in overgrowth of nonsusceptible organisms, including fungi.

Mupirocin Ointment USP, 2% is not formulated for use on mucosal surfaces. Intranasal use has been associated with isolated reports of stinging and drying. A paraffin-based formulation - *Bactroban Nasal® (mupirocin calcium ointment) – is available for intranasal use.

Polyethylene glycol can be absorbed from open wounds and damaged skin and is excreted by the kidneys. In common with other polyethylene glycol-based ointments, Mupirocin Ointment USP, 2% should not be used in conditions where absorption of large quantities of polyethylene glycol is possible, especially if there is evidence of moderate or severe renal impairment.

Information for Patients -

Use this medication only as directed by your healthcare provider. It is for external use only. Avoid contact with the eyes. The medication should be stopped and your healthcare practitioner contacted if irritation, severe itching, or rash occurs. If impetigo has not improved in 3 to 5 days, contact your healthcare practitioner.

Drug Interactions -

The effect of the concurrent application of Mupirocin Ointment USP, 2% and other drug products has not been studied.

Carcinogenesis, Mutagenesis, Impairment of Fertility -

Long-term studies in animals to evaluate carcinogenic potential of mupirocin have not been conducted.

Results of the following studies performed with mupirocin calcium or mupirocin sodium in vitro and in vivo did not indicate a potential for genotoxicity: Rat primary hepatocyte unscheduled DNA synthesis, sediment analysis for DNA strand breaks, Salmonella reversion test (Ames), Escherichia coli mutation assay, metaphase analysis of human lymphocytes, mouse lymphoma assay, and bone marrow micronuclei assay in mice.

Reproduction studies were performed in male and female rats with mupirocin administered subcutaneously at doses up to 14 times a human topical dose (approximately 60 mg mupirocin per day) on a mg/m² basis and revealed no evidence of impaired fertility and reproductive performance from mupirocin.

Pregnancy:

Teratogenic Effects -

Reproduction studies have been performed in rats and rabbits with mupirocin administered subcutaneously at doses up to 22 and 43 times, respectively, the human topical dose (approximately 60 mg mupirocin per day) on a mg/m² basis and revealed no evidence of harm to the fetus due to

mupirocin. There are however, no adequate and well-controlled studies in pregnant women. Because animal studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Nursing Mothers -

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when Mupirocin Ointment USP, 2% is administered to a nursing woman.

Pediatric Use -

The safety and effectiveness of Mupirocin Ointment USP, 2% have been established in the age range of 2 months to 16 years. Use of mupirocin ointment USP, 2% in these age groups is supported by evidence from adequate and well-controlled studies of mupirocin ointment USP, 2% in impetigo in pediatric patients studied as part of the pivotal clinical trials (see CLINICAL STUDIES).

ADVERSE REACTIONS SECTION

The following local adverse reactions have been reported in connection with the use of mupirocin ointment USP, 2%: Burning, stinging, or pain in 1.5% of patients; itching in 1% of patients; rash, nausea, erythema, dry skin, tenderness, swelling, contact dermatitis, and increased exudate in less than 1% of patients. Systemic reactions to mupirocin ointment USP, 2% have occurred rarely.

DOSAGE & ADMINISTRATION SECTION

A small amount of Mupirocin Ointment USP, 2% should be applied to the affected area 3 times daily. The area treated may be covered with a gauze dressing if desired. Patients not showing a clinical response within 3 to 5 days should be re-evaluated.

CLINICAL STUDIES SECTION

- The efficacy of topical mupirocin ointment USP, 2% in impetigo was tested in two studies. In the first, patients with impetigo were randomized to receive either mupirocin ointment USP, 2% or vehicle placebo three times daily for 8 to 12 days. Clinical efficacy rates at end of therapy in the evaluable populations (adults and pediatric patients included) were 71% for mupirocin ointment USP, 2% (n=49) and 35% for vehicle placebo (n=51). Pathogen eradication rates in the evaluable populations were 94% for mupirocin ointment USP, 2% and 62% for vehicle placebo. There were no side effects reported in the group receiving mupirocin ointment USP, 2%.

In the second study, patients with impetigo were randomized to receive either mupirocin ointment USP, 2% three times daily or 30 to 40 mg/kg oral erythromycin ethylsuccinate per day (this was an unblinded study) for 8 days. There was a follow-up visit 1 week after treatment ended. Clinical efficacy rates at the follow-up visit in the evaluable populations (adults and pediatric patients included) were 93% for mupirocin ointment USP, 2% (n=29) and 78.5% for erythromycin (n=28). Pathogen eradication rates in the evaluable patient populations were 100% for both test groups. There were no side effects reported in the group receiving mupirocin ointment.

Pediatrics -

There were 91 pediatric patients aged 2 months to 15 years in the first study described above. Clinical efficacy rates at end of therapy in the evaluable populations were 78% for mupirocin ointment USP, 2% (n=42) and 36% for vehicle placebo (n=49). In the second study described above, all patients were pediatric except two adults in the group receiving mupirocin ointment USP, 2%. The age range of the pediatric patients was 7 months to 13 years. The clinical efficacy rate for mupirocin ointment USP, 2% (n=27) was 96%, and for erythromycin it was unchanged (78.5%).

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

D

MUPIROCIN OINTMENT
2%/0 22 gm

Generic For: **BACTROBAN**
Each gram contains 20mg mupirocin in a polyethylene glycol ointment, NF base

Lot# **Prod# 203-22** Discard After: 01/17

ACKOE
Caution: Federal law prohibits transfer of this drug to any person other than the patient for whom it was prescribed.
RX ONLY-KEEP OUT OF REACH OF CHILDREN
Dosage: See package insert. Store between 66-77 degrees F

M

NDC 61919-203-22

Alpharetta, GA 30005

DirectRX

Diab By: Perrigo
Allergan, MI 48070
NDC 45802-112-22

Mfg Lot: 5/15/2015

MUPIROCIN OINTMENT 2%
NDC 61919-203-22 22 g
Lot Exp Date 01/17
Mfg NDC 45802-112-22

MUPIROCIN OINTMENT 2%
NDC 61919-203-22 22 g
Lot Exp Date 01/17
Mfg NDC 45802-112-22

MUPIROCIN OINTMENT 2%
NDC 61919-203-22 22 g
Lot Exp Date 01/17
Mfg NDC 45802-112-22

MUPIROCIN

mupirocin ointment

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:61919-203(NDC:45802-112)
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MUPIROCIN (UNII: D0GX8630A5) (MUPIROCIN - UNII:D0GX8630A5)	MUPIROCIN	20 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61919-203-22	22 g in 1 TUBE; Type 0: Not a Combination Product	01/01/2015	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA065123	01/01/2015	

Labeler - DirectRX (079254320)

Establishment

Name	Address	ID/FEI	Business Operations
DirectRX		079254320	repack(6 19 19-203)

Revised: 10/2015

DirectRX