

**RYMED- dexchlorpheniramine maleate and phenylephrine hydrochloride tablet, coated**  
**EDWARDS PHARMACEUTICALS, INC.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**RYMED TABLETS**

**Drug Facts**

<b>Active Ingredients (in each tablet)</b>	<b>Purpose</b>
Dexchlorpheniramine Maleate 2 mg	Antihistamine
Phenylephrine HCl 10 mg	Nasal Decongestant

**Uses**

temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- nasal congestion
- reduces swelling of nasal passages

**Warnings**

- **Do not exceed recommended dosage.**

**Do not use this product**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- heart disease
- high blood pressure
- thyroid disease
- diabetes mellitus
- difficulty in urination due to enlargement of the prostate gland

**Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers**

**When using this product**

- excitability may occur, especially in children
- may cause drowsiness
- alcohol, sedatives and tranquilizers may increase the drowsiness effect
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or are accompanied by a fever
- new symptoms occur

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

**Directions**

**Do not exceed recommended dosage.**

Adults and children 12 years of age and over:	1 tablet every 4 hours, not to exceed 6 tablets in 24 hours, or as directed by a doctor
Children 6 to under 12 years of age:	1/2 tablet every 4 hours, not to exceed 3 tablets in 24 hours, or as directed by a doctor
Children under 6 years of age:	Consult a doctor.

**Inactive ingredients**

Magnesium Stearate, Microcrystalline Cellulose, Sodium Starch Glycolate

**Questions or Comments?**

Call 1-800-543-9560

**PRINCIPAL DISPLAY PANEL - 100 Tablet Bottle Label**

**E**

NDC 0485-0080-01

**RYMED TABLETS**

Antihistamine • Nasal Decongestant

**Each tablet contains:**

Dexchlorpheniramine Maleate 2 mg

Phenylephrine HCl 10 mg

This bottle is not to be dispensed to consumer.

Dispense in a tight, light-resistant container  
with a child-resistant cap.

Store at 59°-86° (15°-30°C) [see USP Controlled Room Temperature].

Tamper evident by foil seal under cap.

Do not use if foil seal is broken or missing.

**Manufactured for:**

***EDWARDS***

**Pharmaceuticals, Inc.**

**Ripley, MS 38663**

100 tablets

## Drug Facts

### Active Ingredients Purpose (in each tablet)

Dexchlorpheniramine  
Maleate 2 mg ..... Antihistamine  
Phenylephrine HCl  
10 mg ..... Nasal Decongestant

**Uses** temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose ■ sneezing ■ itching of the nose or throat ■ itchy, watery eyes
- nasal congestion ■ reduces swelling of nasal passages

Peel Here



NDC 0485-0080-01

## RYMED TABLETS

Antihistamine • Nasal Decongestant

### Each tablet contains:

Dexchlorpheniramine Maleate ..... 2 mg  
Phenylephrine HCl ..... 10 mg

This bottle is not to be dispensed to consumer.  
Dispense in a tight, light-resistant container  
with a child-resistant cap.

Store at 59°-86°F (15°-30°C) [see USP Controlled Room Temperature].  
Tamper evident by foil seal under cap.  
Do not use if foil seal is broken or missing.

Manufactured for:  
**EDWARDS**  
Pharmaceuticals, Inc.  
Ripley, MS 38663

100 tablets



Lot: \_\_\_\_\_  
Exp. Date: \_\_\_\_\_

### Drug Facts (continued)

#### Warnings

■ **Do not exceed recommended dosage.**

#### Do not use this product

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

■ a breathing problem such as emphysema or chronic bronchitis  
■ glaucoma ■ heart disease  
■ high blood pressure ■ thyroid disease ■ diabetes mellitus  
■ difficulty in urination due to enlargement of the prostate gland

#### Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

#### When using this product

■ excitability may occur, especially in children  
■ may cause drowsiness  
■ alcohol, sedatives and tranquilizers may increase the drowsiness effect  
■ avoid alcoholic beverages  
■ use caution when driving a

### Drug Facts (continued)

#### Stop use and ask a doctor if

■ nervousness, dizziness, or sleeplessness occur  
■ symptoms do not improve within 7 days or are accompanied by a fever ■ new symptoms occur

**If pregnant or breast-feeding**, ask a health professional before use.

#### Keep out of reach of children.

In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

#### Directions

**Do not exceed recommended dosage.**

Adults and children 12 years of age and over:	1 tablet every 4 hours, not to exceed 6 tablets in 24 hours, or as directed by a doctor
Children 6 to under 12 years of age:	1/2 tablet every 4 hours, not to exceed 3 tablets in 24 hours, or as directed by a doctor
Children under 6 years of age:	Consult a doctor.

#### Inactive ingredients

Magnesium Stearate,  
Microcrystalline Cellulose, Sodium Starch Glycolate

#### Questions or Comments?

**RYMED**

dexchlorpheniramine maleate and phenylephrine hydrochloride tablet, coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0485-0080
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>DEXCHLORPHENIRAMINE MALEATE</b> (UNII: B10YD955QW) (DEXCHLORPHENIRAMINE - UNII:3Q9Q0B929N)	DEXCHLORPHENIRAMINE MALEATE	2 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>SODIUM STARCH GLYCOLATE TYPE A CORN</b> (UNII: AG9B65PV6B)	

**Product Characteristics**

<b>Color</b>	white	<b>Score</b>	2 pieces
<b>Shape</b>	OVAL	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	R;1
<b>Contains</b>			

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:0485-0080-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/15/2011	

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph final	part341	11/15/2011	

**Labeler** - EDWARDS PHARMACEUTICALS, INC. (195118880)

Revised: 8/2023

EDWARDS PHARMACEUTICALS, INC.