FOSTER AND THRIVE DAYTIME COLD AND FLU RELIEF NIGHTTIME COLD AND FLU RELIEF- acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl
Strategic Sourcing Services LLC

McKesson Nighttime & Daytime Cold & Flu Drug Facts

Active ingredients (in each softgel) – Nighttime Cold & Flu

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

Active ingredients (in each softgel) – Daytime Cold & Flu

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Purpose – Nighttime Cold & Flu

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Purpose – Daytime Cold & Flu

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Uses – Nighttime Cold & Flu

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

Uses – Daytime Cold & Flu

temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson’s disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have – Nighttime Cold & Flu

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough as occurs with smoking, asthma, or emphysema

- trouble urinating due to an enlarged prostate gland

Ask a doctor before use if you have – Daytime Cold & Flu

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are – Nighttime Cold & Flu

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

Ask a doctor or pharmacist before use if you are – Daytime Cold & Flu

taking the blood thinning drug warfarin

When using this product – Nighttime Cold & Flu

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

When using the product – Daytime Cold & Flu

do not use more than directed

Stop use and ask a doctor if – Nighttime Cold & Flu

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Stop use and ask a doctor if – Daytime Cold & Flu

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions – Nighttime Cold & Flu

- take only as directed – see overdose warning
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	2 softgels with water every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Directions – Daytime Cold & Flu

- take only as directed – see overdose warning
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	2 softgels with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients – Nighttime Cold & Flu

D&C yellow no. 10, edible ink*, FD&C blue no. 1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution *may contain this ingredient

Inactive ingredients – Daytime Cold & Flu

edible ink*, FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution *may contain this ingredient

Questions or comments?

Call **833-358-6431** Monday to Friday 9:00am to 7:00pm EST

Package/Label Principal Display Panel

COMPARE TO VICKS® DAYQUIL® COLD & FLU ACTIVE INGREDIENTS

Foster & Thrive™

DAYTIME

Cold & Flu Relief

Acetaminophen

Dextromethorphan HBr

Phenylephrine HCl

PAIN RELIEVER, FEVER REDUCER, COUGH SUPPRESSANT, NASAL DECONGESTANT

Alcohol Free | Antihistamine Free | Gluten Free

Non-Drowsy

Multi-Symptom Relief of:

• Aches/Fever

• Sore Throat

• Nasal Congestion

• Cough

• Headache

32 SOFTGELS

ACTUAL SIZE

COMPARE TO VICKS® NYQUIL® COLD & FLU ACTIVE INGREDIENTS

Foster & Thrive™

NIGHTTIME

Cold & Flu Relief

Acetaminophen

Dextromethorphan HBr

Doxylamine succinate

PAIN RELIEVER, FEVER REDUCER, COUGH SUPPRESSANT, ANTIHISTAMINE

Alcohol Free | Gluten Free

Multi-Symptom Relief of:

• Aches/Fever

• Sore Throat

- Sneezing
- Runny Nose
- Cough
- Headache

16 SOFTGELS

ACTUAL SIZE

Nighttime Cold & Flu

Drug Facts

Active ingredients (in each softgel)	*Purpose*
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 15 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Antihistamine

Uses temporarily relieves common cold/flu symptoms:
- cough due to minor throat and bronchial irritation
- minor aches and pains
- sore throat
- headache
- runny nose and sneezing
- fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away. ▼

Drug Facts (continued)

Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are
- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if
- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. ▼

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
- take only as directed – see overdose warning
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	2 softgels with water every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information ■ store at 20-25°C (68-77°F)

Inactive ingredients D&C yellow no. 10, edible ink*, FD&C blue no. 1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution *may contain this ingredient

Questions or comments? Call 833-380-6431 Monday to Friday 9:00am to 7:00pm EST

Daytime Cold & Flu

Drug Facts

Active ingredients (in each softgel)	*Purpose*
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms:
- cough due to minor throat and bronchial irritation
- minor aches and pains
- sore throat
- headache
- nasal congestion
- fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away. ▼

Drug Facts (continued)

Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
- liver disease
- diabetes
- heart disease
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if
- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. ▼

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
- take only as directed – see overdose warning
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	2 softgels with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information ■ store at 20-25°C (68-77°F)

Inactive ingredients edible ink*, FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution *may contain this ingredient

Questions or comments? Call 833-380-6431 Monday to Friday 9:00am to 7:00pm EST

DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND INFORMATION.

**These products are not manufactured or distributed by Procter & Gamble, distributor of Vicks® DayQuil® Cold & Flu and Vicks® NyQuil® Cold & Flu.

PARENTS: Learn about teen medicine abuse www.StopMedicineAbuse.org

Distributed by: McKesson Corp. via Strategic Sourcing Services LLC. Memphis, TN 38141

Money Back Guarantee

Product of Canada

REV 03/23

CONVENIENT RECLOSING TAB

OPEN OTHER END

0 10939 95547 0

58072 KM C1

NDC 70677-1028-1

COMPARE TO VICKS® DAYQUIL® COLD & FLU ACTIVE INGREDIENTS**

Foster & Thrive™

DAYTIME Cold & Flu Relief

Acetaminophen
Dextromethorphan HBr
Phenylephrine HCl

PAIN RELIEVER, FEVER REDUCER, COUGH SUPPRESSANT, NASAL DECONGESTANT

Alcohol Free | Antihistamine Free | Gluten Free
Non-Drowsy

Multi-Symptom Relief of:
- Aches/Fever
- Sore Throat
- Nasal Congestion
- Cough
- Headache

32 SOFTGELS ACTUAL SIZE

COMPARE TO VICKS® NYQUIL® COLD & FLU ACTIVE INGREDIENTS**

Foster & Thrive™

NIGHTTIME Cold & Flu Relief

Acetaminophen
Dextromethorphan HBr
Doxylamine succinate

PAIN RELIEVER, FEVER REDUCER, COUGH SUPPRESSANT, ANTIHISTAMINE

Alcohol Free | Gluten Free

Multi-Symptom Relief of:
- Aches/Fever
- Sore Throat
- Sneezing
- Runny Nose
- Cough
- Headache

16 SOFTGELS ACTUAL SIZE

FOSTER AND THRIVE DAYTIME COLD AND FLU RELIEF NIGHTTIME COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl kit

Product Information			
Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:70677-1028

Packaging				
#	**Item Code**	**Package Description**	**Marketing Start Date**	**Marketing End Date**
1	NDC:70677-1028-1	1 in 1 CARTON; Type 0: Not a Combination Product	10/30/2023	

Quantity of Parts		
Part #	**Package Quantity**	**Total Product Quantity**
Part 1	8 BLISTER PACK	16
Part 2	16 BLISTER PACK	32

Part 1 of 2

FOSTER AND THRIVE NIGHTTIME COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

Product Information	
Item Code (Source)	NDC:0113-1000
Route of Administration	ORAL

Active Ingredient/Active Moiety		
Ingredient Name	**Basis of Strength**	**Strength**
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients	
Ingredient Name	**Strength**
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	

POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics			
Color	GREEN (clear)	**Score**	no score
Shape	OVAL	**Size**	20mm
Flavor		**Imprint Code**	056
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0113-1000-00	8 in 1 CARTON		
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012		

Part 2 of 2

FOSTER AND THRIVE DAYTIME COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule, liquid filled

Product Information	
Item Code (Source)	NDC:0113-2101
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics

Color	ORANGE	**Score**	no score
Shape	OVAL	**Size**	20mm
Flavor		**Imprint Code**	L994
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0113-2101-00	16 in 1 CARTON		
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/30/2023	

Labeler - Strategic Sourcing Services LLC (116956644)

Revised: 11/2023 Strategic Sourcing Services LLC