

**IBUPROFEN- ibuprofen capsule, liquid filled**  
**CVS Pharmacy**

-----

**DRUG FACTS**

**Active ingredient (in each capsule)**

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\*

(present as the free acid and potassium salt)

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/ fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - minor pain of arthritis
  - toothache
  - backache
  - the common cold
  - menstrual cramps
  - temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease or asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery

**Keep out of reach of children**

In case of overdose, get medical help or contact a Poison Control Center right away right away.

**Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over:
  - take 1 capsule every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 capsule, 2 capsules may be used
  - do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**

- each capsule contains: **potassium 20 mg**
- read all warnings and directions before use.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F). Protect from light.
- do not chew, crush, or dissolve capsule before swallowing

### **Inactive ingredients**

FD&C blue #1, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol

### **Questions or Comments ?**

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

### **Principal Display Panel**

Compare to the active ingredient in Advil® Liqui-Gels® \*\*

Liquid Filled Softgels†

### **Ibuprofen**

LIQUID FILLED SOFTGELS†

IBUPROFEN CAPSULES, 200 mg

Pain reliever;

Fever reducer (NSAID)

SOFTGELS†

†Liquid filled capsules

\*\*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liqui-Gels®.


**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

**Distributed by: CVS Pharmacy, Inc.**

One CVS Drive, Woonsocket, RI 02895

CVS.com

### **Product Label**



Compare to the active ingredient in Advil® Liqui-Gels®\*\*

**Liquid Filled Softgels!**


# Ibuprofen

LIQUID FILLED SOFTGELS\*  
IBUPROFEN CAPSULES, 200 mg

**Pain reliever;  
Fever reducer (NSAID)\***

Actual Product Size on Back Panel

**200 SOFTGELS\***  
\*Liquid filled capsules



TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

**Drug Facts**

**Active ingredient (in each capsule)**  
Sulfobenzothiazone equal to 200 mg ibuprofen (NSAID)† ..... Pain reliever/fever reducer  
†nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - muscle aches
  - minor pain of arthritis
  - toothache
- headache
- the common cold
- menstrual cramps
- temporarily relieves redness/itching

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms include: hives, skin rash, facial swelling, difficulty breathing, dizziness, fainting, or loss of consciousness.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug (corticosteroid), or other (aspirin, clozapine, or other) medication
- take other drugs containing prescription or nonprescription NSAIDs, aspirin, ibuprofen, naproxen, or other
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed


**Drug Facts (continued under label)**

This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liqui-Gels®.

**Distributed by:** CVS Pharmacy, Inc.  
One CVS Drive, Woonsocket, RI 02896  
CVS Retail Stores  
CVS.com™ 1-800-SHOP CVS  
V-31884

**CVS Quality**  
Money Back Guarantee

#213638



0 50428 56929 0

Actual Size

Lot No.:  
Exp. Date:

RJ0-AJZ 1004491  
Ref# 0312

PEEL HERE

**Drug Facts (continued)**

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease or asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - vomit bloody or black stools
  - have stomach pain that does not get better
  - stool gets worse or lasts more than 10 days
  - stool gets worse or lasts more than 3 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present in the painful area
  - any new symptoms appear

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use ibuprofen during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over:
  - take 1 capsule every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 capsule, 2 capsules may be used
  - do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**

- each capsule contains: potassium 20 mg
- read all warnings and directions before use
- store at 20° to 25° (68° to 77°F)
- avoid excessive heat above 40°C (104°F). Protect from light.
- do not chew, crush, or dissolve capsule before swallowing
- do not drink, crush, or dissolve capsule before swallowing

**Inactive ingredients** FD&C Blue #1, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol

**Questions or comments?** Call 1-877-755-3838 Monday-Friday 9AM-5PM EST

**CVS HEALTH Ibuprofen Softgels**

<b>IBUPROFEN</b>			
ibuprofen capsule, liquid filled			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:59779-197
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			
<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>	
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg	
<b>Inactive Ingredients</b>			
<b>Ingredient Name</b>	<b>Strength</b>		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)			
GELATIN (UNII: 2G86QN327L)			
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)			
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)			
WATER (UNII: 059QF0K00R)			

<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	

### Product Characteristics

<b>Color</b>	BLUE	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	IB200
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59779-197-88	1 in 1 BOX	09/24/2012	
1		88 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:59779-197-20	1 in 1 BOX	09/24/2012	
2		20 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:59779-197-16	2 in 1 BOX	09/24/2012	
3		80 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:59779-197-48	1 in 1 BOX	09/24/2012	
4		48 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:59779-197-80	1 in 1 BOX	09/24/2012	
5		80 in 1 BOTTLE; Type 0: Not a Combination Product		
6	NDC:59779-197-03	1 in 1 BOX	09/24/2012	
6		300 in 1 BOTTLE; Type 0: Not a Combination Product		
7	NDC:59779-197-40	1 in 1 BOX	09/24/2012	
7		40 in 1 BOTTLE; Type 0: Not a Combination Product		
8	NDC:59779-197-02	200 in 1 BOTTLE; Type 0: Not a Combination Product	09/24/2012	
9	NDC:59779-197-12	120 in 1 BOTTLE; Type 0: Not a Combination Product	09/24/2012	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078682	09/24/2012	

**Labeler** - CVS Pharmacy (062312574)

Revised: 10/2016

CVS Pharmacy