

LOW DOSE ASPIRIN- aspirin 81 mg delayed release tablets tablet WALGREENS

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

481R WALGREENS 0363 0690 Aspirin 81 mg Tablet

DRUG FACTS

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**
- ask your doctor about other uses for enteric-coated 81 mg Aspirin

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problem
- take a blood thinning (anticoagulant) or steroid drug
- take other drug containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Ask a doctor or pharmacist before use if you are

taking a prescription drug for:

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

an allergic reaction occurs. Seek medical help right away.

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days

redness or swelling is present

new symptoms occur

ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

Store at room temperature. Avoid excessive heat above 40°C (104°F).

Inactive ingredients anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide,

triethyl citrate

Questions or comments? Call 1-877-290-4008

NDC 0363-0690-31

Walgreens

Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose

Enteric-Coated

300
TABLETS

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredient (in each tablet)
Aspirin 81 mg (NSAID)*

Purpose
Pain reliever

Nonsteroidal anti-inflammatory drug

Uses ■ for the temporary relief of minor aches and pains or as recommended by your doctor.

Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

■ ask your doctor about other uses for enteric-coated 81 mg Aspirin.

Warnings (See carton for complete warnings information)

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

■ are age 60 or older ■ have had stomach ulcers or bleeding problems

■ take a blood thinning (anticoagulant) or steroid drug

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

■ have 3 or more alcoholic drinks every day while using this product

■ take more or for a longer time than directed

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ drink a full glass of water with each dose

■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: consult a doctor

Other information TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

■ Store at room temperature. Avoid excessive heat above 40°C (104°F)

DISTRIBUTED BY: WALGREEN CO., 200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEED walgreens.com ©2022 Walgreen Co.

MADE IN U.S.A. OF U.S. AND IMPORTED INGREDIENTS ITEM 285642 W00000-0000-0

481R 0222 Lot No. Exp. Date:

Varnish Omit Area

W30B90422-F

Lot No.:
Exp Date:



ITEM 2856942 W00000-0000-0000-0000-0000

TAMPER EVIDENT: DO NOT OPEN THIS PRODUCT IF THE IMPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.

Drug Facts

Active ingredient
(in each tablet)
Aspirin 81 mg (NSAID)*.....Pain reliever/
*nonsteroidal anti-inflammatory drug

Purpose

REMAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Uses

- for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**
- ask your doctor about other uses for enteric coated 81 mg Aspirin.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Drug Facts (continued)

Have 3 or more alcoholic drinks every day while using this product.

Take more or for a longer time than directed.

Do not use

- if you are allergic to aspirin or any other pain reliever/reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug for:

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- ever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-2222) right away.

Directions

- drink a full glass of water with each dose.
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

- Store at room temperature. Avoid excessive heat above 40°C (104°F).

Inactive ingredients

amylaceous lactose, carmelum wax, colloidal silicon dioxide, croscarmellose sodium, D&B yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyborate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

Questions or comments?
Call 1-877-290-4008

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Low Dose Aspirin.

DISTRIBUTED BY: WALGREENS CO.
200 W. LAKE RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
WALGREENS STORES AND PHARMACIES
WALGREENS, WALGREENS.COM
© 2015 WALGREENS CO.
IMPORTED INGREDIENTS

NDC 0363-0690-31



Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose Enteric-Coated

300
TABLETS

ACTUAL SIZE
**Talk to your doctor or other healthcare provider before using this product for your heart.

W30RGM422-F



PAPER BOX
PLASTIC BOTTLE

Walgreens

NDC 0363-0690-20

Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose

Enteric-Coated

200 TABLETS

Drug Facts

Active ingredient (in each tablet)
Aspirin 81 mg (NSAID)*
nonsteroidal anti-inflammatory drug

Purpose
Pain reliever

Uses ■ for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief. ■ ask your doctor about other uses for enteric-coated 81 mg Aspirin.

Warnings (See carton for complete warnings information)
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ take more or for a longer time than directed

■ have 3 or more alcoholic drinks every day while using this product

Do not use
■ if pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ drink a full glass of water with each dose
■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: consult a doctor

Other information
TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING. ■ Store at room temperature. Avoid excessive heat above 40°C (104°F).
W3OR0422-F

DISTRIBUTED BY: WALGREEN CO., 200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED walgreens.com ©2022 Walgreen Co.
MADE IN U.S.A. OF U.S.A. AND IMPORTED INGREDIENTS ITEM 284614 W00000-0000-0
481R 0222

Lot No.:
Exp. Date:

Varnish Omit Area



481R 0222

Our pharmacists recommend the Walgreens Brand Bayer® Low Dose Aspirin. We invite you to compare to national brands. This product is not manufactured or distributed by Bayer AG, owner of the registered trademark.

Drug Facts

Active ingredient (in each tablet)
Aspirin 81 mg (NSAID)*
nonsteroidal anti-inflammatory drug

Purpose
Pain reliever

Uses ■ for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief. ■ ask your doctor about other uses for enteric coated 81 mg Aspirin.

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product

■ take more or for a longer time than directed

Do not use
■ if you are allergic to aspirin or any other pain reliever/fever reducer
■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if
■ you have asthma ■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn

Drug Facts (continued)

■ you are taking a diuretic
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if
■ an allergic reaction occurs. Seek medical help right away.
■ you experience any of the following signs of stomach bleeding: ● feel faint ● vomit blood ● have bloody or black stools

■ have stomach pain that does not get better
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ new symptoms occur
■ ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions
■ drink a full glass of water with each dose
■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
■ children under 12 years: consult a doctor

Other information
■ Store at room temperature. Avoid excessive heat above 40°C (104°F)

Inactive ingredients anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, stearic, talc, titanium dioxide, triethyl citrate

Questions or comments? Call 1-877-290-4008

Walgreens
Compare to the active ingredient in Bayer® Low Dose Aspirin®
Aspirin 81
81 mg / PAIN RELIEVER (NSAID)
Low Dose Enteric-Coated
200 TABLETS

HEART HEALTH
**Talk to your doctor or other healthcare provider before using this product for your heart.

Walgreens
Compare to the active ingredient in Bayer® Low Dose Aspirin®
Aspirin 81
81 mg / PAIN RELIEVER (NSAID)
Low Dose Enteric-Coated

HEART HEALTH
200 TABLETS
ACTUAL SIZE
**Talk to your doctor or other healthcare provider before using this product for your heart.

Drug Facts (continued)

■ you are taking a diuretic
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if
■ an allergic reaction occurs. Seek medical help right away.
■ you experience any of the following signs of stomach bleeding: ● feel faint ● vomit blood ● have bloody or black stools

■ have stomach pain that does not get better
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ new symptoms occur
■ ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions
■ drink a full glass of water with each dose
■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
■ children under 12 years: consult a doctor

Other information
■ Store at room temperature. Avoid excessive heat above 40°C (104°F)

Inactive ingredients anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, stearic, talc, titanium dioxide, triethyl citrate

Questions or comments? Call 1-877-290-4008

W3OR0422-F

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

ITEM 284614 W00000-0000-0



Lot No.:
Exp. Date:

NDC 0363-0690-12

Walgreens

Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose

Enteric-Coated

120 TABLETS

Drug Facts

Active ingredient (in each tablet)
Aspirin 81 mg (NSAID)
nonsteroidal anti-inflammatory drug

Uses ■ for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.** ■ ask your doctor about other uses or enteric-coated 81 mg Aspirin.

Warnings (See carton for complete warnings information)
Reye's syndrome. Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Directions
■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: consult a doctor

Other information TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING ■ Store at room temperature. Avoid excessive heat above 40°C (104°F).

DISTRIBUTED BY: WALGREEN CO., 481R 0222
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walmart.com ©2022 Walgreen Co.
W00000-0000-0
W30RG0422-F
MADE IN U.S.A. OF U.S. AND IMPORTED INGREDIENTS

Lot No.:
Exp. Date:

Varnish Omit Area

Drug Facts (continued)
Other information
Store at room temperature. Avoid excessive heat above 40°C (104°F).

Drug Facts (continued)
Directions
■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: consult a doctor

Other information TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING ■ Store at room temperature. Avoid excessive heat above 40°C (104°F).

Lot No.:
Exp. Date:

Varnish Omit Area



Low Dose
Enteric-Coated

Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

240 TABLETS

3 BOTTLES:
120 TABLETS PER BOTTLE

ACTUAL SIZE
before using this product for your heart.

***Talk to your doctor or other healthcare provider before using this product for your heart.

Drug Facts (continued)
Do not use
■ this product ■ take more or for a longer time than directed

Drug Facts (continued)
Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Other information TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING ■ Store at room temperature. Avoid excessive heat above 40°C (104°F).

Lot No.:
Exp. Date:

Varnish Omit Area

Walgreens

Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

how2recycle.info



TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Low Dose Aspirin, 481R 0222

DISTRIBUTED BY: WALGREEN CO., 481R 0222
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walmart.com ©2022 Walgreen Co.
W00000-0000-0
W30RG0422-F
MADE IN U.S.A. OF U.S. AND IMPORTED INGREDIENTS



Lot No.:
Exp. Date:

Varnish Omit Area

TWIN PACK

NDC 0363-0690-29

Walgreens



Compare to the active ingredient in Bayer® Low Dose Aspirin®

Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose

Enteric-Coated



240
TABLETS

(2 BOTTLES:
120 TABLETS PER BOTTLE)

ACTUAL SIZE

**Talk to your doctor or other healthcare provider before using this product for your heart.

NDC 0363-0690-32

Walgreens

Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose

Enteric-Coated

32 TABLETS

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID) Pain reliever
nonsteroidal anti-inflammatory drug

Purpose
Pain reliever

Uses ■ for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.** ■ ask your doctor about other uses for enteric-coated 81 mg Aspirin.

Warnings (See carton for complete warnings information)
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: consult a doctor

Other Information TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING ■ Store at room temperature. Avoid excessive heat above 40°C (104°F).
DISTRIBUTED BY: WALGREEN CO. 481 F 0222
200 WILMOT RD., DEERFIELD, IL 60015 ITEM 284947
W00000-0000-0 W00000-0000-0
100% SATISFACTION GUARANTEED
Walgreens.com ©2022 Walgreen Co. WSGORG042Z-F
MADE IN U.S.A. OF U.S. AND IMPORTED INGREDIENTS

Lot No.:

Exp. Date:

Varnish Omit Area



ITEM 284947 W00000-0000-0
 3 111917 13644 8

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts
 (in each tablet)

Active ingredient
 Aspirin 81 mg (NSAID) *.....Pain reliever
 *nonsteroidal anti-inflammatory drug

Purpose
 Uses ■ for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.
 ■ ask your doctor about other uses for enteric coated 81 mg Aspirin.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction, which may include:
 ■ hives ■ facial swelling
 ■ asthma (wheezing) ■ shock
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

Drug Facts (continued)

■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 ■ have 3 or more alcoholic drinks every day while using this product
 ■ take more or for a longer time than directed
Do not use
 ■ if you are allergic to aspirin or any other pain reliever/fever reducer
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

■ you have asthma
 ■ stomach bleeding warning applies to you
 ■ you have a history of stomach problems, such as heartburn ■ you are taking a diuretic
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug for

■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if

■ an allergic reaction occurs. Seek medical help right away.
 ■ you experience any of the following signs of stomach bleeding: ● feel faint ● vomit blood ● have bloody or black stools
 ■ have stomach pain that does not get better
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days

Drug Facts (continued)

■ redness or swelling is present
 ■ new symptoms occur
 ■ ringing in the ears or a loss of hearing occurs
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

■ drink a full glass of water with each dose
 ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
 ■ children under 12 years: consult a doctor

Other information

■ Store at room temperature. Avoid excessive heat above 40°C (104°F)

Inactive ingredients

anhydrous lactose, carmalum wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyosorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

NDC 0363-0690-32



Walgreens
 Compare to the active ingredient in Bayer® Low Dose Aspirin**



Aspirin 81
 81 mg / PAIN RELIEVER (NSAID)

Low Dose Enteric-Coated

32 TABLETS

ACTUAL SIZE

**Talk to your doctor or other healthcare provider before using this product for your heart.

W3ORG0422-F

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Low Dose Aspirin.
 DISTRIBUTED BY: WALGREEN CO.
 200 WILMOT RD., DEERFIELD, IL 60015
 100% SATISFACTION GUARANTEED
 481R 0222 walgreens.com ©2022 Walgreen Co.

Drug Facts (continued)
 Questions or comments?
 Call 1-877-290-4008

NDC 0363-0690-12

Walgreens

Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose

Enteric-Coated

120 TABLETS

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Uses ■ for the temporary relief of minor aches and pains, or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.** ■ ask your doctor about other uses for enteric-coated 81 mg Aspirin

Warnings (See carton for complete warnings information)
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert:** Aspirin may cause a severe allergic reaction, which may include: ■ shock ■ asthma (wheezing) ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: consult a doctor

Other information TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING ■ Store at room temperature. Avoid excessive heat above 40°C (104°F)

Questions or comments? Call 1-877-290-4008
Walgreens Pharmacist Recommended. Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
DISTRIBUTED BY: WALGREEN CO. ITEM 757871
200 WILMOT RD., DEERFIELD, IL 60015 W00000-0000-0
100% SATISFACTION GUARANTEED 481R 0921
walgreens.com ©2021 Walgreen Co. W3ORG1221-F

Lot No.:
Exp. Date:

Varnish Omit Area

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredient (in each tablet) Purpose
Aspirin 81 mg (NSAID)* Pain reliever
*nonsteroidal anti-inflammatory drug

Uses
■ for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**
■ ask your doctor about other uses for enteric-coated 81 mg Aspirin

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older

Drug Facts (continued)

■ have had stomach ulcers or bleeding problem
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed
Do not use
■ if you are allergic to aspirin or any other pain reliever/fever reducer
■ if you have ever had an allergic reaction to this product or any of its ingredients
Ask a doctor before use if
■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
■ you are taking a diuretic ■ you have asthma
Ask a doctor or pharmacist before use if you are taking a prescription drug for:
■ gout ■ diabetes ■ arthritis
Stop use and ask a doctor if
■ an allergic reaction occurs. Seek medical help right away.
■ you experience any of the following signs of stomach bleeding: ■ feel faint

Drug Facts (continued)
Inactive ingredients: carboxymethylcellulose, croscarmellose sodium, D&C Yellow #10, aluminum lake, iron oxide other, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyvinylpyrrolidone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate
Questions or comments? Call 1-877-290-4008
©2021 Walgreen Co. All rights reserved. This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer®.

Drug Facts (continued)

■ have bloody or black stools ■ vomit blood
■ have stomach pain that does not get better
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ new symptoms occur
■ ringing in the ears or a loss of hearing occurs
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions
■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: consult a doctor

Other information
■ Store at room temperature. Avoid excessive heat above 40°C (104°F).

Coating/Print Omit Area
Lot No.:
Exp Date:
3 11917 01558 3
ITEM 757871 W00000-0000-0
2021 Walgreen Co.
walgreens.com 1-877-290-4008
100% SATISFACTION GUARANTEED
200 WILMOT RD., DEERFIELD, IL 60015
DISTRIBUTED BY: WALGREEN CO.
481R 0921

NDC 0363-0690-12

HEART HEALTH

Compare to the active ingredient in Bayer® Low Dose Aspirin™

Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose Enteric-Coated

120 TABLETS

ACTUAL SIZE

*Talk to your doctor or other healthcare provider before using this product for your heart.



LOW DOSE ASPIRIN

aspirin 81 mg delayed release tablets tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0690
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
TALC (UNII: 7SEV7J4R1U)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STARCH, CORN (UNII: O8232NY3SJ)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	Heart
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0690-12	1 in 1 CARTON	10/08/2021	
1		120 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0363-0690-32	1 in 1 CARTON	06/15/2022	
2		32 in 1 BOTTLE; Type 0: Not a Combination Product		

3	NDC:0363-0690-29	2 in 1 CARTON	06/15/2022	
3		120 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:0363-0690-20	1 in 1 CARTON	06/15/2022	
4		200 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:0363-0690-31	1 in 1 CARTON	06/15/2022	
5		300 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	10/08/2021	

Labeler - WALGREENS (008965063)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
TIME CAP LABORATORIES, INC.		037052099	manufacture(0363-0690)

Revised: 5/2022

WALGREENS