

# **HYDRALAZINE HYDROCHLORIDE- hydralazine hydrochloride tablet**

## **Aphena Pharma Solutions - Tennessee, LLC**


-----

### **HydrALAZINE Hydrochloride**

#### **Tablets USP**

#### **DESCRIPTION**

Hydralazine Hydrochloride tablets USP, is an antihypertensive, for oral administration. Its chemical name is 1-hydrazinophthalazine monohydrochloride, and its structural formula is:



$C_8H_8N_4 \cdot HCl$  M.W. 196.64

Hydralazine Hydrochloride USP is a white to off-white, odorless crystalline powder. It is soluble in water, slightly soluble in alcohol. It melts at about 275°C, with decomposition. Each tablet for oral administration contains 10 mg, 25 mg, 50 mg or 100 mg hydralazine hydrochloride, USP. Tablets also contain anhydrous lactose, FD&C yellow no. 6 aluminum lake, microcrystalline cellulose, sodium starch glycolate, and stearic acid.

#### **CLINICAL PHARMACOLOGY**

Although the precise mechanism of action of hydralazine is not fully understood, the major effects are on the cardiovascular system.

Hydralazine apparently lowers blood pressure by exerting a peripheral vasodilating effect through a direct relaxation of vascular smooth muscle. Hydralazine, by altering cellular calcium metabolism, interferes with the calcium movements within the vascular smooth muscle that are responsible for initiating or maintaining the contractile state.

The peripheral vasodilating effect of hydralazine results in decreased arterial blood pressure (diastolic more than systolic); decreased peripheral vascular resistance; and an increased heart rate, stroke volume, and cardiac output. The preferential dilatation of arterioles, as compared to veins, minimizes postural hypotension and promotes the increase in cardiac output. Hydralazine usually increases renin activity in plasma, presumably as a result of increased secretion of renin by the renal juxtaglomerular cells in response to reflex sympathetic discharge. This increase in renin activity leads to the production of angiotensin II, which then causes stimulation of aldosterone and consequent sodium reabsorption. Hydralazine also maintains or increases renal and

cerebral blood flow.

Hydralazine is rapidly absorbed after oral administration, and peak plasma levels are reached at 1 to 2 hours. Plasma levels of apparent hydralazine decline with a half-life of 3 to 7 hours. Binding to human plasma protein is 87%. Plasma levels of hydralazine vary widely among individuals. Hydralazine is subject to polymorphic acetylation; slow acetylators generally have higher plasma levels of hydralazine and require lower doses to maintain control of blood pressure. Hydralazine undergoes extensive hepatic metabolism; it is excreted mainly in the form of metabolites in the urine.

## **INDICATIONS AND USAGE**

Essential hypertension, alone or as an adjunct.

## **CONTRAINDICATIONS**

Hypersensitivity to hydralazine; coronary artery disease; mitral valvular rheumatic heart disease.

## **WARNINGS**

In a few patients hydralazine may produce a clinical picture simulating systemic lupus erythematosus including glomerulonephritis. In such patients hydralazine should be discontinued unless the benefit-to-risk determination requires continued antihypertensive therapy with this drug. Symptoms and signs usually regress when the drug is discontinued but residua have been detected many years later. Long-term treatment with steroids may be necessary. (See **PRECAUTIONS, Laboratory Tests.**)

## **PRECAUTIONS**

### **General**

Myocardial stimulation produced by hydralazine can cause anginal attacks and ECG changes of myocardial ischemia. The drug has been implicated in the production of myocardial infarction. It must, therefore, be used with caution in patients with suspected coronary artery disease.

The "hyperdynamic" circulation caused by hydralazine may accentuate specific cardiovascular inadequacies. For example, hydralazine may increase pulmonary artery pressure in patients with mitral valvular disease. The drug may reduce the pressor responses to epinephrine. Postural hypotension may result from hydralazine but is less common than with ganglionic blocking agents. It should be used with caution in patients with cerebral vascular accidents.

In hypertensive patients with normal kidneys who are treated with hydralazine, there is evidence of increased renal blood flow and a maintenance of glomerular filtration rate. In some instances where control values were below normal, improved renal function has been noted after administration of hydralazine. However, as with any antihypertensive agent, hydralazine should be used with caution in patients with advanced renal damage.

Peripheral neuritis, evidenced by paresthesia, numbness, and tingling, has been observed. Published evidence suggests an antipyridoxine effect, and that pyridoxine should be added to the regimen if symptoms develop.

### **Information for Patients**

Patients should be informed of possible side effects and advised to take the medication regularly and continuously as directed.

### **Laboratory Tests**

Complete blood counts and antinuclear antibody titer determinations are indicated before and periodically during prolonged therapy with hydralazine even though the patient is asymptomatic. These studies are also indicated if the patient develops arthralgia, fever, chest pain, continued malaise, or other unexplained signs or symptoms.

A positive antinuclear antibody titer requires that the physician carefully weigh the implications of the test results against the benefits to be derived from antihypertensive therapy with hydralazine.

Blood dyscrasias, consisting of reduction in hemoglobin and red cell count, leukopenia, agranulocytosis, and purpura, have been reported. If such abnormalities develop, therapy should be discontinued.

### **Drug/Drug Interactions**

MAO inhibitors should be used with caution in patients receiving hydralazine. When other potent parenteral antihypertensive drugs, such as diazoxide, are used in combination with hydralazine, patients should be continuously observed for several hours for any excessive fall in blood pressure. Profound hypotensive episodes may occur when diazoxide injection and hydralazine are used concomitantly.

### **Drug/Food Interactions**

Administration of hydralazine with food results in higher plasma levels.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

In a lifetime study in Swiss albino mice, there was a statistically significant increase in the incidence of lung tumors (adenomas and adenocarcinomas) of both male and female mice given hydralazine continuously in their drinking water at a dosage of about 250 mg/kg per day (about 80 times the maximum recommended human dose). In a 2-year carcinogenicity study of rats given hydralazine by gavage at dose levels of 15, 30, and 60 mg/kg/day (approximately 5 to 20 times the recommended human daily dosage), microscopic examination of the liver revealed a small, but statistically significant, increase in benign neoplastic nodules in male and female rats from the high-dose group and in female rats from the intermediate-dose group. Benign interstitial cell tumors of the testes were also significantly increased in male rats from the high-dose group. The tumors observed are common in aged rats and a significantly increased incidence was not observed until 18 months of treatment. Hydralazine was shown to be mutagenic in bacterial systems (Gene Mutation and DNA Repair) and in one of two rats and one rabbit hepatocyte *in vitro* DNA repair studies. Additional *in vivo* and *in vitro* studies using lymphoma cells, germinal cells, and fibroblasts from mice, bone marrow cells from

Chinese hamsters and fibroblasts from human cell lines did not demonstrate any mutagenic potential for hydralazine.

The extent to which these findings indicate a risk to man is uncertain. While long-term clinical observation has not suggested that human cancer is associated with hydralazine use, epidemiologic studies have so far been insufficient to arrive at any conclusions.

## **Pregnancy**

### *Teratogenic Effects*

### *Pregnancy Category C*

Animal studies indicate that hydralazine is teratogenic in mice at 20 to 30 times the maximum daily human dose of 200 to 300 mg and possibly in rabbits at 10 to 15 times the maximum daily human dose, but that it is nonteratogenic in rats. Teratogenic effects observed were cleft palate and malformations of facial and cranial bones.

There are no adequate and well-controlled studies in pregnant women. Although clinical experience does not include any positive evidence of adverse effects on the human fetus, hydralazine should be used during pregnancy only if the expected benefit justifies the potential risk to the fetus.

## **Nursing Mothers**

Hydralazine has been shown to be excreted in breast milk.

## **Pediatric Use**

Safety and effectiveness in pediatric patients have not been established in controlled clinical trials, although there is experience with the use of hydralazine in pediatric patients. The usual recommended oral starting dosage is 0.75 mg/kg of body weight daily in four divided doses. Dosage may be increased gradually over the next 3 to 4 weeks to a maximum of 7.5 mg/kg or 200 mg daily.

## **ADVERSE REACTIONS**

Adverse reactions with hydralazine are usually reversible when dosage is reduced. However, in some cases it may be necessary to discontinue the drug. The following adverse reactions have been observed, but there has not been enough systematic collection of data to support an estimate of their frequency.

### **Common**

headache, anorexia, nausea, vomiting, diarrhea, palpitations, tachycardia, angina pectoris.

### **Less Frequent**

#### ***Digestive***

constipation, paralytic ileus.

#### ***Cardiovascular***

hypotension, paradoxical pressor response, edema.

#### ***Respiratory***

dyspnea.

#### ***Neurologic***

peripheral neuritis, evidenced by paresthesia, numbness, and tingling, dizziness; tremors; muscle cramps; psychotic reactions characterized by depression, disorientation, or anxiety.

**Genitourinary**

difficulty in urination.

**Hematologic**

blood dyscrasias, consisting of reduction in hemoglobin and red cell count, leukopenia, agranulocytosis, purpura, lymphadenopathy; splenomegaly.

**Hypersensitive Reactions**

rash, urticaria, pruritus, fever, chills, arthralgia, eosinophilia, and rarely, hepatitis.

**Other**

nasal congestion, flushing, lacrimation, conjunctivitis.

**OVERDOSAGE**

**Acute Toxicity**

No deaths due to acute poisoning have been reported. Highest known dose survived: adults, 10 g orally.

Oral LD<sub>50</sub> in rats: 173 and 187 mg/kg.

**Signs and Symptoms**

Signs and symptoms of overdosage include hypotension, tachycardia, headache, and generalized skin flushing.

Complications can include myocardial ischemia and subsequent myocardial infarction, cardiac arrhythmia, and profound shock.

**Treatment**

There is no specific antidote.

The gastric contents should be evacuated, taking adequate precautions against aspiration and for protection of the airway. An activated charcoal slurry may be instilled if conditions permit. These manipulations may have to be omitted or carried out after cardiovascular status has been stabilized, since they might precipitate cardiac arrhythmias or increase the depth of shock.

Support of the cardiovascular system is of primary importance. Shock should be treated with plasma expanders. If possible, vasopressors should not be given, but if a vasopressor is required, care should be taken not to precipitate or aggravate cardiac arrhythmia.

Tachycardia responds to beta blockers. Digitalization may be necessary, and renal function should be monitored and supported as required.

No experience has been reported with extracorporeal or peritoneal dialysis.

**DOSAGE AND ADMINISTRATION**

Initiate therapy in gradually increasing dosages; adjust according to individual response. Start with 10 mg four times daily for the first 2 to 4 days, increase to 25 mg four times daily for the balance of the first week. For the second and subsequent weeks, increase

dosage to 50 mg four times daily. For maintenance, adjust dosage to the lowest effective levels.

The incidence of toxic reactions, particularly the L.E. cell syndrome, is high in the group of patients receiving large doses of hydralazine hydrochloride tablets.

In a few resistant patients, up to 300 mg of hydralazine hydrochloride tablets daily may be required for a significant antihypertensive effect. In such cases, a lower dosage of hydralazine hydrochloride tablets combined with a thiazide and/or reserpine or a beta blocker may be considered. However, when combining therapy, individual titration is essential to ensure the lowest possible therapeutic dose of each drug.

## HOW SUPPLIED

Hydralazine Hydrochloride Tablets USP are available as:

<b>10 mg</b>	Orange, round tablets debossed with SG on one side and 182 on other side. Available in Bottles of 100 (NDC 77771-182-01) and 1000 (NDC 77771-182-10).
--------------	---

<b>25 mg</b>	Orange, round tablets debossed with SG on one side and 183 on other side. Available in Bottles of 100 (NDC 77771-183-01) and 1000 (NDC 77771-183-10).
--------------	---

<b>50 mg</b>	Orange, round tablets debossed with SG on one side and 184 on other side. Available in Bottles of 100 (NDC 77771-184-01) and 1000 (NDC 77771-184-10).
--------------	---

<b>100 mg</b>	Orange, round tablets debossed with SG on one side and 185 on other side. Available in Bottles of 100 (NDC 77771-185-01).
---------------	---

Dispense in a tight, light-resistant container as defined in the USP, with a child-resistant closure (as required).

Store at 20° to 25° C (68° to 77° F); excursions permitted to 15° to 30° C (59° to 86° F). [See USP Controlled Room Temperature].

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Manufactured by:  
ScieGen Pharmaceuticals, Inc.  
Hauppauge, NY 11788 USA

Distributed by:  
Radha Pharmaceuticals, Inc.  
Hauppauge, NY 11788 USA

Rev: 2/2021

## Repackaging Information

Please reference the **How Supplied** section listed above for a description of individual tablets. This drug product has been received by Aphenia Pharma - TN in a manufacturer or distributor packaged configuration and repackaged in full compliance with all applicable cGMP regulations. The package configurations available from Aphenia are listed below:

---

Count	25 mg	50 mg	100 mg
90	71610-561-60	71610-562-60	-
180	71610-561-80	71610-562-80	-
270	71610-561-92	71610-562-92	-
540	-	71610-562-98	-
2000	-	-	71610-549-02
4200	-	-	71610-549-05

---

Store between 20°-25°C (68°-77°F). See USP Controlled Room Temperature. Dispense in a tight light-resistant container as defined by USP. Keep this and all drugs out of the reach of children.

Repackaged by:



Cookeville, TN 38506

20210601JH

### PRINCIPAL DISPLAY PANEL - 25 mg



NDC 71610-561 - Hydralazine HCl, USP 25 mg Tablets - Rx Only

Hydralazine HCl, USP  
25mg 90 Tablets  
NDC# 71610-0561-60

Store between 20-25 degrees C(68-77 degrees F).  
See USP Controlled Room Temperature.  
Dispense in a tight light-resistant container  
as defined by USP.  
Keep this and all drugs out of the reach of children.

MFG BY: SolisGen Pharmaceuticals Inc  
Hauppauge, NY  
RPKG BY: Aphenix Pharma, Cookeville, TN 38506  
RX Only

MFG LOT: GTIN: 10371610561809 S/N: P018757748  
BATCH: 1 EXP:

### PRINCIPAL DISPLAY PANEL - 50 mg

NDC 71610-562 - Hydralazine HCl, USP 50 mg Tablets - Rx Only

Hydralazine HCl, USP  
50mg 90 Tablets  
NDC# 71610-0562-60

Store between 20-25 degrees C(68-77 degrees F).  
See USP Controlled Room Temperature.  
Dispense in a tight light-resistant container  
as defined by USP.  
Keep this and all drugs out of the reach of children.

MFG BY: SolisGen Pharmaceuticals Inc  
Hauppauge, NY  
RPKG BY: Aphenix Pharma, Cookeville, TN 38506  
RX Only

MFG LOT: GTIN: 10371610562606 S/N: P018757750  
BATCH: 2 EXP:




### PRINCIPAL DISPLAY PANEL - 100 mg



NDC 71610-549 - Hydralazine HCl, USP 100 mg Tablets - Rx Only

Hydralazine HCl, USP  
100mg 2000 Tablets  
NDC# 71610-0549-02

Store between 20-25 degrees C(68-77 degrees F).  
See USP Controlled Room Temperature.  
Dispense in a tight light-resistant container  
as defined by USP.  
Keep this and all drugs out of the reach of children.

MFG BY: SolisGen Pharmaceuticals Inc  
Hauppauge, NY  
RPKG BY: Aphenix Pharma, Cookeville, TN 38506  
RX Only

MFG LOT: GTIN: 10371610549027 S/N: P018634273  
BATCH: 3 EXP:

## HYDRALAZINE HYDROCHLORIDE

hydralazine hydrochloride tablet

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:71610-561(NDC:77771-183)
<b>Route of Administration</b>	ORAL		



**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
HYDRALAZINE HYDROCHLORIDE (UNII: FD171B778Y) (HYDRALAZINE - UNII:26NAK24LS8)	HYDRALAZINE HYDROCHLORIDE	25 mg

**Inactive Ingredients**

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

**Product Characteristics**

Color	orange	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	SG;183
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71610-561-60	90 in 1 BOTTLE; Type 0: Not a Combination Product	05/25/2021	
2	NDC:71610-561-80	180 in 1 BOTTLE; Type 0: Not a Combination Product	05/25/2021	
3	NDC:71610-561-92	270 in 1 BOTTLE; Type 0: Not a Combination Product	05/25/2021	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA205236	03/04/2021	

**HYDRALAZINE HYDROCHLORIDE**

hydralazine hydrochloride tablet

**Product Information**

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:71610-562(NDC:77771-184)
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
HYDRALAZINE HYDROCHLORIDE (UNII: FD171B778Y) (HYDRALAZINE - UNII:26NAK24LS8)			HYDRALAZINE HYDROCHLORIDE	50 mg
Inactive Ingredients				
Ingredient Name			Strength	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)				
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
Product Characteristics				
Color	orange	Score	no score	
Shape	ROUND	Size	8mm	
Flavor		Imprint Code	SG;184	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71610-562-60	90 in 1 BOTTLE; Type 0: Not a Combination Product	05/25/2021	
2	NDC:71610-562-80	180 in 1 BOTTLE; Type 0: Not a Combination Product	05/25/2021	
3	NDC:71610-562-92	270 in 1 BOTTLE; Type 0: Not a Combination Product	05/25/2021	
4	NDC:71610-562-98	540 in 1 BOTTLE; Type 0: Not a Combination Product	05/25/2021	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA205236	03/04/2021		

## HYDRALAZINE HYDROCHLORIDE

hydralazine hydrochloride tablet

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:71610-549(NDC:77771-185)
Route of Administration	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
HYDRALAZINE HYDROCHLORIDE (UNII: FD171B778Y) (HYDRALAZINE - UNII:26NAK24LS8)	HYDRALAZINE HYDROCHLORIDE	100 mg

**Inactive Ingredients**

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

**Product Characteristics**

Color	orange	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	SG;185
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71610-549-02	2000 in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2024	
2	NDC:71610-549-05	4200 in 1 BOTTLE; Type 0: Not a Combination Product	04/30/2021	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA205236	03/04/2021	

**Labeler** - Aphenia Pharma Solutions - Tennessee, LLC (128385585)**Establishment**

Name	Address	ID/FEI	Business Operations
Aphenia Pharma Solutions - Tennessee, LLC		128385585	REPACK(71610-561, 71610-562, 71610-549)

Revised: 4/2024

Aphenia Pharma Solutions - Tennessee, LLC