

ASPIRIN- aspirin tablet, coated
Amerisource Bergen

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Good Neighbor Pharmacy 44-227

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - headache
 - menstrual pain
 - a cold
 - toothache
 - muscle pain
- or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- asthma (wheezing)
- hives
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks every day while using this product
- are age 60 or older
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 to 2 tablets every 4 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

black iron oxide, cellulose, colloidal anhydrous silica, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, propylene glycol, shellac wax, simethicone, sodium alginate, sodium bicarbonate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-426-9391

Principal display panel

**GOOD
NEIGHBOR
PHARMACY®**

Compare to Ecotrin® active ingredient†

NDC 46122-596-02

REGULAR STRENGTH

**Aspirin 325 mg
Pain Reliever (NSAID)**

Enteric Coated

125 Tablets | 325 mg Each

Actual Size

†This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Ecotrin®.

50844 REV0421A22757

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

Distributed By
AmerisourceBergen
1 West First Avenue
Conshohocken, PA 19428
Questions or Concerns?
www.mygnp.com

GOOD NEIGHBOR PHARMACY
BRAND PRODUCTS

SATISFACTION GUARANTEED

Drug Facts (continued)

Directions

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 to 2 tablets every 4 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients black iron oxide, cellulose, colloidal amorphous silica, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, polyethylene glycol, wax, stearic acid, sodium alginate, sodium bicarbonate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate

(Questions or comments?) 1-800-426-9391

†This product is not manufactured or distributed by Mediatech Products Inc., owner of the registered trademark Ecotrin®.

50944 REV0421A22757

Compare to Ecotrin® active ingredient†

NDC 46122-596-02

REGULAR STRENGTH

Aspirin 325 mg

Pain Reliever (NSAID)

Enteric Coated

125 Tablets | 325 mg Each

Actual Size 

SATISFACTION GUARANTEED

Distributed By
AmerisourceBergen
1 West First Avenue
Conshohocken, PA 19428
Questions or Concerns?
www.mygnp.com

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

ABC#: 10212526



0 87701 43194 4

No Print Area
Lot no. & Exp. Date

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts	Purpose
Active ingredient (in each tablet) Aspirin 325 mg (NSAID)*, Pain reliever, nonsteroidal anti-inflammatory drug	
Uses	temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> minor pain of arthritis headache menstrual pain a cold toothache muscle pain or as recommended by your doctor
Warnings	<p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Aspirin may cause a severe allergic reaction, which may include: <ul style="list-style-type: none"> hives throat swelling asthma (wheezing) hives shock </p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: <ul style="list-style-type: none"> take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have had stomach ulcers or bleeding problems have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug have 3 or more alcoholic drinks every day while using this product are age 60 or older take more or for a longer time than directed </p>
Drug Facts (continued)	<p>Do not use</p> <ul style="list-style-type: none"> if you are allergic to aspirin or any other pain reliever/fever reducer if you have ever had an allergic reaction to this product or any of its ingredients <p>Ask a doctor before use if</p> <ul style="list-style-type: none"> stomach bleeding warning applies to you you have a history of stomach problems, such as heartburn you are taking a diuretic you have high blood pressure, heart disease, liver cirrhosis or kidney disease you have asthma <p>Ask a doctor or pharmacist before use if you are taking a prescription drug for:</p> <ul style="list-style-type: none"> gout diabetes arthritis <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> an allergic reaction occurs. Seek medical help right away. you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> feeling faint vomiting blood have bloody or black stools have stomach pain that does not get better ringing in the ears or a loss of hearing occurs pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur <p>These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>

B-0045-227-57
 REV0421A22757

GOOD NEIGHBOR PHARMACY 44-227

ASPIRIN

aspirin tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:46122-596
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SHELLAC (UNII: 46N107B71O)	
SODIUM ALGINATE (UNII: C269C4G2ZQ)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
Polyvinyl Acetate Phthalate (UNII: 58QVG85GW3)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Product Characteristics

Color	orange	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;227
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46122-596-02	1 in 1 CARTON	06/01/2019	
1		125 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	06/01/2019	

Labeler - Amerisource Bergen (007914906)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(46122-596)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(46122-596) , pack(46122-596)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(46122-596)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(46122-596)

Revised: 9/2021

Amerisource Bergen