

**ASPIRIN- aspirin tablet 325mg tablet  
WinCo Foods LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**011R Winco Foods 67091 382 Aspirin tablet 325mg 100 ct and 300ct**

**DRUG FACTS**

***Active ingredient (in each tablet)***

Aspirin 325 mg (NSAID\*)

\*nonsteroidal anti-inflammatory drug

***Purposes***

Pain reliever/fever reducer

***Uses***

- for the temporary relief of minor aches and pains due to
  - o headache
  - o colds
  - o muscle pain
  - o menstrual pain
  - o toothache
  - o minor pain of arthritis
- temporarily reduces fever

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use** if you have ever had an allergic to Aspirin or any other pain reliever/fever reducer

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have not been drinking fluids
- you have lost a lot of fluid due to vomiting or diarrhea

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug
- under a doctor's care for any serious condition

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - o feel faint
  - o have bloody or black stools
  - o vomit blood
  - o have stomach pain that does not get better
    - pain gets worse or lasts more than 10 days
    - fever gets worse or lasts more than 3 days
    - redness or swelling is present in the painful area
    - any new symptoms appear
    - ringing in the ears or a loss of hearing occurs

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 to 2 tablets every 4 hours while symptoms last. Do not take more than 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information**

- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

**Inactive ingredients** corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide

Questions? Call 1-800-824-1706 Monday - Friday 9am - 4pm MST

**Drug Facts (continued)**

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**■ you have not been drinking fluids**

- you have lost a lot of fluid due to vomiting or diarrhea

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug
- under a doctor's care for any serious condition

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - have bloody or black stools
  - vomit blood
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present in the painful area
  - any new symptoms appear
- ringing in the ears or a loss of hearing occurs

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use aspirin during the last 3 months of pregnancy, unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 to 2 tablets every 4 hours while symptoms last. Do not take more than 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other Information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

**Inactive ingredients** Corn Starch, Croscarmellose Sodium, Hydroxyethylcellulose, Microcrystalline Cellulose, Mineral Oil, Titanium Dioxide

**Questions?** Call 1-800-824-1706 Monday - Friday 9am - 4pm MST

\*\*Compare to the active ingredient in GENUINE BAYER®



**Aspirin**

PAIN RELIEVER - FEVER REDUCER (NSAID)

**300**  
Tablets

325 mg EACH

**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

**Drug Facts**

**Active ingredient (in each tablet)** **Purposes**

Aspirin 325 mg (NSAID)\* ..... Pain reliever/fever reducer  
nonsteroidal anti-inflammatory drug

**Uses**

- for the temporary relief of minor aches and pains due to
  - headache
  - colds
  - muscle pain
  - minor pain of arthritis
- temporarily reduces fever

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer

**Drug Facts (continued under label)**

**DISTRIBUTED BY:** WINCO FOODS, LLC. BOISE, ID 83704  
 \*This product is not manufactured or distributed by BAYER HEALTHCARE LLC, distributor of GENUINE BAYER®.



0170 0222 WFF202

**PEEL HERE** LOT NO: EXP. DATE:

**UNVARNISHED AREA**

**Adhesive Area**

**Drug Facts (continued)**

**Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have not been drinking fluids
- you have lost a lot of fluid due to vomiting or diarrhea

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug
- under a doctor's care for any serious condition

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding:

- feel faint
- have bloody or black stools
- vomit blood

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- ringing in the ears or a loss of hearing occurs

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ drink a full glass of water with each dose

- adults and children 12 years and over: take 1 to 2 tablets every 4 hours while symptoms last. Do not take more than 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information** ■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)

- use by expiration date on package

**Inactive ingredients** corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide

**Questions?** Call 1-800-824-1706 Monday - Friday 9am - 4pm MST

\*\*Compare to the active ingredient in GENUINE BAYER®

**WinCo FOODS**

# Aspirin

PAIN RELIEVER - FEVER REDUCER (NSAID)

100

Tablets  
325 mg EACH

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

**Drug Facts**

**Active ingredient (in each tablet)**  
Aspirin 325 mg (NSAID)\*, . . . . . Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses**

- for the temporary relief of minor aches and pains due to: ■ headache ■ colds ■ muscle pain
- menstrual pain ■ toothache ■ minor pain of arthritis
- temporarily reduces fever

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives ■ facial swelling
- shock ■ asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**DISTRIBUTED BY:** WINCO FOODS, LLC, BOISE, ID 83704

\*\*This product is not manufactured or distributed by BAYER-HEALTHCARE LLC, distributor of GENUINE BAYER®.

0170552123008

017R 0821 WF2102

Varnish Omit Area

LOT #:      EXP. DATE:  
PEEL HERE FOR MORE DRUG FACTS

<b>ASPIRIN</b>			
aspirin tablet 325mg tablet			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:67091-382
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			
	<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
	ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
<b>Inactive Ingredients</b>			
	<b>Ingredient Name</b>		<b>Strength</b>

<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>MINERAL OIL</b> (UNII: T5L8T28FGP)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	TCL011
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67091-382-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	01/18/2022	
2	NDC:67091-382-30	300 in 1 BOTTLE; Type 0: Not a Combination Product	01/18/2022	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	01/18/2022	

**Labeler** - WinCo Foods LLC (056098817)

**Registrant** - TIME CAP LABORATORIES, INC. (037052099)

### Establishment

Name	Address	ID/FEI	Business Operations
TIME CAP LABORATORIES, INC.		037052099	manufacture(67091-382)

Revised: 5/2022

WinCo Foods LLC