

REGENCY POST MILKING TEAT DIP- post-milking phenol teat dip solution
ABS Corporation

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click [here](#).

Regency Post-Milking Teat Dip

ACTIVE INGREDIENTS:

ACTIVE INGREDIENTS	Purpose
Phenol (1.56% w/v)	Disinfectant
Phenolate Sodium (0.06% w/v)	Disinfectant

PURPOSE:

Active Ingredients	Purpose
Phenol (1.56% w/v)	Disinfectant
Phenolate Sodium (0.06% w/v)	Disinfectant

Keep Out of Reach of Children

☐ If swallowed, get medical help or contact Poison Control Center right away.

USES:

FOR ANIMAL USE ONLY

Aids in controlling mastitis organisms with a softening effect on teats.

Directions for Use:

Before milking, wash udder and teats with damp cloth. Immediately after milking, dip each teat 1 to 2 inches into REGENCY Teat Dip.

Allow each teat to air dry. Do not wipe.

During the winter months, keep cows inside until the teats have completely dried to prevent freezing and chapping.

When cow is being dried out, dip each teat in REGENCY 1X Post-Milking Teat Dip once a day for 3 to 4 days after the last milking.

CAUTION:

Avoid eye contact. Do not inhale fumes. For external use only.

FIRST AID:

For Ingestion - Immediately give castor oil or vegetable oil (15 -30cc). Get medical attention immediately.

Be ready to induce vomiting at the advice of a physician or poison control center.

For Eyes - Immediately flush eyes with water for at least 15 minutes. Wash affected skin with water.

When Using this product:

Do not Store in Direct Sunlight.

Avoid Excessive Heat.

USER SAFETY WARNING:

Do Not Allow Product To Freeze. If frozen, allow product to completely thaw in a warm room.

Shake and thoroughly mix product before using.

Rinse teat dip cup with water after each milking and invert to dry.

Never pour used teat dip into original container.

Keep cap on container when not in use.

INACTIVE INGREDIENTS:

Water, Glycerin, Linear Alkylbenzene Sulfonate, Sodium Borate, Fragrance, FD&C Blue # 1



Aids in controlling mastitis organisms with a softening effect on the teats.

1.56% (w/v) Phenol with 0.06% (w/v) Sodium Phenate
Teat Dip Will Not Burn, Dry, Irritate Or Chap Teats. Contains a special blend of emollients and phenol.

NOTE: Store at room temperature away from direct sunlight.

NET CONTENTS: 1 U.S. Gallon (3.785 L)

FOR ANIMAL USE ONLY
KEEP OUT OF REACH OF CHILDREN
DO NOT STORE IN DIRECT SUNLIGHT
AVOID EXCESSIVE HEAT



FIRST AID: For Ingestion - Immediately give castor oil or vegetable oil (15 - 30cc). Get medical attention immediately. Be ready to induce vomiting at the advice of a physician or poison control center. For Eyes - Immediately flush eyes with water for at least 15 minutes. Wash affected skin with water.

CAUTION: Avoid eye contact. For external use only.

DIRECTIONS: Before milking, wash udder and teats with damp cloth. Immediately after milking, dip each teat 1 to 2 inches into Regency 1X Post-Milking Teat Dip.

Allow each teat to air dry. Do not wipe. During the winter months, keep cows inside until the teats have completely dried to prevent freezing and chapping. If the solution in the dip cup becomes contaminated, discard and use fresh solution. Rinse teat dip cup with water after each milking and invert to dry. Never pour used teat dip into the original container. Keep cap on container when not in use.

When a cow is being dried out, dip each teat in Regency 1X Post-Milking Teat Dip once a day for 3 to 4 days after the last milking. Do not allow product to freeze. If frozen, allow the product to completely thaw in a warm room. Shake and thoroughly mix product before using.

INGREDIENTS: Water, Glycerin, Phenol, LAS (Linear Alkylbenzene Sulfonate), Sodium Borate, Fragrance, FD&C Blue #1.

Distributed Exclusively By:

CLSolutions, LLC
PO BOX 837
Fulton, TX 78358
(361) 729-3993

www.CoastalLivestock.com

Sporicidin* is a registered trademark of Contec, Inc.
United States Patent # US6,586,477 B1

CLS-1X-1GAL



RegencyPost-Milking Teat Dip Masticid® with Sporicidin™ Technology

"Ready To Use Udder Wash"

Aids in controlling mastitis organisms with a softening effect on the teats.

1.56% (w/v) Phenol with 0.06% (w/v) Sodium Phenate

Teat Dip will Not Burn, Dry, Irritate of Chap Teats. Contains a special blend of emollients and ophenol.

NET CONTENTS: 1 U.S. Gallon (3.785 L)

RegencyTM

Masticide[®]
"Ready To Use Teat Dip"

1X Post-Milking Teat Dip



Aids in controlling mastitis organisms with a softening effect on the teats.
1.56% (w/v) Phenol with 0.06% (w/v) Sodium Phenate
 Teat Dip Will Not Burn, Dry, Irritate Or Chap Teats. Contains a special blend of emollients and phenol.
 NOTE: Store at room temperature away from direct sunlight.

FOR ANIMAL USE ONLY
 KEEP OUT OF REACH OF CHILDREN
 DO NOT STORE IN DIRECT SUNLIGHT
 AVOID EXCESSIVE HEAT



NET CONTENTS: 5 U.S. Gallons (18.927 L)

FIRST AID: For Ingestion - Immediately give castor oil or vegetable oil (15 - 30cc). Get medical attention immediately. Be ready to induce vomiting at the advice of a physician or poison control center. For Eyes - Immediately flush eyes with water for at least 15 minutes. Wash affected skin with water.

CAUTION: Avoid eye contact. For external use only.
DIRECTIONS: Before milking, wash udder and teats with damp cloth. Immediately after milking, dip each teat 1 to 2 inches into Regency 1X Post-Milking Teat Dip.

Allow each teat to air dry. Do not wipe. During the winter months, keep cows inside until the teats have completely dried to prevent freezing and chapping. If the solution in the dip cup becomes contaminated, discard and use fresh solution. Rinse teat dip cup with water after each milking and invert to dry. Never pour used teat dip into the original container. Keep cap on container when not in use.

When a cow is being dried out, dip each teat in Regency 1X Post-Milking Teat Dip once a day for 3 to 4 days after the last milking. Do not allow product to freeze. If frozen, allow the product to completely thaw in a warm room. Shake and thoroughly mix product before using.

INGREDIENTS: Water, Glycerin, Phenol, LAS (Linear Alkylbenzene Sulfonate), Sodium Borate, Fragrance, FD&C Blue #1.

Distributed Exclusively By:
 CLSolutions, LLC
 PO BOX 837
 Fulton, TX 78358
 (361) 729-3993
 www.CoastalLivestock.com
 Sporicidin[®] is a registered trademark of Contec, Inc.
 United States Patent # US6,586,477 B1
 CLS-1X-5GAL



Regency Post-Milking Teat Dip Masticid[®] with SporicidinTM Technology "Ready To Use Udder Wash"

Aids in controlling mastitis organisms with a softening effect on the teats.

1.56% (w/v) Phenol with 0.06% (w/v) Sodium Phenate

Teat Dip will Not Burn, Dry, Irritate of Chap Teats. Contains a special blend of emollients and ophenol.

NET CONTENTS: 5 U.S. Gallon (18.927 L)

RegencyTM

Masticide[®]
"Ready To Use Teat Dip"

1X Post-Milking Teat Dip



Aids in controlling mastitis organisms with a softening effect on the teats.
1.56% (w/v) Phenol with 0.06% (w/v) Sodium Phenate
 Teat Dip Will Not Burn, Dry, Irritate Or Chap Teats. Contains a special blend of emollients and phenol.
 NOTE: Store at room temperature away from direct sunlight.

FOR ANIMAL USE ONLY
 KEEP OUT OF REACH OF CHILDREN
 DO NOT STORE IN DIRECT SUNLIGHT
 AVOID EXCESSIVE HEAT



NET CONTENTS: 55 U.S. Gallons (208.2 L)

FIRST AID: For Ingestion - Immediately give castor oil or vegetable oil (15 - 30cc). Get medical attention immediately. Be ready to induce vomiting at the advice of a physician or poison control center. For Eyes - Immediately flush eyes with water for at least 15 minutes. Wash affected skin with water.

CAUTION: Avoid eye contact. For external use only.
DIRECTIONS: Before milking, wash udder and teats with damp cloth. Immediately after milking, dip each teat 1 to 2 inches into Regency 1X Post-Milking Teat Dip.

Allow each teat to air dry. Do not wipe. During the winter months, keep cows inside until the teats have completely dried to prevent freezing and chapping. If the solution in the dip cup becomes contaminated, discard and use fresh solution. Rinse teat dip cup with water after each milking and invert to dry. Never pour used teat dip into the original container. Keep cap on container when not in use.

When a cow is being dried out, dip each teat in Regency 1X Post-Milking Teat Dip once a day for 3 to 4 days after the last milking. Do not allow product to freeze. If frozen, allow the product to completely thaw in a warm room. Shake and thoroughly mix product before using.

INGREDIENTS: Water, Glycerin, Phenol, LAS (Linear Alkylbenzene Sulfonate), Sodium Borate, Fragrance, FD&C Blue #1.

Distributed Exclusively By:
 CLSolutions, LLC
 PO BOX 837
 Fulton, TX 78358
 (361) 729-3993
 www.CoastalLivestock.com
 Sporicidin[®] is a registered trademark of Contec, Inc.
 United States Patent # US6,586,477 B1
 CLS-1X-55GAL



Regency Post-Milking Teat Dip Masticid[®] with SporicidinTM Technology "Ready To Use Udder Wash"

Aids in controlling mastitis organisms with a softening effect on the teats.

1.56% (w/v) Phenol with 0.06% (w/v) Sodium Phenate

Teat Dip will Not Burn, Dry, Irritate of Chap Teats. Contains a special blend of emollients and ophenol.
NET CONTENTS: 55 U.S. Gallon (208.2 L)



1X Post-Milking Teat Dip



Aids in controlling mastitis organisms with a softening effect on the teats.

1.56% (w/v) Phenol with 0.06% (w/v) Sodium Phenate
 Teat Dip Will Not Burn, Dry, Irritate Or Chap Teats. Contains a special blend of emollients and phenol.

NOTE: Store at room temperature away from direct sunlight.

FOR ANIMAL USE ONLY
 KEEP OUT OF REACH OF CHILDREN
 DO NOT STORE IN DIRECT SUNLIGHT
 AVOID EXCESSIVE HEAT



NET CONTENTS: 270 U.S. Gallons

FIRST AID: For Ingestion - Immediately give castor oil or vegetable oil (15 - 30cc). Get medical attention immediately. Be ready to induce vomiting at the advice of a physician or poison control center. For Eyes - Immediately flush eyes with water for at least 15 minutes. Wash affected skin with water.

CAUTION: Avoid eye contact. For external use only.

DIRECTIONS: Before milking, wash udder and teats with damp cloth. Immediately after milking, dip teat entirely into Regency® Masticide® 1X Post-Milking Teat Dip.

Allow each teat to air dry. Do not wipe. During the winter months, keep cows inside until the teats have completely dried to prevent freezing and chapping. If the solution in the dip cup becomes contaminated, discard and use fresh solution. Rinse teat dip cup with water after each milking and invert to dry. Never pour used teat dip into the original container. Keep cap on container when not in use.

When a cow is being dried out, dip each teat in Regency® Masticide® 1X Post-Milking Teat Dip once a day for 3 to 4 days after the last milking.

Do not allow product to freeze. If frozen, allow the product to completely thaw in a warm room. Shake and thoroughly mix product before using.

INGREDIENTS: Water, Glycerin, Phenol, LAS (Linear Alkylbenzene Sulfonate), Sodium Borate, Fragrance, FD&C Blue #1.

Distributed Exclusively By:
 CLSolutions, LLC™
 7031 North 16th St.
 Omaha, NE 68112
 (402) 502-3928
 www.elsag.com
 CLS-1X-REV-8

Sporicidin® is a registered trademark of Contec, Inc.
 United States Patent # U56,586,477 B1

LOT NO. _____

Regency Post-Milking Teat Dip Masticid® with Sporicidin™ Technology

"Ready To Use Udder Wash"

Aids in controlling mastitis organisms with a softening effect on the teats.

1.56 % (w/v) Phenol with 0.06 % (w/v) Sodium Phenate

Teat Dip will Not Burn, Dry, Irritate of Chap Teats. Contains a special blend of emollients and ophenol.

NET CONTENTS: 270 U.S. Gallon

REGENCY POST MILKING TEAT DIP

post-milking phenol teat dip solution

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:18472-010
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENOL (UNII: 339NCG44TV) (PHENOL - UNII:339NCG44TV)	PHENOL	156 g in 1 L
PHENOLATE SODIUM (UNII: 4NC0T56V35) (PHENOL - UNII:339NCG44TV)	PHENOL	0.06 g in 1 L

Inactive Ingredients

Ingredient Name	Strength
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
SODIUM BORATE (UNII: 91MBZ8H3QO)	

GLYCERIN (UNII: PDC6A3C0OX)	
P-DODECYLBENZENESULFONIC ACID (UNII: OC21S23N1O)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:18472-010-02	1021.95 L in 1 CONTAINER		
2	NDC:18472-010-03	3.785 L in 1 JUG		
3	NDC:18472-010-04	18.925 L in 1 PAIL		
4	NDC:18472-010-05	208.175 L in 1 DRUM		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		06/13/2011	

Labeler - ABS Corporation (048022727)

Registrant - ABS Corporation (048022727)

Establishment

Name	Address	ID/FEI	Business Operations
ABS Corporation		048022727	api manufacture, pack, label