

**TOPCARE ACID REDUCER- famotidine tablet, film coated**  
**Topco Associates LLC**

-----  
**Topco Associates LLC. Acid Reducer Drug Facts**

**Active ingredient (in each tablet)**

Famotidine 20 mg

**Purpose**

Acid reducer

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings**

**Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes** before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

**Other information**

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture

**Inactive ingredients**

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, titanium dioxide

**Questions or comments?**

**1-888-423-0139**

**Principal Display Panel**

TopCare® health

COMPARE TO MAXIMUM STRENGTH PEPCID® AC ACTIVE INGREDIENT

MAXIMUM STRENGTH

Acid Reducer

FAMOTIDINE TABLETS, 20 mg

ACID REDUCER

Just One Tablet

Prevents & Relieves Heartburn


Due to Acid Indigestion

50 TABLETS

actual size

19471 88 C0

CODE AREA



**Drug Facts (continued)**

**Directions**  
 ■ Adults and children 12 years and over: Take 1 tablet with a glass of water. Do not chew.  
 ■ If you have symptoms, swallow 1 tablet with a glass of water. Do not chew.  
 ■ If you are taking other acid-reducing drugs, ask your doctor.  
 ■ Do not use more than 2 tablets in 24 hours.  
 ■ Children under 12 years: Ask a doctor.

**Other information**  
 ■ Read the directions and warnings on the carton before use.  
 ■ Keep the carton, it contains important information.

**Drug Facts (continued)**

■ Store at 20°-25°C (68°-77°F); excursions permitted to 15°-30°C (59°-86°F) as indicated on the label.  
 ■ Protect from moisture.  
**Inactive ingredients** can include water, colloidal silicon dioxide, croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, titanium dioxide.

**Questions or comments?** 1-888-423-0139

**Drug Facts**

**Active ingredient**  
 Famotidine 20 mg

**Purpose**  
 Acid reducer

**Uses**  
 ■ Relieves heartburn associated with acid indigestion and sour stomach.  
 ■ Prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages.

**Warnings**  
 ■ Allergy alert: Do not use if you are allergic to famotidine or other acid reducers.  
 ■ Do not use if you have certain swallowing problems, vomit, or are taking other acid-reducing drugs.

**Directions**  
 ■ Ask a doctor before use if you have heartburn more often than once a week, or if you have trouble swallowing, or if you have lost weight without trying to lose it.  
 ■ If pregnant or breastfeeding, ask a health professional before use.  
 ■ Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center for help. (1-800-222-1222)

**Drug Facts (continued)**

**Warnings**  
 ■ Allergy alert: Do not use if you are allergic to famotidine or other acid reducers.  
 ■ Do not use if you have certain swallowing problems, vomit, or are taking other acid-reducing drugs.

**Directions**  
 ■ Ask a doctor or pharmacist before use if you are taking a prescription drug for acid reflux or other conditions.  
 ■ Stop use and ask a doctor if:  
 ■ You are not getting better after 14 days of use.  
 ■ You are having trouble swallowing or vomiting.  
 ■ You have unexplained weight loss.  
 ■ You have frequent heartburn, especially with heartburn or indigestion.  
 ■ You have chest pain or a sharp pain spreading to arms, neck, or shoulders.  
 ■ You have dizziness or lightheadedness when sitting or standing.

DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

**Drug Facts (continued)**

**Warnings**  
 ■ Allergy alert: Do not use if you are allergic to famotidine or other acid reducers.  
 ■ Do not use if you have certain swallowing problems, vomit, or are taking other acid-reducing drugs.

**Directions**  
 ■ Ask a doctor or pharmacist before use if you are taking a prescription drug for acid reflux or other conditions.  
 ■ Stop use and ask a doctor if:  
 ■ You are not getting better after 14 days of use.  
 ■ You are having trouble swallowing or vomiting.  
 ■ You have unexplained weight loss.  
 ■ You have frequent heartburn, especially with heartburn or indigestion.  
 ■ You have chest pain or a sharp pain spreading to arms, neck, or shoulders.  
 ■ You have dizziness or lightheadedness when sitting or standing.

**Scan here for more information or call 1-888-423-0139**

**TopCare®** product is laboratory tested to guarantee its highest quality. Your local pharmacist is guaranteed.

\* This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Maltum Steingly Peptic® AC.

**GF GLUTEN FREE**

**TopCare®** health  
**MAXIMUM STRENGTH**  
**Acid Reducer**  
 FAMOTIDINE TABLETS, 20 mg  
 ACID REDUCER

**QUESTIONS?** 1-888-423-0139  
 ©TOPCO PERFORMANCE  
 ELK GROVE VILLAGE, IL 60007  
 TOPCO ASSOCIATES, LLC  
 DISTRIBUTED BY



**TopCare®** health

**MAXIMUM STRENGTH**


**Acid Reducer**

FAMOTIDINE TABLETS, 20 mg  
 ACID REDUCER

**Just One Tablet**  
 Prevents & Relieves Heartburn

NDC 36800-194-71

COMPARE TO MAXIMUM STRENGTH PEPCID® AC ACTIVE INGREDIENT\*



Due to Acid Indigestion

50 TABLETS

actual  
size**TOPCARE ACID REDUCER**

famotidine tablet, film coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:36800-194
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>FAMOTIDINE</b> (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ05DW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

**Product Characteristics**

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	8mm
<b>Flavor</b>		<b>Imprint Code</b>	L194
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-194-71	1 in 1 CARTON	10/16/2006	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:36800-194-	50 in 1 CARTON	10/16/2006	12/31/2020

2	51	1 in 1 CARTON	10/10/2006	12/31/2020
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:36800-194-02	25 in 1 CARTON	09/28/2006	
3		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
4	NDC:36800-194-78	1 in 1 CARTON	02/02/2021	
4		100 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077351	09/28/2006	

**Labeler** - Topco Associates LLC (006935977)

Revised: 11/2021

Topco Associates LLC