MOTION SICKNESS RELIEF- dimenhydrinate tablet Amerisource Bergen

Good Neighbor Pharmacy 44-198

Active ingredient (in each tablet)

Dimenhydrinate 50 mg

Purpose

Antiemetic

Uses

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

Warnings

Do not use

for children under 2 years of age unless directed by a doctor.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- marked drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

 to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity

adults and children 12 years and over	1 to 2 tablets every 4-6 hours; do not exceed 8 tablets in 24 hours, or as directed by a doctor
children 6 to under 12 years	$\frac{1}{2}$ to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours, or as directed by a doctor
Childran / Fo	$\frac{1}{2}$ tablet every 6-8 hours; do not exceed $1\frac{1}{2}$ tablets in 24 hours, or as directed by a doctor

Other information

- each tablet contains: calcium 35 mg
- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from moisture
- see end flap for expiration date and lot number

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid

Questions or comments?

1-800-426-9391

Principal display panel

GOOD NEIGHBOR PHARMACY®

NDC 46122-536-53

Compare to Dramamine® Original Formula active ingredient*

Motion Sickness Relief

dimenhydrinate 50 mg / antiemetic

Original Formula

Prevents Nausea, Vomiting & Dizziness Children & Adults

12 Tablets | 50 mg Each

Actual Size

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

50844 REV0518B19802

Distributed By AmerisourceBergen 1 West First Avenue Conshohocken, PA 19428 Questions or Concerns? www.mygnp.com



Product Information Product Type HUMAN OTC DRUG Item Code (Source) NDC:46122-536 Route of Administration ORAL Active Ingredient/Active Moiety Ingredient Name Basis of Strength

Strength

MOTION SICKNESS RELIEF

Inactive Ingredients	
Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETI7Z6XBU4)	

Product Characteristics			
Color	white	Score	2 pieces
Shape	ROUND	Size	9mm
Flavor		Imprint Code	44;198
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:46122-536- 53	2 in 1 CARTON	12/01/1992		
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M009	12/01/1992	

Labeler - Amerisource Bergen (007914906)

STEARIC ACID (UNII: 4ELV7Z65AP)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(46122-536)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(46122-536) , pack(46122-536)

Establishment			
Name	Address	ID/FEI	Business Operations

LNK International, Inc. 117025878 manufacture(46122-536)

Revised: 3/2024 Amerisource Bergen