

KANDESN CC CREAM MEDIUM- sunscreen broad spectrum spf30 cream Sunrider Manufacturing L.P.

Kandesn CC Cream Medium

USES:

- helps prevent sunburn
- if used as directed with other sun protection measures (see Directions), decreased the risk of skin cancer and early skin aging caused by the sun.

DIRECTIONS:

- apply liberally 15 minutes before sun exposure
- Reapply:
 - after 40 minutes of swimming or sweating
 - immediately after towel drying
 - at least every 2 hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. - 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses
- Children under 6 months: Ask a doctor

ACTIVE INGREDIENTS:

Titanium Dioxide 11.4%

Zinc Oxide 5.5%

OTHER INGREDIENTS:

Water (Deionized)
Caprylic/Capric
Triglyceride, Octyldodecyl
Neopentanoate, VP/Eicosene Copolymer,
Squalane (Veg. Origin),
Cyclopentasiloxane, Glyceryl Stearate,
PEG-100 Stearate, Cetyl Alcohol,
Hydrogen Dimethicone, Alumina,
Glycerin, Iron Oxides, Olea Europaea
(Olive) Fruit Oil, Persea Gratissima
(Avocado) Oil, Simmondsia Chinensis
(Jojoba) Seed Oil, Tocopheryl Acetate
(Natural), Tocopherol, Lysine PCA,
Ceramide-3, Panicum Miliaceum (Millet)
Seed Extract, Ceratonia Siliqua (Carob)
Gum, Phospholipids, Ruscus Aculeatus
Root Extract (Butcherbroom), Avena
Sativa (Oat) Kernel Protein, Symphytum
Officinale Extract (Comfrey), Plantago
Ovata Seed Extract (Plantain), Hydrolyzed

Wheat Protein, Xanthan Gum,
Cyclomethicone, Triethoxycaprylylsilane,
Sodium Polyacrylate, Propylene Glycol,
Cellulose Gum, Dimethiconol, Carbomer,
Triethanolamine, Magnesium Aluminum
Silicate, Polyhydroxystearic Acid, Phenyl
Trimethicone, Hexyldecanol, Citric Acid,
Trisodium EDTA, Phenoxyethanol,
Ethylhexylglycerin, Sodium Benzoate,
and Fragrance.

WARNINGS:

FOR EXTERNAL USE ONLY

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.


Stop use and ask a doctor if rash occurs.

Keep out of reach of children. If product is swallowed, get medical help or contact a
Poison Control Center right away.

KEEP OUT OF REACH OF CHILDREN. If product is swallowed, get medical help or contact
a Poison Control Center right away.

CC Cream Medium

Carton



BOX EXTERIOR PRINTS
 SPOT BLUE PMS 3035
 SPOT UV (SEE LAYER) PROCESS

BOX INTERIOR PRINTS PROCESS

Label

Kandesn®

CC
CREAM
SUNSCREEN
BROAD SPECTRUM
SPF 30

medium

1.25 fl. oz./37 mL

Uses

- helps prevent sunburn
- If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions

- apply liberally 15 minutes before sun exposure

• reapply:

- after 40 minutes of swimming or sweating
- Immediately after towel drying
- at least every 2 hours

• **Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. – 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor

Other Information

- protect this product from excessive heat and direct sun

Manufactured by
Sunrider Manufacturing, L.P.
Midlothian, Texas 76065, USA
Product Inquiry: 1-310-781-8096



SUNRIDER®

US S870770134

KANDESN CC CREAM MEDIUM

sunscreen broad spectrum spf30 cream

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:62191-013
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	114 mg in 1 g
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	55 mg in 1 g
Inactive Ingredients		
Ingredient Name	Strength	
WATER (UNII: 059QF0KO0R)		
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)		
OCTYLDODECYL NEOPENTANOATE (UNII: X8725R883T)		
VINYLPYRROLIDONE/EICOSENE COPOLYMER (UNII: 035MV9S1C3)		
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)		
SQUALANE (UNII: GW89575KF9)		
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)		
PEG-100 STEARATE (UNII: YD01N1999R)		
CETYL ALCOHOL (UNII: 936JST6JCN)		
METHICONE (20 CST) (UNII: 6777U11MKT)		
ALUMINUM OXIDE (UNII: LMI26O6933)		
OLIVE OIL (UNII: 6UYK2W1W1E)		
AVOCADO OIL (UNII: 6VNO72PFC1)		
JOJOBA OIL (UNII: 724GKU717M)		
ALPHA-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0)		
PHENOXYETHANOL (UNII: HIE492ZZ3T)		
MAGNESIUM ALUMINUM SILICATE (UNII: 6M3P64V0NC)		
POLYHYDROXYSTEARIC ACID (2300 MW) (UNII: YXH47AOU0F)		
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)		
HEXYLDECANOL (UNII: 151Z7P1317)		
CYCLOMETHICONE (UNII: NMQ347994Z)		
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)		
EDETATE TRISODIUM (UNII: 420IP921MB)		
XANTHAN GUM (UNII: TTV12P4NEE)		
GLYCERIN (UNII: PDC6A3C0OX)		
RUSCUS ACULEATUS ROOT (UNII: ZW12V95I1Q)		
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)		
ETHYLHEXYLGLYCERIN (UNII: 147D247K3P)		
.DELTA.-TOCOPHEROL (UNII: JU84X1I10N)		
LYSINE PIDOLATE (UNII: 89WJG4A348)		
Ceramide NP (UNII: 4370DF050B)		
MILLET (UNII: TJR6B3R47P)		
SODIUM BENZOATE (UNII: OJ245FE5EU)		
DIMETHICONOL (40 CST) (UNII: 343C7U75XW)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
EGG PHOSPHOLIPIDS (UNII: 1Z74184RGV)		
CARBOMER HOMOPOLYMER, UNSPECIFIED TYPE (UNII: 0A5MM307FC)		
Oatmeal (UNII: 8PI54V663Y)		
SYMPHYTUM X UPLANDICUM LEAF (UNII: D05H XK6R3G)		
PLANTAGO OVATA SEED (UNII: UD50RBY30F)		
HYDROLYZED WHEAT PROTEIN (ENZYMATIC, 3000 MW) (UNII: J2S07SB0YL)		
LOCUST BEAN GUM (UNII: V4716MY704)		
FERRIC OXIDE RED (UNII: 1K09F3G675)		

CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)

SODIUM POLYACRYLATE (2500000 MW) (UNII: 05I15JN12J)

FERRIC OXIDE YELLOW (UNII: EX438O2MRT)

FERROSOFERRIC OXIDE (UNII: XM0M87F357)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62191-013-01	1 in 1 PACKAGE	12/20/2021	
1		41.75 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	part352	09/01/2016	

Labeler - Sunrider Manufacturing L.P. (786951475)

Revised: 12/2023

Sunrider Manufacturing L.P.