CARE ONE CLEARLAX- polyethylene glycol 3350 powder, for solution
American Sales Company

American Sales Company ClearLax® Drug Facts

Active ingredient (in each dose)

Polyethylene Glycol 3350, 17 g (cap filled to line)

Purpose

Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use

if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are

taking a prescription drug

When using this product

you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not take more than directed unless advised by your doctor
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
- fill to top of white section in cap which is marked to indicate the correct dose (17 g)
- stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- use once a day
- use no more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20˚-25˚C (68˚-77˚F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

Inactive ingredients

none

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to the Active Ingredient in MiraLAX®

CLEARLAX®

Polyethylene Glycol 3350

Powder for Solution

Osmotic Laxative

UNFLAVORED POWDER – GRIT FREE

Relieves Occasional Constipation / Irregularity

Softens Stool

14 ONCE-DAILY DOSES

NET WT 8.3 OZ (238 g)

Compare to the Active Ingredient in MiraLAX®*

NDC 41520-306-02

CareOne®

CLEARLAX®

Polyethylene Glycol 3350 Powder for Solution Osmotic Laxative

UNFLAVORED POWDER - GRIT FREE

- Relieves Occasional Constipation / Irregularity
- Softens Stool

14 ONCE-DAILY DOSES

NET WT 8.3 OZ (238 g)

: 306E1 D4 F4

Peel Here

Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool.

Dissolves in Any Beverage

Drug Facts

Active ingredient (in each dose) **Purpose**

Polyethylene Glycol 3350, 17 g (cap filled to line) ..... Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have
- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if
- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

3 41520 31948 8

: 30E1 D4 B3

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
  - fill to top of white section in cap which is marked to indicate the correct dose (17 g)
  - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
  - use once a day
  - use no more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

Inactive ingredients none

Questions or comments? 1-800-719-9260

*This product is not manufactured or distributed by MSD Consumer Care, Inc., distributor of MiraLAX®. CLEARLAX® is a registered trademark of L. Perrigo Company.

• SUGAR FREE GLUTEN FREE

DISTRIBUTED BY FOODHOLD U.S.A., LLC
LANDOVER, MD 20785 - 1-877-846-9949
©2014 Ahold Licensing, Sarl

Quality guaranteed or your money back.

CARE ONE CLEARLAX

polyethylene glycol 3350 powder, for solution

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:41520-306
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL 3350 - UNII:G2M7P15E5P)	POLYETHYLENE GLYCOL 3350	17 g in 17 g

Product Characteristics

Color	WHITE	**Score**	
Shape		**Size**	
Flavor		**Imprint Code**	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41520-306-02	238 g in 1 BOTTLE		
2	NDC:41520-306-03	510 g in 1 BOTTLE		
3	NDC:41520-306-01	119 g in 1 BOTTLE		
4	NDC:41520-306-18	153 g in 1 BOTTLE		

5	NDC:41520-306-17	289 g in 1 BOTTLE		
6	NDC:41520-306-19	595 g in 1 BOTTLE		
7	NDC:41520-306-04	850 g in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090685	10/09/2009	

Labeler - American Sales Company (809183973)

Revised: 8/2014 American Sales Company