

## METHOCARBAMOL - methocarbamol tablet Unit Dose Services


-----

### Methocarbamol Tablets USP Rx Only

#### DESCRIPTION

Methocarbamol tablets USP a carbamate derivative of guaifenesin, is a central nervous system (CNS) depressant with sedative and musculoskeletal relaxant properties.

The chemical name of methocarbamol is 3-(2-methoxyphenoxy)-1,2-propanediol 1-carbamate and has the empirical formula  $C_{11}H_{15}NO_5$ . Its molecular weight is 241.24. The structural formula is shown below.



Methocarbamol is a white powder, sparingly soluble in water and chloroform, soluble in alcohol (only with heating) and propylene glycol, and insoluble in benzene and n-hexane.

Methocarbamol tablets USP are available as 500 mg and 750 mg tablets for oral administration.

Methocarbamol tablets USP 500 mg and 750 mg contain the following inactive ingredients: sodium lauryl sulfate, sodium starch glycolate, povidone K 90, polyethylene glycol, magnesium stearate, colloidal silicon dioxide, low substituted hydroxypropyl cellulose and stearic acid.

#### CLINICAL PHARMACOLOGY

The mechanism of action of methocarbamol in humans has not been established, but may be due to general central nervous system (CNS) depression. It has no direct action on the contractile mechanism of striated muscle, the motor end plate or the nerve fiber.

#### Pharmacokinetics

In healthy volunteers, the plasma clearance of methocarbamol ranges between 0.20 and 0.80 L/h/kg, the mean plasma elimination half-life ranges between 1 and 2 hours, and the plasma protein binding ranges between 46% and 50%.

Methocarbamol is metabolized via dealkylation and hydroxylation. Conjugation of methocarbamol also is likely. Essentially all methocarbamol metabolites are eliminated in the urine. Small amounts of unchanged methocarbamol also are excreted in the urine.

#### Special populations

##### *Elderly*

The mean ( $\pm$ SD) elimination half-life of methocarbamol in elderly healthy volunteers (mean ( $\pm$ SD) age, 69 ( $\pm$ 4) years) was slightly prolonged compared to a younger (mean ( $\pm$ SD) age, 53.3 ( $\pm$ 8.8) years),

healthy population (1.5 ( $\pm$ 0.4) hours versus 1.1 ( $\pm$ 0.27) hours, respectively). The fraction of bound methocarbamol was slightly decreased in the elderly versus younger volunteers (41 to 43% versus 46 to 50%, respectively).

#### *Renally impaired*Renally impaired

The clearance of methocarbamol in 8 renally-impaired patients on maintenance hemodialysis was reduced about 40% compared to 17 normal subjects, although the mean ( $\pm$ SD) elimination half-life in these two groups was similar: 1.2 ( $\pm$  0.6) versus 1.1 ( $\pm$ 0.3) hours, respectively.

#### *Hepatically impaired*Hepatically impaired

In 8 patients with cirrhosis secondary to alcohol abuse, the mean total clearance of methocarbamol was reduced approximately 70% compared to that obtained in 8 age- and weight-matched normal subjects. The mean ( $\pm$ SD) elimination half-life in the cirrhotic patients and the normal subjects was 3.38 ( $\pm$ 1.62) hours and 1.11 ( $\pm$ 0.27) hours, respectively. The percent of methocarbamol bound to plasma proteins was decreased to approximately 40 to 45% compared to 46 to 50% in the normal subjects.

## **INDICATIONS & USAGE**

Methocarbamol tablets USP are indicated as an adjunct to rest, physical therapy, and other measures for the relief of discomfort associated with acute, painful musculoskeletal conditions. The mode of action of methocarbamol has not been clearly identified, but may be related to its sedative properties. Methocarbamol does not directly relax tense skeletal muscles in man.

## **CONTRAINDICATIONS**

Methocarbamol tablets USP are contraindicated in patients hypersensitive to methocarbamol or to any of the tablet components.

## **WARNINGS**

Since methocarbamol may possess a general CNS depressant effect, patients receiving methocarbamol tablets USP should be cautioned about combined effects with alcohol and other CNS depressants. Safe use of methocarbamol tablets USP has not been established with regard to possible adverse effects upon fetal development. There have been reports of fetal and congenital abnormalities following in utero exposure to methocarbamol. Therefore, methocarbamol tablets USP should not be used in women who are or may become pregnant and particularly during early pregnancy unless in the judgment of the physician the potential benefits outweigh the possible hazards (**see PRECAUTIONS, Pregnancy**).(see PRECAUTIONSPRECAUTIONS, PregnancyPregnancy).

### **Use In Activities Requiring Mental Alertness**

Methocarbamol may impair mental and/or physical abilities required for performance of hazardous tasks, such as operating machinery or driving a motor vehicle. Patients should be cautioned about operating machinery, including automobiles, until they are reasonably certain that methocarbamol therapy does not adversely affect their ability to engage in such activities.

## **PRECAUTIONS**

### **Information for Patients**

Patients should be cautioned that methocarbamol may cause drowsiness or dizziness, which may impair their ability to operate motor vehicles or machinery.

Because methocarbamol may possess a general CNS-depressant effect, patients should be cautioned about combined effects with alcohol and other CNS depressants.

### **Drug Interactions**

**See WARNINGS and PRECAUTIONS** See WARNINGS WARNINGS and PRECAUTIONS PRECAUTIONS for interaction with CNS drugs and alcohol.

Methocarbamol may inhibit the effect of pyridostigmine bromide. Therefore, methocarbamol should be used with caution in patients with myasthenia gravis receiving anticholinesterase agents.

### **Drug & OR Laboratory Test Interactions**

Methocarbamol may cause a color interference in certain screening tests for 5-hydroxyindoleacetic acid (5-HIAA) using nitrosonaphthol reagent and in screening tests for urinary vanillylmandelic acid (VMA) using the Gitlow method.

### **Carcinogenesis & Mutagenesis & Impairment Of Fertility**

Long-term studies to evaluate the carcinogenic potential of methocarbamol have not been performed. No studies have been conducted to assess the effect of methocarbamol on mutagenesis or its potential to impair fertility.

### **Pregnancy**

*Teratogenic effects -Pregnancy Category C* Teratogenic effects -Pregnancy Category C

Animal reproduction studies have not been conducted with methocarbamol. It is also not known whether methocarbamol can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Methocarbamol tablets USP should be given to a pregnant woman only if clearly needed. Safe use of methocarbamol tablets USP has not been established with regard to possible adverse effects upon fetal development. There have been reports of fetal and congenital abnormalities following in utero exposure to methocarbamol. Therefore, methocarbamol tablets USP should not be used in women who are or may become pregnant and particularly during early pregnancy unless in the judgment of the physician the potential benefits outweigh the possible hazards (see **WARNINGS** see WARNINGS WARNINGS).

### **Nursing Mothers**

Methocarbamol and/or its metabolites are excreted in the milk of dogs; however, it is not known whether methocarbamol or its metabolites are excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when methocarbamol tablets USP are administered to a nursing woman.

### **Pediatric Use**

Safety and effectiveness of methocarbamol tablets USP in pediatric patients below the age of 16 have not been established.

## **ADVERSE REACTIONS**

Adverse reactions reported coincident with the administration of methocarbamol include:

*Body as a whole*Body as a whole: Anaphylactic reaction, angioneurotic edema, fever, headache

*Cardiovascular system*Cardiovascular system: Bradycardia, flushing, hypotension, syncope, thrombophlebitis

*Digestive system*:Digestive system: Dyspepsia, jaundice (including cholestatic jaundice), nausea and vomiting

*Hemic and lymphatic system*:Hemic and lymphatic system: Leukopenia

*Immune system*Immune system: Hypersensitivity reactions

*Nervous system*:Nervous system: Amnesia, confusion, diplopia, dizziness or lightheadedness, drowsiness, insomnia, mild muscular incoordination, nystagmus, sedation, seizures (including grand mal), vertigo

*Skin and special senses*Skin and special senses: Blurred vision, conjunctivitis, nasal congestion, metallic taste, pruritus, rash, urticarial

## **OVERDOSAGE**

Limited information is available on the acute toxicity of methocarbamol. Overdose of methocarbamol is frequently in conjunction with alcohol or other CNS depressants and includes the following symptoms: nausea, drowsiness, blurred vision, hypotension, seizures, and coma.

In post-marketing experience, deaths have been reported with an overdose of methocarbamol alone or in the presence of other CNS depressants, alcohol or psychotropic drugs.

### **Treatment**

Management of overdose includes symptomatic and supportive treatment. Supportive measures include maintenance of an adequate airway, monitoring urinary output and vital signs, and administration of intravenous fluids if necessary. The usefulness of hemodialysis in managing overdose is unknown.

## **DOSAGE & ADMINISTRATION**

Methocarbamol Tablets USP 500 mg – Adults:

Initial dosage: 3 tablets q.i.d.

Maintenance dosage: 2 tablets q.i.d.

Methocarbamol Tablets USP 750 mg – Adults:

Initial dosage: 2 tablets q.i.d.

Maintenance dosage: 1 tablet q.4h. or 2 tablets t.i.d.

Six grams a day are recommended for the first 48 to 72 hours of treatment. (For severe conditions 8 grams a day may be administered). Thereafter, the dosage can usually be reduced to approximately 4 grams a day.

## **HOW SUPPLIED**

Product: 50436-0534

NDC: 50436-0534-1 30 TABLET in a BOTTLE

NDC: 50436-0534-2 60 TABLET in a BOTTLE

NDC: 50436-0534-3 90 TABLET in a BOTTLE

NDC: 50436-0534-4 45 TABLET in a BOTTLE

# METHOCARBAMOL TABLET

NDC: 50436-0533-1  
**METHOCARBAMOL**  
**500 MG / 30 TAB**

Each uncoated tablet contains  
 Methocarbamol USP 500mg  
 Dispense in tight container

WARNING: KEEP OUT OF  
 REACH OF CHILDREN.STORE AT  
 20-25 ° C(68-77 ° F) CONTROLLED  
 ROOM TEMPERATURE. SEE PACKAGE  
 INSERT FOR ADDITIONAL INFORMATION.



MFG BY: HETERO LABS LTD  
 HYDERABAD-500 055,INDIA  
 MFG NDC: 31722-533-01  
 MFG LOT: XXXXX  
 LOT: XXXXX EXP: XX/XX/XX  
 Pkg by: Unit Dose Services, LLC  
 Dania, FL 33004 **Rev.1 RX ONLY**

NDC: 50436-0533-1  
 DRUG: METHOCARBAMOL  
 500 MG / 30 TAB  
 LOT: XXXXX EXP: XX/XX/XX  
 NDC: 50436-0533-1  
 DRUG: METHOCARBAMOL  
 500 MG / 30 TAB  
 LOT: XXXXX EXP: XX/XX/XX  
 NDC: 50436-0533-1  
 DRUG: METHOCARBAMOL  
 500 MG / 30 TAB  
 LOT: XXXXX EXP: XX/XX/XX  
 NDC: 50436-0533-1  
 DRUG: METHOCARBAMOL  
 500 MG / 30 TAB  
 LOT: XXXXX EXP: XX/XX/XX

# METHOCARBAMOL TABLET

NDC: 50436-0534-1  
**METHOCARBAMOL**  
**750 MG / 30 TAB**

Each uncoated tablet contains  
 Methocarbamol USP 750mg  
 Dispense in tight container

WARNING: KEEP OUT OF  
 REACH OF CHILDREN.STORE AT  
 20-25 ° C(68-77 ° F) CONTROLLED  
 ROOM TEMPERATURE. SEE PACKAGE  
 INSERT FOR ADDITIONAL INFORMATION.



MFG BY: HETERO LABS LTD  
 HYDERABAD-500 055,INDIA  
 MFG NDC: 31722-534-01  
 MFG LOT: XXXXX  
 LOT: XXXXX EXP: XX/XX/XX  
 Pkg by: Unit Dose Services, LLC  
 Dania, FL 33004 **Rev.1 RX ONLY**

NDC: 50436-0534-1  
 DRUG: METHOCARBAMOL  
 750 MG / 30 TAB  
 LOT: XXXXX EXP: XX/XX/XX  
 NDC: 50436-0534-1  
 DRUG: METHOCARBAMOL  
 750 MG / 30 TAB  
 LOT: XXXXX EXP: XX/XX/XX  
 NDC: 50436-0534-1  
 DRUG: METHOCARBAMOL  
 750 MG / 30 TAB  
 LOT: XXXXX EXP: XX/XX/XX  
 NDC: 50436-0534-1  
 DRUG: METHOCARBAMOL  
 750 MG / 30 TAB  
 LOT: XXXXX EXP: XX/XX/XX

## METHOCARBAMOL

methocarbamol tablet

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:50436-0533(NDC:31722-533)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
METHOCARBAMOL (UNII: 125OD7737X) (METHOCARBAMOL - UNII:125OD7737X)	METHOCARBAMOL	500 mg

### Inactive Ingredients

Ingredient Name	Strength
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

<b>POVIDONE K90</b> (UNII: RDH86HJV5Z)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>HYDROXYPROPYL CELLULOSE (1200000 MW)</b> (UNII: RFW2ET671P)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	

### Product Characteristics

<b>Color</b>	WHITE (White to Offwhite)	<b>Score</b>	no score
<b>Shape</b>	OVAL (Capsule shaped)	<b>Size</b>	15mm
<b>Flavor</b>		<b>Imprint Code</b>	114;H
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50436-0533-1	30 in 1 BOTTLE; Type 0: Not a Combination Product	10/12/2017	
2	NDC:50436-0533-2	60 in 1 BOTTLE; Type 0: Not a Combination Product	10/12/2017	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090200	03/20/2013	

## METHOCARBAMOL

methocarbamol tablet

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:50436-0534(NDC:31722-534)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>METHOCARBAMOL</b> (UNII: 125OD7737X) (METHOCARBAMOL - UNII:125OD7737X)	METHOCARBAMOL	750 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>POVIDONE K90</b> (UNII: RDH86HJV5Z)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	

**HYDROXYPROPYL CELLULOSE (1200000 MW)** (UNII: RFW2ET671P)

**STEARIC ACID** (UNII: 4ELV7Z65AP)

### Product Characteristics

<b>Color</b>	WHITE (White to Offwhite)	<b>Score</b>	no score
<b>Shape</b>	OVAL (Capsule shaped)	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	115;H
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50436-0534-1	30 in 1 BOTTLE; Type 0: Not a Combination Product	10/12/2017	
2	NDC:50436-0534-2	60 in 1 BOTTLE; Type 0: Not a Combination Product	10/12/2017	
3	NDC:50436-0534-3	90 in 1 BOTTLE; Type 0: Not a Combination Product	10/12/2017	
4	NDC:50436-0534-4	45 in 1 BOTTLE; Type 0: Not a Combination Product	10/12/2017	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090200	03/20/2013	

**Labeler** - Unit Dose Services (831995316)

### Establishment

Name	Address	ID/FEI	Business Operations
Unit Dose Services		831995316	REPACK(50436-0534, 50436-0533) , RELABEL(50436-0533, 50436-0534)

Revised: 10/2017

Unit Dose Services