# ESOMEPRAZOLE MAGNESIUM- esomeprazole capsule, delayed release Rite Aid Corporation

-----

# **Rite Aid Corporation Esomeprazole Magnesium Drug Facts**

#### Active ingredient (in each capsule)

Esomeprazole 20 mg

(Each delayed-release capsule corresponds to 22 mg esomeprazole magnesium dihydrate)

#### Purpose

Acid reducer

# Uses

- treats frequent heartburn (occurs **<u>2 or more</u>** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

# Warnings

Allergy alert: Do not use if you are allergic to esomeprazole

# Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness**, **sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain

These may be signs of a serious condition. See your doctor.

# Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

# Ask a doctor or pharmacist before use if you are

• taking a prescription drug. Acid reducers may interact with certain prescription drugs.

# Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

# If pregnant or breast-feeding,

ask a health professional before use.

# Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

# Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- may take 1 to 4 days for full effect

# 14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

# **Repeated 14-Day Courses (if needed)**

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

# Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F)

# **Inactive ingredients**

FD&C blue no. 1, FD&C blue no. 1 aluminum lake, FD&C red no. 3, ferric oxide, gelatin, glyceryl monostearate, hypromellose, magnesium stearate, meglumine, methacrylic acid and ethyl acrylate copolymer dispersion, polyethylene glycol, polysorbate 80, shellac, sodium lauryl sulfate, sugar spheres, talc, titanium dioxide, triethyl citrate

# Questions or comments?

1-800-719-9260

# Package/Label Principal Display Panel

SEE NEW WARNING INFORMATION Compare to the active ingredient of Nexium<sup>®</sup> 24 HR Clear Minis<sup>™</sup> TREATS FREQUENT HEARTBURN MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT ESOMEPRAZOLE MAGNESIUM Delayed Release Capsules 20 mg Acid Reducer MINI CAPSULES 24 HR ACTUAL SIZE 42 CAPSULES THREE 14-DAY COURSES OF TREATMENT

	Tips for Managing Heartburn <ul> <li>Avoid foods or drinks that are more likely to cause heartburn, such as rich, spior, fang and fried foods, chocolaro, criffeine, alochel and even some acidic fruits and vegeschles.</li> <li>Eat sloudy and do not eat big meals.</li> <li>Do not eat late at right or just before beddime.</li> <li>Do not eat late at right or just before beddime.</li> <li>Do not eat late at right or just before beddime.</li> <li>Do not is flat or bend over soon after eating.</li> <li>Raise the head of your bed.</li> <li>User loose-fixing dotting aroundyour stormach.</li> <li>If you are overweight, lose weight.</li> </ul> <li>Exat FREQUENT HEARTBURN</li>	
ESOMEPRAZO Delayed Release Acid F	Arys FOR FULL EFFECT	ACTUAL SIZE
7U4D7 83 C1	COURSES OF TREATMENT	BROKEN OR MISSING.
KEEP CARTON FOR COMPLETE WARK         Drug Facts         Active ingredient (in each capsule)       Purpose         Boomeprace 20 mg.       Add reduced         (Each delayed-release capsule corresponds to       22 mg esomeprace magnesium ditydrate)         Uses       treats frequent hearitum (occurs 2 or more days a week)         In of intended for immediate relief of hearitum; this drug may take 1 to 4 days for full effect         Warnings         Allergy alert. Do not use if you are allergic to esomeprazole         Do not use if you have:         In trubie or pain swaltowing food, wording with blood, or bloody or black stools         In heartburn with lightheadedness, swesting or dizziness         Cheartburn with lightheadedness, swesting or dizziness         In capital pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness         Incommit to able of a serious conting.		Drug Facts (continued)         Other information         = read the directions and warnings before use         = keep the carlon. It contains important information.         = store at 20-25°C (68-77°F)         Inactive ingredients         RO&C blue no. 1, FD&C blue no. 1         aluminum lake, FD&C red no. 3, femic odde, gelstin, glycery monochearste, hypormitose, magnesium stearste, meglumine, methacrytic add and ethyl acrylate copolymer dispersion, polyethylene glycol, polysortate 80, shellac, socium lauyi suffate, sugar spheres, tak, titanium dixide, itehyl ctrate         Questions or comments?         1-800-719-9260         *This moduct is not manufactured or distilization.

ESOMEPRAZOLE M	rearthum or womting stomach pain se if you are sers may interact with certain	nce than 14 days unless directed by your doct ( Courses (if needed) a 14-day course every 4 months more than 14 days or more often than ever as directed by a doctor set directed by a doctor wettimes be caused by a serious condition.	SATISFACTI GUARANT	HLL, PA 17011
someprazole capsule, delaye	d release			
Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source	) NDC:11822	-0024
Route of Administration	ORAL	item coue (source	)	0024
	Ingredient Name		<b>Basis of Strength</b> ESOMEPRAZOLE	Strengtl 20 mg
<b>ESOMEPRAZOLE</b> (UNII: N3PA65	59F1) (ESOMEPRAZOLE - UNII:1	N3PA6559FT)		20 mg
ESOMEPRAZOLE (UNII: N3PA65 Inactive Ingredients				
Inactive Ingredients	Ingredient Nan			Strength
Inactive Ingredients FD&C BLUE NO. 1 (UNII: H3R47k	Ingredient Nan (3TBD)			
Inactive Ingredients FD&C BLUE NO. 1 (UNII: H3R47k FD&C RED NO. 3 (UNII: PN2ZH51	Ingredient Nan (3TBD) .OQY)			
Inactive Ingredients FD&C BLUE NO. 1 (UNII: H3R47k FD&C RED NO. 3 (UNII: PN2ZH5I FERROSOFERRIC OXIDE (UNII:	Ingredient Nan (3TBD) LOQY) XM0 M8 7F357)			
Inactive Ingredients FD&C BLUE NO. 1 (UNII: H3R47k FD&C RED NO. 3 (UNII: PN2ZH5I FERROSOFERRIC OXIDE (UNII: GELATIN, UNSPECIFIED (UNII: 2	<b>Ingredient Nan</b> (3TBD) LOQY) XM0 M8 7F357) 2G86QN327L)			
Inactive Ingredients FD&C BLUE NO. 1 (UNII: H3R47k FD&C RED NO. 3 (UNII: PN2ZH5I FERROSOFERRIC OXIDE (UNII: GELATIN, UNSPECIFIED (UNII: 2 GLYCERYL MONOSTEARATE (	Ingredient Nan (3TBD) .OQY) XM0M87F357) 2G86QN327L) UNII: 230OU9XXE4)			
Inactive Ingredients FD&C BLUE NO. 1 (UNII: H3R47k FD&C RED NO. 3 (UNII: PN2ZH5I FERROSOFERRIC OXIDE (UNII: GELATIN, UNSPECIFIED (UNII: 2	<b>Ingredient Nan</b> (3TBD) .OQY) XM0M87F357) (386QN327L) (UNII: 230OU9XXE4) <b>D</b> (UNII: 3NXW29V3WO)			
Inactive Ingredients FD&C BLUE NO. 1 (UNII: H3R47F FD&C RED NO. 3 (UNII: PN2ZH5I FERROSOFERRIC OXIDE (UNII: GELATIN, UNSPECIFIED (UNII: 2 GLYCERYL MONOSTEARATE ( HYPROMELLOSE, UNSPECIFIE	<b>Ingredient Nan</b> (3TBD) LOQY) XM0M87F357) 2G86QN327L) (UNII: 230OU9XXE4) <b>D</b> (UNII: 3NXW29V3WO) 70097M6I30)			
Inactive Ingredients FD&C BLUE NO. 1 (UNII: H3R47k FD&C RED NO. 3 (UNII: PN2ZH51 FERROSOFERRIC OXIDE (UNII: GELATIN, UNSPECIFIED (UNII: 2 GLYCERYL MONOSTEARATE ( HYPROMELLOSE, UNSPECIFIE MAGNESIUM STEARATE (UNII: 2	<b>Ingredient Nan</b> (3TBD) LOQY) XM0 M8 7F357) (3G86 QN327L) (UNII: 230 O U9 XXE4) <b>D</b> (UNII: 3NXW29 V3WO) 70 0 9 7M6 I30)	ne		
Inactive Ingredients FD&C BLUE NO. 1 (UNII: H3R47k FD&C RED NO. 3 (UNII: PN2ZH5I FERROSOFERRIC OXIDE (UNII: GELATIN, UNSPECIFIED (UNII: 2 GLYCERYL MONOSTEARATE ( HYPROMELLOSE, UNSPECIFIE MAGNESIUM STEARATE (UNII: 3 MEGLUMINE (UNII: 6 HG8 UB2MU	<b>Ingredient Nan</b> (3TBD) .OQY) XM0M87F357) (2G86QN327L) (UNII: 230OU9XXE4) <b>D</b> (UNII: 3NXW29V3WO) 70097M6I30) (Y) <b>ACRYLATE COPOLYMER (1:1)</b>	ne TYPE A (UNII: NX76LV51		
Inactive Ingredients FD&C BLUE NO. 1 (UNII: H3R47k FD&C RED NO. 3 (UNII: PN2ZH51 FERROSOFERRIC O XIDE (UNII: GELATIN, UNSPECIFIED (UNII: 2 GLYCERYL MONOSTEARATE ( HYPROMELLOSE, UNSPECIFIE MAGNESIUM STEARATE (UNII: 5 MEGLUMINE (UNII: 6 HG8 UB2MU METHACRYLIC ACID - ETHYL #	Ingredient Nan (3TBD) LOQY) XM0M87F357) 2G86QN327L) UNII: 230OU9XXE4) D (UNII: 3NXW29V3WO) 70097M6I30) IY) ACRYLATE COPOLYMER (1:1) SPECIFIED (UNII: 3WJQ0SDW1A	ne TYPE A (UNII: NX76LV51		
Inactive Ingredients FD&C BLUE NO. 1 (UNII: H3R47) FD&C RED NO. 3 (UNII: PN2ZH5I FERROSOFERRIC OXIDE (UNII: GELATIN, UNSPECIFIED (UNII: 2 GLYCERYL MONOSTEARATE ( HYPROMELLOSE, UNSPECIFIE MAGNESIUM STEARATE (UNII: 7 MEGLUMINE (UNII: 6 HG8 UB2MU METHACRYLIC ACID - ETHYL A POLYETHYLENE GLYCOL, UN	Ingredient Nan (3TBD) (OQY) XM0M87F357) (2G86QN327L) (UNII: 230OU9XXE4) D (UNII: 3NXW29V3WO) 70097M6I30) (Y) ACRYLATE COPOLYMER (1:1) SPECIFIED (UNII: 3WJQ0SDW1A	ne TYPE A (UNII: NX76LV51		
Inactive Ingredients FD&C BLUE NO. 1 (UNII: H3R47F FD&C RED NO. 3 (UNII: PN2ZH5I FERROSOFERRIC OXIDE (UNII: GELATIN, UNSPECIFIED (UNII: 2 GLYCERYL MONOSTEARATE ( HYPROMELLOSE, UNSPECIFIE MAGNESIUM STEARATE (UNII: 3 MEGLUMINE (UNII: 6 HG8 UB2MU METHACRYLIC ACID - ETHYL A POLYETHYLENE GLYCOL, UNI POLYSORBATE 80 (UNII: 6 OZP)	<b>Ingredient Nan</b> (3TBD) .OQY) XM0M87F357) (2G86QN327L) (UNII: 230OU9XXE4) D (UNII: 3NXW29V3WO) 70097M6I30) JY) ACRYLATE COPOLYMER (1:1) SPECIFIED (UNII: 3WJQ0SDW1A 39ZG8H)	ne TYPE A (UNII: NX76LV51		
Inactive Ingredients FD&C BLUE NO. 1 (UNII: H3R47) FD&C RED NO. 3 (UNII: PN2ZH5I FERROSOFERRIC OXIDE (UNII: GELATIN, UNSPECIFIED (UNII: 2 GLYCERYL MONOSTEARATE ( HYPROMELLOSE, UNSPECIFIE MAGNESIUM STEARATE (UNII: 4 MEGLUMINE (UNII: 6 HG8 UB2MU METHACRYLIC ACID - ETHYL A POLYETHYLENE GLYCOL, UNI POLYSORBATE 80 (UNII: 60ZP) SHELLAC (UNII: 46N107B710)	<b>Ingredient Nan</b> (3TBD) .OQY) XM0M87F357) (2G86QN327L) (UNII: 230OU9XXE4) D (UNII: 3NXW29V3WO) 70097M6I30) JY) ACRYLATE COPOLYMER (1:1) SPECIFIED (UNII: 3WJQ0SDW1A 39ZG8H)	ne TYPE A (UNII: NX76LV51		
Inactive Ingredients FD&C BLUE NO. 1 (UNII: H3R47) FD&C RED NO. 3 (UNII: PN2ZH51 FERROSOFERRIC OXIDE (UNII: GELATIN, UNSPECIFIED (UNII: 2 GLYCERYL MONOSTEARATE ( HYPROMELLOSE, UNSPECIFIE MAGNESIUM STEARATE (UNII: 5 MEGLUMINE (UNII: 6 HG8 UB2MU METHACRYLIC ACID - ETHYL A POLYETHYLENE GLYCOL, UNS POLYSORBATE 80 (UNII: 6 OZP SHELLAC (UNII: 46 N107B710) SODIUM LAURYL SULFATE (UN	<b>Ingredient Nan</b> (3TBD) .OQY) XM0M87F357) (3G6QN327L) (UNII: 230OU9XXE4) <b>D</b> (UNII: 3NXW29V3WO) 70097M6I30) (Y) <b>ACRYLATE COPOLYMER (1:1)</b> <b>SPECIFIED</b> (UNII: 3WJQ0SDW1A 39ZG8H) NII: 368GB5141J)	ne TYPE A (UNII: NX76LV51		

Product Characteristics						
Color		BLUE	Score		no score	
Shape		CAPSULE	Size		11mm	
Flavor			Imprint Code		7U4	
Contains						
Packaging						
# Item Code		Package Description		Marketing Start Date	Marketing End Date	
1 NDC:11822-0024-5	1 in 1 C	CAPSULE		07/01/2020		
1	14 in 1	in 1 BOTTLE; Type 0: Not a Combination Product				
<b>2</b> NDC:11822-0024-7	3 in 1 0	in 1 CARTON		06/26/2020		
2	14 in 1 BOTTLE; Type 0: Not a Combination Product					
Marketing Information						
Marketing Category	, Ap	oplication Number or Mono	graph Citation	Marketing Start Date	Marketing End Date	
ANDA	AND	A207193		06/26/2020		

# Labeler - Rite Aid Corporation (014578892)

Revised: 7/2020

Rite Aid Corporation