

CARE ONE IBUPROFEN CHILDRENS- ibuprofen suspension
American Sales Company

American Sales Company Ibuprofen Oral Suspension Drug Facts

Active ingredient (in each 5 mL)

Ibuprofen 100 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever,

headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- give with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance
- wash dosage cup after each use

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)**
under 24 lbs	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

**or as directed by a doctor

Other information

- **each 5 mL contains:** sodium 2 mg
- **do not use if printed neckband is broken or missing**
- store at 20-25°C (68-77°F)
- do not freeze

Inactive ingredients

anhydrous citric acid, FD&C red #40, glycerin, high fructose corn syrup, hypromellose, natural and artificial bubblegum flavors, polysorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to the active ingredient in Children's Motrin®

For Ages 2 to 11 Years

Children's

IBUPROFEN Oral Suspension 100mg Per 5mL

Pain Reliever/Fever Reducer (NSAID)

Lasts up to 8hrs

Alcohol Free

dosage cup included

Gluten Free

See New Warnings

OUR PHARMACISTS RECOMMEND

Bubble Gum Flavor

4 FL OZ (118mL)



CARE ONE IBUPROFEN CHILDRENS

ibuprofen suspension

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:41520-166

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	100 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	PINK (light)	Score	
Shape		Size	
Flavor	BUBBLE GUM	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41520-166-26	1 in 1 CARTON	04/28/2004	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:41520-166-28	1 in 1 CARTON	08/11/2013	
2		147 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074937	04/28/2004	

Labeler - American Sales Company (809183973)

Revised: 1/2019

American Sales Company