

**FLU RELIEF THERAPY NIGHT TIME- acetaminophen, diphenhydramine hcl, phenylephrine hcl liquid
Family Dollar (FAMILY WELLNESS)**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Diphenhydramine HCl 25 mg

Phenylephrine HCl 10 mg

Purposes

Pain reliever/fever reducer

Antihistamine/Cough suppressant

Nasal decongestant

Uses

- temporarily relieves
 - minor aches and pains
 - sneezing
 - headache
 - minor sore throat pain
 - itchy, watery eyes
 - nasal and sinus congestion
 - runny nose
 - cough due to minor throat and bronchial irritation
 - itchy nose or throat
- temporarily reduces fever

Warnings

Allergy alert: Acetaminophen may cause severe skin reactions. symptoms may include:

- skin reddening
- blisters
- rash

Liver warning: This product contains acetaminophen. Severe liver damage may occur

if you take:

- more than 6 doses (12 tablespoons or 180 mL) in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other product containing diphenhydramine, even one used on skin
- with any other product containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have

- liver disease
- glaucoma
- heart disease
- thyroid disease
- high blood pressure
- diabetes
- cough accompanied by excessive phlegm (mucus)
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking

- the blood thinning drug warfarin
- sedatives or tranquilizers

When using this product

- **do not use more than directed (see overdose warning)**
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- redness or swelling is present
- nervousness, dizziness, or sleeplessness occurs
- pain, cough or nasal congestion gets worse or lasts more than 7 days

- new symptoms occur
- fever gets worse or lasts more than 3 days
- cough comes back or occurs with a fever, rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) immediately. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device
- mL=milliliter
- keep dosing cup with product
- adults and children 12 years and over
 - 30 mL every 4 hours
- children under 12 years of age; do not use

Other information

- **each 30 mL contains:** potassium 10 mg
- **each 30 mL contains:** sodium 14 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

acesulfame potassium, alcohol, citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavors, glycerin, maltitol solution, propylene glycol, purified water, sodium benzoate, sodium citrate

Questions or comments?

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

Principal Display Panel

Compare to the active ingredients in Theraflu® Nighttime Severe Cold & Cough*

For ages 12 & over

Nighttime

Flu Relief Therapy

Severe Cold and Flu

Acetaminophen 650 mg

Pain reliever; fever reducer

Diphenhydramine HCl 25 mg

Cough suppressant; antihistamine

Phenylephrine HCl 10 mg

Nasal decongestant

Alcohol 10%

Cherry Flavor

FL OZ (mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by Novartis Consumer Health, Inc., owner of the registered trademark Theraflu® Nighttime Severe Cold & Cough.

DISTRIBUTED BY: FAMILY DOLLAR SERVICES, INC

10401 MONROE RD

MATTHEWS, NC 28105 USA

VISIT FAMILYDOLLAR.COM

Product Label

NDC 55319-317-08
COMPARE TO THE ACTIVE INGREDIENTS OF THERAFLU® NIGHTTIME SEVERE COLD & COUGH

FAMILY wellness™

100% SATISFACTION OR YOUR MONEY BACK GUARANTEED

Nighttime Flu Relief Therapy
 Severe Cold & Flu

Pain Reliever/Fever Reducer - **Acetaminophen** 650 mg
 Cough Suppressant/Antihistamine - Diphenhydramine HCl 25 mg
 Nasal Decongestant - Phenylephrine HCl 10 mg

For Relief of:
 Nasal Congestion
 Sore Throat, Runny Nose
 Headache, Body Ache
 Cough, Fever

Adults 12 years and over

CHERRY FLAVOR
 Alcohol Content 10%
 PLD-C265B LB003431

8.3 FL OZ (245 mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by Novartis Consumer Health, Inc., distributor of Theraflu® Nighttime Severe Cold & Cough.

DISTRIBUTED BY:
 FAMILY DOLLAR SERVICES, INC.,
 10401 MONROE RD
 MATH-HEWS, NC 28105 USA
 VISIT FAMILYDOLLAR.COM

Not 100% satisfied? Return package and unused product within 30 days to any Family Dollar store for a refund (with receipt) or exchange.

SKU 0913244

PLD-C265B
 LB003432 0 3 2 2 5 1 0 0 5 0 7 7

Drug Facts

Active ingredients (in each 30 mL) **Purposes**

Acetaminophen 650 mg.....Pain reliever/fever reducer
 Diphenhydramine HCl 25 mg.....Antihistamine/Cough suppressant
 Phenylephrine HCl 10 mg.....Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold
 - minor aches and pains
 - sore throat
 - itchy, watery eyes
 - nasal and sinus congestion
- headache
- runny nose
- sneezing

PEEL CORNER FOR MORE DRUG FACTS ▲

Drug Facts (continued)

- itching of the nose or throat
- cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- to make a child sleepy
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen,

Drug Facts (continued)

ask a doctor or pharmacist.

- with any other product containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- high blood pressure
- heart disease
- diabetes
- thyroid disease
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking

- the blood thinning drug warfarin
- sedatives or tranquilizers

Drug Facts (continued)

When using this product

- do not exceed recommended dosage
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- marked drowsiness may occur
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for

Drug Facts (continued)

adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- keep dosing cup with product
- adults and children 12 years and over
 - 30 mL every 4 hours
- children under 12 years of age: do not use

Other information

- each 30 mL contains: potassium 10 mg
- each 30 mL contains: sodium 14 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients aceulfame potassium, alcohol, citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, maltitol, propylene glycol, purified water, sodium benzoate, sodium citrate

Questions or comments?
 Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Family Wellness Nighttime Flu Relief Therapy

FLU RELIEF THERAPY NIGHT TIME

acetaminophen, diphenhydramine hcl, phenylephrine hcl liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55319-317
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL	

DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg in 30 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ALCOHOL (UNII: 3K9958V90M)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
MALTITOL (UNII: D65DG142WK)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-317-08	245 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/31/2015	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/31/2015	

Labeler - Family Dollar (FAMILY WELLNESS) (024472631)