

**RAPID COMFORT ASPIRIN- aspirin tablet, film coated**  
**Lab Safety Supply, Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**RAPID Comfort™**

***Drug Facts***

**Active ingredient (in each tablet)**

Aspirin (NSAID\*) 325 mg

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/fever reducer

**Uses**

**Temporarily relieves minor aches and pains associated with**

- headache
- minor arthritis pain
- common cold
- menstrual cramps
- muscular aches
- backache
- toothache

**Temporarily reduces fever.**

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- skin reddening
- facial swelling
- rash
- asthma (wheezing)
- blisters
- shock

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcohol drinks every day while using this product
- take more or for a longer time than directed

#### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/ fever reducer
- right before or after heart surgery
- if you are taking prescription drugs for gout, diabetes or arthritis

#### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

#### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other drug

#### **When using this product**

- take with food or milk if stomach upset occurs

#### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- if ringing in the ears or loss of hearing occurs
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### **Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- do not take longer than 10 days, unless directed by a doctor (see Warnings)
- drink a full glass of water with each dose

**Adults and children: (12 years and older)** Take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

**Children under 12 years:** Do not give to children under 12 years of age.

**Other information**

- read all product information before using
- store at room temperature 59°-86°F (15°-30°C)
- avoid excessive heat and humidity
- tamper evident sealed packets
- do not use any opened or torn packets

**Inactive ingredients**

corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide

**Questions or comments? 1-800-356-0783**

**116R LSS -Aspirin 325 mg Label**

Rapid Comfort TM

ASPIRIN

5 grain (325 mg) (NSAID)

Easy to Swallow

Film-Coated Tablets

Pull to Open

Compare active ingredients to:

Bayer®

Registered Trademark of Bayer Consumer

100 Tablets (50x2s)

Tamer-Evident Unit Dose Packets

**RAPIDComfort™**

**ASPIRIN**

5 grain (325mg) (NSAID)

Easy to Swallow  
Film-Coated Tablets

**Drug Facts**

**Active ingredient (in each tablet) Purpose**  
Aspirin 325 mg (NSAID)\*..... Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses**

Temporarily relieves minor aches and pains associated with

- headache
- muscular aches
- minor arthritis pain
- backache
- common cold
- toothache
- menstrual cramps

Temporarily reduces fever.

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
  - skin reddening
  - facial swelling
  - rash
  - asthma (wheezing)
  - blisters
  - shock
- If an allergic reaction occurs, stop use and seek medical help right away.

**Drug Facts (continued)**

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcohol drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery
- if you are taking prescription drugs for gout, diabetes or arthritis

**Ask a doctor before use if**

- stomach bleeding warning applies to you
  - you have a history of stomach problems such as heartburn
  - you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
  - you are taking a diuretic
  - you have asthma
- (Continued on opposite panel)

**RAPIDComfort™**

**ASPIRIN**

5 grain (325mg) (NSAID)

Easy to Swallow  
Film-Coated Tablets

Pull to Open

Compare active ingredient to:

**Bayer®**

Registered Trademark of Bayer Consumer

100 Tablets (50x2s)

Retain carton for complete product information

Tamper-Evident Unit Dose Packets

Rev. 7/8/11, Mfg. 22

**Drug Facts (continued)**

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other drug

When using this product

**Drug Facts (continued)**

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- do not take longer than 10 days, unless directed by a

**RAPIDComfort™**

ASPIRIN

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- if ringing in the ears or loss of hearing occurs
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

doctor (see Warnings)

- drink a full glass of water with each dose

**Adults and children: (12 years and older)** Take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

**Children under 12 years:** Do not give to children under 12 years of age.

**Other information**

- read all product information before using
- store at room temperature 15°-30°C (59°-86°F)
- avoid excessive heat and humidity
- tamper evident sealed packets
- do not use any opened or torn packets

**Inactive ingredients**  
corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide

**Questions or comments?** 1-800-356-0783

# ASPIRIN

## 5 grain (325mg) (NSAID)

Easy to Swallow  
Film-Coated Tablets

Compare active ingredient to:  
**Bayer®**  
Registered Trademark of Bayer Consumer

100 Tablets (50x2s)

© 2011 GHC Specialty Brands, LLC  
1-800-356-0783 • Product #148804/3XKX9

Tamper-Evident Unit Dose Packets

Product Information			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:60683-106
<b>Route of Administration</b>	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg	
Inactive Ingredients			
Ingredient Name	Strength		
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)			
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)			
HYPROMELLOSES (UNII: 3NXW29V3WO)			
MINERAL OIL (UNII: T5L8T28FGP)			
STARCH, CORN (UNII: O8232NY3SJ)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
Product Characteristics			
<b>Color</b>	white (white)	<b>Score</b>	no score
<b>Shape</b>	ROUND (ROUND)	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	TCL;011
<b>Contains</b>			
Packaging			

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:60683-106-13	250 in 1 BOX, UNIT-DOSE		
1		2 in 1 PACKET		
2	NDC:60683-106-33	50 in 1 BOX, UNIT-DOSE		
2		2 in 1 PACKET		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part343	10/14/2011	

**Labeler** - Lab Safety Supply, Inc. (042851410)

### Establishment

Name	Address	ID/FEI	Business Operations
Time Cap Labs, Inc.		037052099	MANUFACTURE

Revised: 10/2011

Lab Safety Supply, Inc.