

**BENZETHONIUM CHLORIDE PLUS DYCLONINE HYDROCHLORIDE- liquid  
bandage liquid  
Geiss Destin & Dunn Inc.**

-----  
**Good Sense Liquid Bandage**

***Active ingredients***

Benzethonium chloride 0.2%

Dyclonine hydrochloride 0.75%

***Purpose***

First aid antiseptic

Topical analgesic

***Uses***

- first aid to prevent bacteria contamination in minor cuts, scrapes and burns
- for the temporary relief of pain

***Warnings***

**For external use only.**

**Flammable:**

Keep away from fire or flame.

**Do not use**

- in the eyes
- over large areas of the body
- longer than 1 week unless directed by a doctor

**Ask a doctor before use if you have**

- deep or puncture wounds
- deep cuts
- animal bites
- serious burns

**When using this product**

- avoid contact with eyes

**Stop use and ask a doctor if**

- conditions persists or gets worse
- infection occurs

- symptoms persist for more than 7 days or clear up and occur again within a few days

**Keep out of reach of children.**

If swallowed, get medical help or call a Poison Control Center right away.

***Directions***

- for adults and children 2 years of age and older
- clean affected area
- apply a small amount on the area 1 to 3 times daily and let dry
- a second coating may be applied for extra protection
- children under 2 years of age: consult a doctor

***Other information***

- **cap bottle tightly**
- store between 20°C to 25°C(68° to 77°F)
- may temporarily sting upon application
- to remove, apply more Liquid Bandage and quickly wipe off

***Inactive ingredients***

acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

***Questions***

**Call 1-866-964-0939**

**Principal Display Panel**

***GOODSENSE***

***Liquid***

***Bandage***

*Topical analgesic and antiseptic seals to protect*

*For small cuts and wounds*

- *Prevents infection*
- *Water-resistant*
- *Invisible and flexible*
- *Built-in applicator brush*

*Use Liquid Bandage for:*

*Blisters, chapped and cracked fingertips, hangnails, paper cuts, shaving nicks and help in preventing the formation of calluses.*

*Great for:*

Athletes, active lifestyles, outdoor sportsmen and musicians.

NET 0.3 FL OZ (9 mL)

NDC 00000-000-00

**GOODSENSE®**

# Liquid Bandage

**Compare to active ingredient in New-Skin\*\***



**Topical analgesic and antiseptic seals to protect For small cuts and wounds**

- Prevents infection
- Water-resistant
- Invisible and flexible
- Built-in applicator brush

0.3 FL OZ (9 mL)

---

**Use Liquid Bandage for:**  
*Blisters, chapped and cracked fingertips, hangnails, paper cuts, shaving nicks and help in preventing the formation of calluses.*

**Great for:**  
*Athletes, active lifestyles, outdoor sportsmen and musicians.*

53-033GS

---

**GOODSENSE®**

# Liquid Bandage

**Topical analgesic and antiseptic seals to protect For small cuts and wounds**

0.3 FL OZ (9 mL)

---

Distributed by: Perrigo Direct, Inc.  
 Peachtree City, GA 30269  
 www.PerrigoDirect.com  
 1-866-964-0939  
 GoodSense® is a registered trademark of L. Perrigo Company.  
 Made in USA with U.S. and imported parts

8 46036 00269 0

**FPO**

\*This product is not manufactured or distributed by Advantice Health, LLC, owner of the registered trademark New-Skin®.

---

**Warnings**  
 For external use only.  
 Flammable: Keep away from fire or flame.  
 Do not use ■ in the eyes ■ over large areas of the body  
 ■ longer than 1 week unless directed by a doctor  
 Ask a doctor before use if you have ■ deep or puncture wounds ■ deep cuts  
 ■ animal bites ■ serious burns  
 When using this product ■ avoid contact with eyes  
 Stop use and ask a doctor if ■ condition persists or gets worse ■ infection occurs  
 ■ symptoms persist for more than 7 days or clear up and occur again within a few days  
 Keep out of reach of children. If swallowed, get medical help or call a Poison Control Center right away.

**Directions**  
 ■ for adults and children 2 years of age and older ■ clean affected area  
 ■ apply a small amount on the area 1 to 3 times daily and let dry ■ a second coating may be applied for extra protection ■ children under 2 years of age: consult a doctor

**Other information**  
 ■ cap bottle tightly ■ store between 20° to 25°C (68° to 77°F) ■ may temporarily sting upon application ■ to remove, apply more Liquid Bandage and quickly wipe off

**Inactive ingredients**  
 acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

**Questions? Call 1-866-964-0939**

**Care** ■ Wipe top of bottle clean after each use.

**Uses**  
 ■ first aid to prevent bacterial contamination in minor cuts, scrapes and burns  
 ■ for the temporary relief of pain

**DO NOT USE IF NECKBAND IS BROKEN**

**Drug Facts**

**Active Ingredients**  
Benzethonium chloride 0.2%  
Dyclonine hydrochloride 0.75%

**Purpose**  
First aid antiseptic  
Topical analgesic

NDC 0000-000-00

**GOODSENSE®**

**Liquid Bandage**

*Topical analgesic and antiseptic seals to protect*

*For small cuts and wounds*

*Water-resistant*

0.3 FLOZ (9 mL)

**DO NOT USE IF NECKBAND IS BROKEN**

**Active ingredients:** Benzethonium Chloride 0.2% and Dyclonine Hydrochloride 0.75% **Purpose:** First aid antiseptic and topical analgesic **Uses:** First aid to prevent bacterial contamination in minor cuts, scrapes, and burns. For the temporary relief of pain.

**Warnings:** For external use only. **Flammable:** Keep away from fire or flame. **Do not use** in eyes, over large areas of the body, and longer than 1 week unless directed by a doctor.

**Ask a doctor before use if you have** deep or puncture wounds, deep cuts, animal bites and serious burns.

**When using this product** avoid contact with eyes.

**Stop use and ask a doctor** if condition persists or gets worse, symptoms persist for more than 7 days or clear up and occur again within a few days, and infection occurs.

**Keep out of reach of children.** If swallowed, get medical help or call a Poison Control Center right away.

**Directions:** For adults and children 2 years of age and older. Clean affected area. Apply a small amount on the area 1 to 3 times daily and let dry. A second coating may be applied for extra protection. Children under 2 years of age: consult a doctor.

**Other information: Cap bottle tightly.** Store between 20° to 25°C (68° to 77°F). May temporarily sting upon application. To remove, apply more Liquid Bandage and quickly wipe off.

**Inactive ingredients:** acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

**Questions? Call 1-866-964-0939**

**MADE IN USA WITH U.S. AND IMPORTED PARTS**  
Distributed by: Perrigo Direct, Inc. Peachtree, GA 30269  
www.PerrigoDirect.com

50-033GS

## BENZETHONIUM CHLORIDE PLUS DYCLONINE HYDROCHLORIDE

liquid bandage liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50804-755
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BENZETHONIUM CHLORIDE</b> (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	0.2 mg in 9 mL
<b>DYCLONINE HYDROCHLORIDE</b> (UNII: ZEC193879Q) (DYCLONINE - UNII:078A24Q300)	DYCLONINE HYDROCHLORIDE	0.75 mg in 9 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ACETONE</b> (UNII: 1364PS73AF)	
<b>AMYL ACETATE</b> (UNII: 92Q24NH7AS)	
<b>CASTOR OIL</b> (UNII: D5340Y219G)	
<b>ETHYL ACETATE</b> (UNII: 76845O8NMZ)	

**PYROXYLIN** (UNII: KYR8BR2X6O)

**ALCOHOL** (UNII: 3K9958V90M)

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50804-755-03	1 in 1 CARTON	07/09/2021	
1		9 mL in 1 BOTTLE, WTH APPLICATOR; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	07/09/2021	

**Labeler** - Geiss Destin & Dunn Inc. (076059836)

Revised: 1/2024

Geiss Destin & Dunn Inc.