

**BENZETHONIUM CHLORIDE PLUS DYCLONINE HYDROCHLORIDE- liquid bandage liquid**  
**Geiss Destin & Dunn Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Good Sense Liquid Bandage**

***Active ingredients***

Benzethonium chloride 0.2%

Dyclonine hydrochloride 0.75%

***Purpose***

First aid antiseptic

Topical analgesic

***Uses***

- first aid to prevent bacteria contamination in minor cuts, scrapes and burns
- for the temporary relief of pain

***Warnings***

**For external use only.**

**Flammable:**

Keep away from fire or flame.

**Do not use**

- in the eyes
- over large areas of the body
- longer than 1 week unless directed by a doctor

**Ask a doctor before use if you have**

- deep or puncture wounds
- deep cuts
- animal bites
- serious burns

**When using this product**

- avoid contact with eyes

**Stop use and ask a doctor if**

- conditions persists or gets worse
- infection occurs
- symptoms persist for more than 7 days or clear up and occur again within a few days

**Keep out of reach of children.**

If swallowed, get medical help or call a Poison Control Center right away.

**Directions**

- for adults and children 2 years of age and older
- clean affected area
- apply a small amount on the area 1 to 3 times daily and let dry
- a second coating may be applied for extra protection
- children under 2 years of age: consult a doctor

**Other information**

- **cap bottle tightly**
- store between 20°C to 25°C(68° to 77°F)
- may temporarily sting upon application
- to remove, apply more Liquid Bandage and quickly wipe off

**Inactive ingredients**

acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

**Questions**

**Call 1-866-964-0939**

**Principal Display Panel****GOODSENSE****Liquid****Bandage**

*Topical analgesic and antiseptic seals to protect*

*For small cuts and wounds*

- *Prevents infection*
- *Water-resistant*
- *Invisible and flexible*
- *Built-in applicator brush*

*Use Liquid Bandage for:*


Blisters, chapped and cracked fingertips, hangnails, paper cuts, shaving nicks and help in preventing the formation of calluses.

Great for:

Athletes, active lifestyles, outdoor sportsmen and musicians.

NET 0.3 FL OZ (9 mL)

NDC 00000-000-00



# Liquid Bandage

**Topical analgesic and antiseptic seals to protect**  
**For small cuts and wounds**

- Prevents infection
- Invisibile and flexible

- Water-resistant
- Built-in applicator brush



0.3 FL OZ (9 mL)

**Compare to active ingredient in New-Skin\*\***

**Use Liquid Bandage for:**  
*Blisters, chapped and cracked fingertips, hangnails, paper cuts, shaving nicks and help in preventing the formation of calluses.*

**Great for:**  
*Athletes, active lifestyles, outdoor sportsmen and musicians.*

53-033GS

---

GOODSENSE®

**Liquid Bandage**


*For small cuts and wounds*

## Liquid Bandage

**Topical analgesic and antiseptic seals to protect**  
**For small cuts and wounds**

0.3 FL OZ (9 mL)

Distributed by: Perrigo Direct, Inc.  
Peachtree City, GA 30269  
www.PerrigoDirect.com  
1-866-964-0939  
GoodSense® is a registered trademark of L. Perrigo Company.  
Made in USA with U.S. and imported parts



8 46036 00269 0

\*This product is not manufactured or distributed by Advantice Health, LLC, owner of the registered trademark New-Skin®.

---

**Care** ■ Wipe top of bottle clean after each use.

**Questions? Call 1-866-964-0939**

**Inactive ingredients** ■ acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

**Other information** ■ cap bottle tightly ■ store between 20° to 25°C (68° to 77°F) ■ may temporarily sting upon application ■ to remove, apply more Liquid Bandage and quickly wipe off

**Directions** ■ for adults and children 2 years of age and older ■ clean affected area ■ apply a small amount on the area 1 to 3 times daily and let dry ■ a second coating may be applied for extra protection ■ children under 2 years of age: consult a doctor

**Keep out of reach of children.** If swallowed, get medical help or call a Poison Control Center right away.

**Stop use and ask a doctor if** ■ condition persists or gets worse ■ infection occurs

**When using this product** ■ avoid contact with eyes

**Ask a doctor before use if you have** ■ deep or puncture wounds ■ deep cuts ■ animal bites ■ serious burns

■ longer than 1 week unless directed by a doctor

**DO NOT USE IF NECKBAND IS BROKEN**

**Drug Facts**

**Active Ingredients**  
Benzethonium chloride 0.2%  
Dyclonine hydrochloride 0.75%  
.....First aid antiseptic  
.....Topical analgesic

**Uses**  
■ first aid to prevent bacterial contamination in minor cuts, scrapes and burns  
■ for the temporary relief of pain

**Warnings**  
For external use only.  
Flammable: Keep away from fire or flame.  
Do not use ■ in the eyes ■ over large areas of the body

**DO NOT USE IF NECKBAND IS BROKEN**

NDC 00000-000-00

**GOODSENSE®**

**Liquid Bandage**

**Topical analgesic and antiseptic seals to protect**

**For small cuts and wounds**

**Water-resistant**

0.3 FL OZ (9 mL)

**Active ingredients:** Benzethonium Chloride 0.2% and Dyclonine Hydrochloride 0.75% **Purpose:** First aid antiseptic and topical analgesic **Uses:** First aid to prevent bacterial contamination in minor cuts, scrapes, and burns. For the temporary relief of pain.

**Warnings:** For external use only. **Flammable:** Keep away from fire or flame. **Do not use** in eyes, over large areas of the body, and longer than 1 week unless directed by a doctor.

**Ask a doctor before use if you have** deep or puncture wounds, deep cuts, animal bites and serious burns.  
**When using this product** avoid contact with eyes.  
**Stop use and ask a doctor** if condition persists or gets worse, symptoms persist for more than 7 days or clear up and occur again within a few days, and infection occurs.  
**Keep out of reach of children.** If swallowed, get medical help or call a Poison Control Center right away.

**Directions:** For adults and children 2 years of age and older. Clean affected area. Apply a small amount on the area 1 to 3 times daily and let dry. A second coating may be applied for extra protection. Children under 2 years of age: consult a doctor.

**Other information: Cap bottle tightly.** Store between 20° to 25°C (68° to 77°F). May temporarily sting upon application. To remove, apply more Liquid Bandage and quickly wipe off.

**Inactive ingredients:** acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

**Questions? Call 1-866-964-0939**

**MADE IN USA WITH U.S. AND IMPORTED PARTS**  
Distributed by: Perrigo Direct, Inc. Peachtree, GA 30269  
www.PerrigoDirect.com

50-033GS

## BENZETHONIUM CHLORIDE PLUS DYCLONINE HYDROCHLORIDE

liquid bandage liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50804-755
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BENZETHONIUM CHLORIDE</b> (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	0.2 mg in 9 mL
<b>DYCLONINE HYDROCHLORIDE</b> (UNII: ZEC193879Q) (DYCLONINE - UNII:078A24Q300)	DYCLONINE HYDROCHLORIDE	0.75 mg in 9 mL

### Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

<b>ACETONE</b> (UNII: 1364PS73AF)	
<b>AMYL ACETATE</b> (UNII: 92Q24NH7AS)	
<b>CASTOR OIL</b> (UNII: D5340Y2I9G)	
<b>ETHYL ACETATE</b> (UNII: 76845O8NMZ)	
<b>PYROXYLIN</b> (UNII: KYR8BR2X6O)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50804-755-03	1 in 1 CARTON	07/09/2021	
1		9 mL in 1 BOTTLE, WTH APPLICATOR; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	07/09/2021	

**Labeler** - Geiss Destin & Dunn Inc. (076059836)

Revised: 1/2023

Geiss Destin & Dunn Inc.