SUPHEDRINE PE SINUS PRESSURE PLUS PAIN- acetaminophen, phenylephrine hcl tablet

ARMY AND AIR FORCE EXCHANGE SERVICE

Exchange Select 44-502-Sinus Pressure Plus Pain

Active ingredients (in each caplet)

Acetaminophen 325 mg Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer Nasal decongestant

Uses

- temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies, and the common cold:
 - minor aches and pains
 - nasal congestion
 - headache
 - sinus congestion and pressure
- promotes sinus drainage
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

 with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- heart disease
- thyroid disease
- diabetes
- liver disease
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- adults and children 12 years and over
 - take 2 caplets every 4 hours
 - do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, croscarmellose sodium, crospovidone, FD&C yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, povidone, silicon dioxide, stearic acid

Questions or comments?

1-800-426-9391

Principal display panel

exchange **∕select**™

Pseudoephedrine Free

Compare To The Active Ingredients of SUDAFED PE® SINUS PRESSURE + PAIN*

SUPHEDRINE PE SINUS PRESSURE + PAIN

Acetaminophen, Phenylephrine HCl Pain Reliever/Fever Reducer, Nasal Decongestant

- Maximum Strength
- Nasal Congestion
- Sinus Headache
- Sinus Pressure

NON-DROWSY

Actual Size

24 Caplets

✓ quality value

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark SUDAFED PE® SINUS PRESSURE + PAIN.

"SATISFACTION GUARANTEED OR YOUR MONEY BACK"

Manufactured For Your Military Exchanges Distributed by: LNK International, Inc. Hauppauge, NY 11788 1-800-426-9391

50844 REV0820D50208

SUPHEDRINE PE

SINUS PRESSURE + PAIN



TAMPER EVIDENT: DO NOT USE
IF PACKAGE IS OPENED OR IF
BLISTER UNIT IS TORN,
BROKEN OR SHOWS
ANY SIGNS OF TAMPERING

exchange select

Pseudoephedrine Free

Compare To The Active Ingredients of SUDAFED PE® SINUS PRESSURE + PAIN*

PHEDRIN

Acetaminophen, Phenylephrine HCI

Pain Reliever/Fever Reducer, Nasal Decongestant

- Maximum Strength
- Nasal Congestion
- Sinus Headache
- Sinus Pressure

NON-DROWSY

quality value

print/No varnish area Lot no/Exp date

B-0066E-502-08-F REV0820D50208

Drug Facts

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Active ingredients (in each caplet)

Actual Size

Caplets

Purpose Phenylephrine HCl 5 mg Nasal decongestant

Drug Facts (continued)

temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies, and the common cold:
■ minor aches and pains ■ nasal congestion ■ headache

- sinus congestion and pressure

Drug Facts (continued)

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away
- Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- \blacksquare if you have ever had an allergic reaction to this product or any of its

Ask a doctor before use if you have ■ heart disease

■ thyroid disease ■ diabetes ■ liver disease ■ high blood pressure difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are taking the blood thinning

When using this product do not exceed recommended dosage.

Drug Facts (continued)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
 new symptoms occur These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions on do not use more than directed

- adults and children 12 years and over
- take 2 caplets every 4 hours
- do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

Other information ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients corn starch, croscarmellose sodium, crospovidone, FD&C yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, povidone, silicon dioxide, stearic acid

Questions or comments? 1-800-426-9391

Exchange Select 44-502 REV0820D

SUPHEDRINE PE SINUS PRESSURE PLUS PAIN

acetaminophen, phenylephrine hcl tablet

Product Information					
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55301-902		
Route of Administration	ORAL				

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	

Inactive Ingredients			
Ingredient Name	Strength		
STARCH, CORN (UNII: O8232NY3SJ)			
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)			
CROSPOVIDONE, UNSPECIFIED (UNII: 2S7830E561)			
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)			
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)			
SILICON DIOXIDE (UNII: ETJ7Z 6XBU4)			
STEARIC ACID (UNII: 4ELV7Z65AP)			

Product Characteristics				
Color	orange	Score	no score	
Shape	OVAL	Size	18mm	
Flavor		Imprint Code	44;502	
Contains				

Packaging						
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:55301- 902-08	2 in 1 CARTON	09/16/2021			
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product				

Marketing Information			
Marketing Category			Marketing End Date
OTC Monograph Drug	M012	09/16/2021	

Establishment				
Name	Address	ID/FEI	Business Operations	
LNK International, Inc.		832867837	manufacture(55301-902) , pack(55301-902)	

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(55301-902)

Revised: 9/2023 ARMY AND AIR FORCE EXCHANGE SERVICE