

**DAYTIME NON DROWS COLD AND FLU AND NIGHTTIME COLD AND-
acetaminophen, dextromethorphan hydrobromide, doxylamine succinate,
phenylephrine hydrochloride
Spirit Pharmaceuticals LLC**

Daytime Non Drowsy Cold and Flu and Nighttime Cold and Flu

Active ingredients (in each softgel)

COLD & FLU NON-DROWSY DAY RELIEF

Acetaminophen 325 mg
Dextromethorphan hydrobromide 10 mg
Phenylephrine hydrochloride 5 mg

COLD & FLU NIGHT RELIEF

Acetaminophen 325 mg
Dextromethorphan hydrobromide 10 mg
Doxylamine succinate 6.25 mg

Purposes

COLD & FLU NON DROWSY DAY RELIEF

Pain reliever/fever reducer
Cough suppressant
Nasal decongestant

COLD & FLU NIGHT RELIEF

Pain reliever/fever reducer
Cough suppressant
Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- fever
- headache
- minor aches and pain
- cough due to minor throat and bronchial irritation
- sore throat
- nasal congestion (Daytime only)
- runny nose and sneezing (Nighttime only)

Warnings

Liver warning This product contains acetaminophen. Severe liver damage may occur if you take: ● more than 4 doses in 24 hours, which is the maximum daily amount for this product ● with other drugs containing acetaminophen ● 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ● skin reddening ● blisters ● rash
If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- to make a child sleepy (Nighttime only)

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- liver disease
- trouble urinating due to enlarged prostate gland
- diabetes (Daytime only)
- heart disease (Daytime only)
- thyroid disease (Daytime only)
- high blood pressure (Daytime only)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema (Daytime only)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema (Nighttime only)
- glaucoma (Nighttime only)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (Nighttime only)

When using this product

- do not take more than directed
- marked drowsiness may occur (Nighttime only)
- avoid alcoholic drinks (Nighttime only)
- excitability may occur, especially in children (Nighttime only)
- be careful when driving a motor vehicle or operating machinery (Nighttime only)
- alcohol, sedatives, and tranquilizers may increase drowsiness (Nighttime only)

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless (Daytime only)
- pain, nasal congestion, or cough gets worse or lasts more than 7 days (Daytime only)
- pain or cough gets worse or lasts more than 7 days (Nighttime only)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

Directions

- when using other DAYTIME and NIGHTTIME products, carefully read each label to ensure correct dosing

Directions (Daytime only)

- take only as directed - see Overdose warning
- do not exceed 4 doses per 24 hours

adults & children 12 years & over	take 2 softgels with water every 4 hours
children 4 to under 12 years	ask a doctor
children under 4 years of age	do not use

- when using other DAYTIME and NIGHTTIME products, carefully read each label to ensure correct dosing

Directions (Nighttime only)

- take only as directed - see Overdose warning
- do not exceed 4 doses per 24 hours

adults & children 12 years & over	take 2 softgels with water every 6 hours
children 4 to under 12 years	ask a doctor
children under 4 years of age	do not use

Other information

- store at room temperature.

Inactive ingredients

DAYTIME

FD&C Red# 40, FD&C Yellow# 6, gelatin, glycerin, polyethylene glycol, lecithin, myglyol, povidone, propylene glycol, purified water, sorbitol sorbitan, titanium dioxide

NIGHTTIME

D&C Yellow# 10, FD&C Blue# 1, gelatin, glycerin, myglyol, lecithin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan, polysorb, sorbitol sorbitan, titanium dioxide

Questions or comments?

1-888-333-9792

Principal Display Panel



DAYTIME
NON-DROWSY

COLD & FLU

Acetaminophen

Pain Reliever / Fever Reducer
Dextromethorphan HBr
Cough Suppressant
Phenylephrine HCl
Nasal Decongestant

MULTI-SYMPTOM

NIGHTTIME

COLD & FLU

Acetaminophen

Pain Reliever / Fever Reducer
Dextromethorphan HBr
Cough Suppressant
Doxylamine Succinate
Antihistamine

MULTI-SYMPTOM

COMBO PACK



Compare to Vicks' DayQuil® &
NyQuil® Cold & Flu LiquiCaps®
active ingredients†

DAYTIME
NON-DROWSY

COLD & FLU

Acetaminophen

Pain Reliever / Fever Reducer
Dextromethorphan HBr
Cough Suppressant
Phenylephrine HCl
Nasal Decongestant

MULTI-SYMPTOM

NIGHTTIME

COLD & FLU

Acetaminophen

Pain Reliever / Fever Reducer
Dextromethorphan HBr
Cough Suppressant
Doxylamine Succinate
Antihistamine

MULTI-SYMPTOM



Actual Size

48 total softgels**
(**Liquid-filled capsules)

32 softgels 16 softgels



Actual Size

Drug Facts	
Active ingredients for Daytime (in each softgel)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant
Active ingredients for Nighttime (in each softgel)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 15 mg	Cough suppressant
Doxylamine succinate 0.25 mg	Antihistamine
Uses	
<ul style="list-style-type: none"> temporarily relieves common cold/flu symptoms: fever headache sore throat minor aches and pains cough due to minor throat and bronchial irritation nasal congestion (Daytime only) runny nose and sneezing (Nighttime only) 	
Warnings	
<p>Liver warning This product contains acetaminophen. Severe liver damage may occur if you take:</p> <ul style="list-style-type: none"> more than 4 doses in 24 hours, which is the maximum daily amount for this product with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> skin reddening blisters rash <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Sore throat warning If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p>	
Drug Facts (continued)	
<p>Do not use</p> <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. if you have ever had an allergic reaction to this product or any of its ingredients to make a child sleepy (Nighttime only) <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> cough that occurs with too much phlegm (mucus) trouble urinating due to enlarged prostate gland diabetes (Daytime only) heart disease (Daytime only) thyroid disease (Daytime only) high blood pressure (Daytime only) persistent or chronic cough such as occurs with smoking, asthma, or emphysema (Daytime only) a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema (Nighttime only) glaucoma (Nighttime only) <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> taking the blood thinning drug warfarin taking sedatives or tranquilizers (Nighttime only) <p>When using this product</p> <ul style="list-style-type: none"> do not take more than directed marked drowsiness may occur (Nighttime only) avoid alcoholic drinks (Nighttime only) excitability may occur, especially in children (Nighttime only) be careful when driving a motor vehicle or operating machinery (Nighttime only) alcohol, sedatives, and tranquilizers may increase drowsiness (Nighttime only) 	

Drug Facts (continued)	
<p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> you get nervous, dizzy or sleepless (Daytime only) pain, nasal congestion, or cough gets worse or lasts more than 7 days (Daytime only) pain or cough gets worse or lasts more than 7 days (Nighttime only) fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur cough comes back or occurs with rash or headache that lasts <p>These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.</p> <p>Overdose warning Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.</p>	
Directions (Daytime only)	
<ul style="list-style-type: none"> take only as directed - see Overdose warning do not exceed 4 doses per 24 hours adults and children 12 years and over: take 2 softgels with water every 4 hours children 4 to under 12 years: ask a doctor children under 4 years: do not use when using other DAYTIME and NIGHTTIME products, carefully read each label to ensure correct dosing 	
Drug Facts (continued)	
Directions (Nighttime only)	
<ul style="list-style-type: none"> take only as directed - see Overdose warning do not exceed 4 doses per 24 hours adults and children 12 years and over: take 2 softgels with water every 6 hours children 4 to under 12 years: ask a doctor children under 4 years: do not use when using other DAYTIME and NIGHTTIME products, carefully read each label to ensure correct dosing 	
Other information	
<ul style="list-style-type: none"> store at room temperature 	
Inactive ingredients (Daytime only)	
FD&C Red No. 40, FD&C Yellow No. 8, gelatin, glycerin, lecithin, mylglycol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan, titanium dioxide	
Inactive ingredients (Nighttime only)	
D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, lecithin, mylglycol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan, titanium dioxide	
Questions or comments? Call 1-888-333-8782	

DAYTIME NON DROWS COLD AND FLU AND NIGHTTIME COLD AND acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68210-5008
---------------------	----------------	---------------------------	----------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68210-5008-4	1 in 1 CARTON; Type 1: Convenience Kit of Co-Package	12/12/2022	

Quantity of Parts		
Part #	Package Quantity	Total Product Quantity
Part 1	1 BLISTER PACK	8
Part 2	1 BLISTER PACK	8

Part 1 of 2

DAYTIME NON DROWSY COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled

Product Information	
Item Code (Source)	NDC:68210-5006
Route of Administration	ORAL

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE	5 mg

Inactive Ingredients	
Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SORBITOL (UNII: 506T60A25R)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	red	Score	no score
Shape	OVAL	Size	21mm
Flavor		Imprint Code	512;A09;AP01

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		4 in 1 CARTON		
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/12/2022	

Part 2 of 2**NIGHTTIME COLD AND FLU**

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride capsule, liquid filled

Product Information

Item Code (Source)	NDC:68210-5007
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POVIDONE (UNII: FZ989GH94E)	

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	green	Score	no score
Shape	OVAL	Size	21mm
Flavor		Imprint Code	116;A07;AP02
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		2 in 1 CARTON		
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/19/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/19/2022	

Labeler - Spirit Pharmaceuticals LLC (179621011)