

IBUPROFEN MINI- ibuprofen mini 200 mg capsule, liquid filled
TARGET CORPORATION

Target 11673 683 Ibuprofen 200 mg Mini liquid filled capsules

DRUG FACTS

Active ingredient (in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*

(present as the free acid and potassium salt)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Uses

temporarily relieves minor aches and pains due to:

- headache
- toothache
- backache
- menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis

temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood-thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin,

ibuprofen, naproxen, or others]

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke.

These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

you have symptoms of heart problems or stroke:

- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling

pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days

redness or swelling is present in the painful area

any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist

if pain or fever does not respond to 1 capsule, 2 capsules may be used

do not exceed 6 capsules in 24 hours, unless directed by a doctor

- children under 12 years: ask a doctor

Other information

- **each capsule contains:** potassium 20 mg
- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients ammonium hydroxide, FD&C green no. 3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, shellac, sorbitan monooleate, sorbitol sorbitan

Questions or comments? call **1-800-910-6874**

Ibuprofen Minis
200 mg Solubilized Ibuprofen Capsules
Pain Reliever / Fever Reducer (NSAID)
Minis - smaller capsule, same strength
up&up
80 SOFTGELS** (**LIQUID-FILLED CAPSULES)

NDC 11673-683-80

DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Uses ■ temporarily relieves minor aches and pains due to: headache, toothache, backache, menstrual cramps, the common cold, muscular aches, minor pain of arthritis
■ temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs (containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ **do not take more than directed** ■ **the smallest effective dose should be used** ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor

Other information
■ Children under 12 years: ask a doctor

each capsule contains: potassium 20 mg
Questions or comments? Call 1-800-910-6874

094-015-008 R00 Dist. by Target Corp. Target Brands, Inc.
C-022262-01-034-0000 Mpls., MN 554-03 7001.1023
Code No.: 60/DRIUGS/515
4q127003 **Made in India**
TM & © 2024

Lot No.:
Exp. Date:

Varnish Omit Area

Ibuprofen Minis

200 mg Solubilized Ibuprofen Capsules
Pain Reliever / Fever Reducer (NSAID)
Minis – smaller capsule, same strength



160 SOFTGELS** (**LIQUID-FILLED CAPSULES)

NDC:11673-683-661

DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Uses

- temporarily relieves minor aches and pains due to: headache, toothache, backache, menstrual cramps, the common cold, muscular aches, minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions **do not take more than directed**
the smallest effective dose should be used
adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist

if pain or fever does not respond to 1 capsule, 2 capsules may be used

do not exceed 6 capsules in 24 hours, unless directed by a doctor

children under 12 years: ask a doctor

Other information
each capsule contains: potassium 20 mg

Questions or comments? Call 1-800-910-6874

094-01 7840 R00 Made in India 700T 1023

C-002262-01-034-0000 TM & ©2024

Dist. by Target Corp. Target Brands, Inc. Mpls., MN 55403

40127103 Code Inc.: 6010US515

Lot No.:
Exp. Date:

Varnish Omit Area

Compare to active ingredient in Advil® Liqui-Gels® minis*

Ibuprofen Minis

200 mg Solubilized Ibuprofen Capsules
Pain Reliever / Fever Reducer (NSAID)
Minis – smaller capsule, same strength



Actual Size

300 SOFTGELS** (**LIQUID-FILLED CAPSULES)

NDC 11673-683-32

DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each capsule)

Solubilized Ibuprofen (NSAID)*
200 mg (present as the free acid and potassium salt)
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued under label)

This product is not manufactured or distributed by Haleon group of companies, owner of the registered trademark Advil® and Cabotant Pharma Solutions, Inc., owner of the registered trademark Liqui-Gels®.

094-01 8119 R00
C-002262-01-034-0000
Distributed by Target Corporation
Minneapolis, MN 55403
Made in India

700T 1023
40127103 Code Inc.: 6010US515

Lot No.:

Exp. Date:

Varnish Omit Area

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

right before or after heart surgery

if you have problems or serious side effects from taking pain relievers or fever reducers

you have a history of stomach problems, such as heartburn

you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke

you are taking a diuretic

Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition

taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin

taking any other drug

When using this product:

take with food or milk if stomach upset occurs

Stop use and ask a doctor if:

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

you have symptoms of heart problems or stroke:

- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling

pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days

redness or swelling is present in the painful area

any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions **do not take more than directed**
adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist

if pain or fever does not respond to 1 capsule, 2 capsules may be used

do not exceed 6 capsules in 24 hours, unless directed by a doctor

children under 12 years: ask a doctor

Other information
each capsule contains: potassium 20 mg

Questions or comments? Call 1-800-910-6874

094-01 7840 R00 Made in India 700T 1023

C-002262-01-034-0000 TM & ©2024

Dist. by Target Corp. Target Brands, Inc. Mpls., MN 55403

40127103 Code Inc.: 6010US515

Lot No.:

Exp. Date:

Varnish Omit Area

Adhesive Area

RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Drug Facts

Active ingredient (in each capsule)	Purpose
Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)* Pain reliever/Fever reducer (present as the free acid and potassium salt) *nonsteroidal anti-inflammatory drug	

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

Drug Facts (continued)

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Code No.: 6018055515

ibuprofen
Minis
200 mg Solubilized Ibuprofen Capsules
Pain Reliever / Fever Reducer (NSAID)

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

Drug Facts (continued)

When using this product


- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely

DO NOT USE IF SEAL UNDER BOTTLE CAP OR PROTECTIVE SEAL UNDER BOTTLE CAP PROTECTION IS BROKEN OR MISSING



Lot No.:
Exp Date:
Varnish Omit

Drug Facts (continued)

directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains: potassium 20 mg
- read all warnings and directions before use

Drug Facts (continued)

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

ammonium hydroxide, FD&C green no. 3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, shellac, sorbitan monooleate, sorbitol sorbitan

Questions or comments? Call 1-800-910-6874

Satisfaction guaranteed – Love it or your money back.

700T.1023

Distributed by Target Corporation
Minneapolis, MN 55403
Made in India
TM & ©2024 Target Brands, Inc.

*This product is not manufactured or distributed by Haleon group of companies, owner of the registered trademark Advil® and Catalent Pharma Solutions, Inc., owner of the registered trademark Liqui-Gels®.

4003503

Compare to active ingredient in Advil® Liqui-Gels® minis*

Ibuprofen Minis

NDC 11673-683-66



200 mg Solubilized Ibuprofen Capsules
Pain Reliever / Fever Reducer (NSAID)

Minis – smaller capsule, same strength


160 SOFTGELS**
(**LIQUID-FILLED CAPSULES)



Varnish/Print Omit Area

Varnish Omit

Varnish Omit



IBUPROFEN MINI

ibuprofen mini 200 mg capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-683
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
GELATIN (UNII: 2G86QN327L)	
WATER (UNII: 059QF0KO0R)	
SORBITAN MONOOLEATE (UNII: 06XEA2VD56)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
FERROUS OXIDE (UNII: G7036X8B5H)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
AMMONIA (UNII: 5138Q19F1X)	
SORBITOL (UNII: 506T60A25R)	
SHELLAC (UNII: 46N107B71O)	

Product Characteristics

Color	green	Score	no score
Shape	CAPSULE	Size	6mm
Flavor		Imprint Code	62
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-683-32	300 in 1 BOTTLE; Type 0: Not a Combination Product	12/17/2021	
2	NDC:11673-683-80	1 in 1 CARTON	12/17/2021	
2		80 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:11673-683-66	1 in 1 CARTON	12/17/2021	
3		160 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079205	12/17/2021	

Labeler - TARGET CORPORATION (006961700)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	manufacture(11673-683)

Revised: 1/2024

TARGET CORPORATION