

ZICAM COLD REMEDY ZINC AND ELDERBERRY LOZENGE- zinc acetate anhydrous, zinc gluconate, and sambucus nigra flowering top lozenge Church & Dwight Co., Inc.

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

Zicam® Cold Remedy Zinc and Elderberry Lozenge

Active ingredients (in each lozenge)

Zincum aceticum 2x

Zincum gluconicum 2x

Sambucus nigra (elderberry) 4x

Purpose

Active ingredients (in each tablet)	Purpose
Zincum aceticum 2x	Reduces duration of the common cold and helps relieve nasal congestion
Zincum gluconicum 2x	
Sambucus nigra (elderberry) 4x	Helps relieve cold symptoms *

* Claims based on traditional homeopathic practice, not accepted medical evidence. Not FDA evaluated.

Uses

Reduces duration of the common cold and helps relieve nasal congestion

Zicam® Cold Remedy was not formulated to be effective for flu and allergies.

Warnings

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

If you are allergic or sensitive to zinc, consult a doctor before using.

Stop use and ask a doctor if symptoms, including cough, persist, recur or are

accompanied by a fever, rash, or persistent headache, or if new symptoms occur. A persistent cough that lasts for more than one week may be a sign of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **for best results, use at the first sign of a cold and continue to use until symptoms completely subside**
- adults and children 12 years of age and older:
 - take 1 tablet at the onset of symptoms. Completely dissolve lozenge in mouth. Do not chew. Do not swallow whole.
- repeat every 2 - 3 hours, not to exceed 7 tablets in 24 hours. Take until symptoms are gone.
- to avoid minor stomach upset, do not take on an empty stomach
- do not eat or drink for 15 minutes after use. Do not eat or drink citrus fruits or juices for 30 minutes before or after use. Otherwise, drink plenty of fluids.
- children under 12 years of age: ask a doctor before use

Other information

- store between 15-29°C (59-84°F)
- avoid freezing and excessive heat

Inactive ingredients

citric acid, corn syrup, FD&C Blue No. 1, FD&C Red No. 40, flavor, glycerin, malic acid, purified water, soy lecithin, sucralose, sucrose

Questions? Comments?

call 877-942-2626 toll free

www.zicam.com

PRINCIPAL DISPLAY PANEL - 25 INDIVIDUALLY WRAPPED CENTER FILLED LOZENGES

NEW!

Plus ELDERBERRY

ZICAM®

COLD REMEDY

PRE-COLD®

FIGHTER
 ZINC &
 ELDERBERRY
 LOZENGE
 Elderberry Flavor
 HOMEOPATHIC
 TAKE AT
 THE FIRST SIGN
 25 INDIVIDUALLY WRAPPED
 CENTER FILLED
 LOZENGES



NEW! Plus **ELDERBERRY**





SHORTENS COLDS

DO NOT USE IF SEALED INNER POUCH IS DAMAGED OR TORN.
 This product is not required to go through the FDA's New Drug Application approval process.

Drug Facts	
Active ingredients (in each lozenge)	Purpose
Zincum acetatum 2x	Reduces duration of the common cold
Zincum gluconicum 2x	and helps relieve nasal congestion
Sambucus nigra 4x (elderberry)	Helps relieve cold symptoms†
Uses	
Reduces duration of the common cold and helps relieve nasal congestion	
<small>Zicam® Cold Remedy was not formulated to be effective for flu or allergies.</small>	
Warnings	
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	
If you are allergic or sensitive to zinc, consult a doctor before using.	
Stop use and ask a doctor if symptoms, including cough, persist, recur or are accompanied by a fever, rash, or persistent headache, or if new symptoms occur. A persistent cough that lasts for more than one week may be a sign of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions	
■ for best results, use at the first sign of a cold and continue to use until symptoms completely subside	
■ adults and children 12 years of age and older:	
■ take 1 lozenge at the onset of symptoms. Completely dissolve lozenge in mouth. Do not chew. Do not swallow whole.	
■ repeat every 2–3 hours, not to exceed 7 lozenges in 24 hours. Take until symptoms are gone.	
■ to avoid minor stomach upset, do not take on an empty stomach	
■ do not eat or drink for 15 minutes after use. Do not eat or drink citrus fruits or juices for 30 minutes before or after use. Otherwise, drink plenty of fluids.	
■ children under 12 years of age: ask a doctor before use	
Other information	
■ store between 15–29°C (59–84°F)	
■ avoid freezing and excessive heat.	
Inactive ingredients citric acid, corn syrup, FD&C Blue No. 1, FD&C Red No. 40, flavor, glycerin, malic acid, purified water, soy lecithin, sucralose, sucrose	

GET OVER YOUR COLD FASTER™

- Great tasting elderberry flavor
- Non-drowsy
- Non-habit forming

SATISFACTION GUARANTEED

Questions? Comments? call 877-942-2626 toll free www.zicam.com

KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

†Claims based on traditional homeopathic practice, not accepted medical evidence. Not FDA evaluated.

DISTRIBUTED BY CHURCH & DWIGHT CO., INC. EWING, NJ 08628



TAKE AT THE FIRST SIGN

Elderberry Flavor

25 INDIVIDUALLY WRAPPED CENTER FILLED LOZENGES

Take every 2-3 HOURS until symptoms are gone.

That first snuffle, sneeze or throat tickle... you have a Pre-Cold® the first sign a full blown cold is coming.

Take Zicam® Lozenges now to shorten a cold.

GO FROM PRE-COLD® TO NO COLD FASTER.

Zicam® Cold Remedy products are available in a wide range of forms and flavors.

70101541
ZCFC-30182-01

HOMEOPATHIC
HOMEOPATHIC
HOMEOPATHIC

ZICAM COLD REMEDY ZINC AND ELDERBERRY LOZENGE

zinc acetate anhydrous, zinc gluconate, and sambucus nigra flowering top lozenge

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:10237-462
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ZINC ACETATE ANHYDROUS (UNII: H2ZEY72PME) (ZINC CATION - UNII:13S1S8SF37)	ZINC CATION	2 [hp_X]
ZINC GLUCONATE (UNII: U6WSN5SQ1Z) (ZINC CATION - UNII:13S1S8SF37)	ZINC GLUCONATE	2 [hp_X]
SAMBUCUS NIGRA FLOWERING TOP (UNII: CT03BSA18U) (SAMBUCUS NIGRA FLOWERING TOP - UNII:CT03BSA18U)	SAMBUCUS NIGRA FLOWERING TOP	4 [hp_X]

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID ACETATE (UNII: DSO12WL7AU)	
CORN SYRUP (UNII: 9G5L16BK6N)	
WATER (UNII: 059QF0KO0R)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
METHYL SALICYLATE (UNII: LAV5U5022Y)	
GLYCERIN (UNII: PDC6A3C0OX)	
MALIC ACID (UNII: 817L1N4CKP)	
SUCROSE (UNII: C151H8M554)	

Product Characteristics

Color	PURPLE	Score	no score
Shape	ROUND	Size	15mm
Flavor	BERRY	Imprint Code	Z
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:10237-462-25	1 in 1 CARTON	08/02/2021	
1		25 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
UNAPPROVED HOMEOPATHIC		08/02/2021	

Labeler - Church & Dwight Co., Inc. (001211952)

