

**MUCINEX DM- guaifenesin and dextromethorphan hydrobromide tablet, extended release**  
**Select Corporation**

-----  
**Mucinex® DM**

**Drug Facts**

---

| <b>Active ingredients (in each extended-release bi-layer tablet)</b> | <b>Purpose</b>    |
|--|-------------------|
| Dextromethorphan HBr 30 mg   | Cough suppressant |
| Guaifenesin 600 mg   | Expectorant       |

---

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
  - the intensity of coughing
  - the impulse to cough to help you get to sleep

**Warnings**

**Do not use**

- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

**When using this product**

- do not use more than directed

**Stop use and ask a doctor if**

- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a

Poison Control Center right away.

### **Directions**

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: 1 or 2 tablets every 12 hours; not more than 4 tablets in 24 hours
- children under 12 years of age: do not use

### **Other information**

- tamper evident: do not use if carton is damaged or if pouch is broken or torn
- store at 20-25°C (68-77°F)

### **Inactive ingredients**

carbomer homopolymer type B; D&C Yellow #10 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

### **Questions?**

**1-866-MUCINEX (1-866-682-4639)** You may also report side effects to this phone number.

Product Repackaged and Distributed with permission of manufacturer by: **Select Corporation**, Carrollton, TX 75007

Distributed by: Reckitt Benckiser  
Parsippany, NJ 07054-0224

### **PRINCIPAL DISPLAY PANEL - 2 Tablet Pouch Blister Pack Carton**

Mucinex® DM

600 mg guaifenesin & 30 mg dextromethorphan HBr  
extended-release bi-layer tablets

EXPECTORANT & COUGH SUPPRESSANT

12 HOUR

- ✓ Controls Cough
- ✓ Thins And Loosens Mucus

20 Pouches –  
2 tablets each

- ← extended-release
- ← immediate-release



**Mucinex<sup>®</sup>DM**  
600 mg guaifenesin & 30 mg dextromethorphan HBr  
extended-release bi-layer tablets  
**EXPECTORANT & COUGH SUPPRESSANT**




- ✓ Controls Cough
- ✓ Thins And Loosens Mucus

**Mucinex<sup>®</sup>DM**  
600 mg guaifenesin & 30 mg dextromethorphan HBr  
extended-release bi-layer tablets  
**EXPECTORANT & COUGH SUPPRESSANT**



- ✓ Controls Cough
- ✓ Thins And Loosens Mucus

20 pouches —  
2 tablets each




**Mucinex<sup>®</sup>DM**  
600 mg guaifenesin & 30 mg dextromethorphan HBr  
extended-release bi-layer tablets  
**EXPECTORANT & COUGH SUPPRESSANT**



↑  
TO OPEN  
PUSH IN TAB AND PULL OUT

- ✓ Controls Cough
- ✓ Thins And Loosens Mucus

20 pouches —  
2 tablets each



**Mucinex<sup>®</sup>DM**  
600 mg guaifenesin & 30 mg dextromethorphan HBr  
extended-release bi-layer tablets  
**EXPECTORANT & COUGH SUPPRESSANT**



**Drug Facts**

**Active Ingredients** **Purpose**  
**(in each extended-release bi-layer tablet)**  
Dextromethorphan HBr 30 mg...Cough suppressant  
Guaifenesin 600 mg.....Expectorant

**Uses** ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive  
■ temporarily relieves:  
■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants  
■ the intensity of coughing  
■ the impulse to cough to help you get to sleep

**Warnings**  
**Do not use** ■ for children under 12 years of age  
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

**Drug Facts (continued)**

**When using this product**  
■ do not use more than directed  
**Stop use and ask a doctor if**  
■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

■ **pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
■ do not crush, chew, or break tablet  
■ take with a full glass of water  
■ this product can be administered without regard for timing of meals  
■ adults and children 12 years and older:  
1 or 2 tablets every 12 hours; not more than 4 tablets in 24 hours  
■ children under 12 years of age: do not use

**Other information**



**12 HOUR**

✓ Controls Cough  
✓ Thins And Loosens Mucus

20 pouches —  
2 tablets each

← extended-release  
← immediate-release

depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

■ tamper evident: do not use if carton is damaged or if pouch is broken or torn

■ store at 20-25°C (68-77°F)

**Inactive ingredients** carbomer homopolymer type B; D&C Yellow #10 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

**Questions? 1-888-MUCINEX (1-888-682-4639)**  
You may also report side effects to this phone number.

**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

Product Repackaged and Distributed with permission of manufacturer by: **Select Corporation** Carrollton, TX 75007

Distributed by: **Reckitt Benckiser** Parsippany, NJ 07054-0224 ©2014 RB. Made in England 081514



1401650227

## MUCINEX DM

guaifenesin and dextromethorphan hydrobromide tablet, extended release

### Product Information

|                                |                |                           |                              |
|--------------------------------|----------------|---------------------------|------------------------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:52904-650(NDC:63824-056) |
| <b>Route of Administration</b> | ORAL           |                           |                              |

### Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength             | Strength |
|--|-------------------------------|----------|
| <b>guaifenesin</b> (UNII: 495W7451VQ) (guaifenesin - UNII:495W7451VQ)                        | guaifenesin                   | 600 mg   |
| <b>dextromethorphan hydrobromide</b> (UNII: 9D2RTI9KYH) (dextromethorphan - UNII:7355X3ROTS) | dextromethorphan hydrobromide | 30 mg    |

### Inactive Ingredients

| Ingredient Name  | Strength |
|--|----------|
| <b>Carbomer Homopolymer Type B (Allyl Pentaerythritol Crosslinked)</b> (UNII: HHT01Z NK31) |          |
| <b>D&amp;C yellow no. 10</b> (UNII: 35SW5USQ3G)  |          |
| <b>aluminum oxide</b> (UNII: LMI26O6933)   |          |
| <b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)  |          |
| <b>magnesium stearate</b> (UNII: 70097M6I30)   |          |
| <b>microcrystalline cellulose</b> (UNII: OP1R32D61U)                                       |          |

### Product Characteristics

|                 |                          |                     |             |
|-----------------|--------------------------|---------------------|-------------|
| <b>Color</b>    | WHITE (yellow and white) | <b>Score</b>        | no score    |
| <b>Shape</b>    | OVAL                     | <b>Size</b>         | 16mm        |
| <b>Flavor</b>   |                          | <b>Imprint Code</b> | Mucinex;600 |
| <b>Contains</b> |                          |                     |             |

**Packaging**

| # | Item Code        | Package Description                             | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:52904-650-03 | 1 in 1 BLISTER PACK                             | 06/26/2012           |                    |
| 1 |                  | 2 in 1 POUCH; Type 0: Not a Combination Product |                      |                    |
| 2 | NDC:52904-650-20 | 20 in 1 CARTON                                  | 06/26/2012           |                    |
| 2 |                  | 2 in 1 POUCH; Type 0: Not a Combination Product |                      |                    |

**Marketing Information**

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| NDA                | NDA021620                                | 06/26/2012           |                    |

**Labeler** - Select Corporation (053805599)

Revised: 4/2022

Select Corporation