

NAUSEACALM- anamirta cocculus seed, ipecac, strychnos nux-vomica seed, tobacco leaf, kerosene tablet

Boiron

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

NauseaCalm Tablets

Active ingredient** (in each tablet)

Cocculus indicus 4C HPUS (0.5 mg) (contains less than 10⁻¹⁰ mg picrotoxinin)

Ipecacuanha 4C HPUS (0.5 mg) (contains less than 10⁻¹⁰ mg emetine alkaloids)

Nux Vomica 4C HPUS (0.5 mg) (contains less than 10⁻¹⁰ strychnine alkaloids)

Tabacum 4C HPUS (0.5 mg) (contains less than 10⁻¹⁰ nicotine alkaloids)

Petroleum 4C HPUS (0.5 mg) (contains less than 10⁻¹⁰ aromatic hydrocarbons)

The letters "HPUS" indicate that the components in this product are officially monographed in the Homeopathic Pharmacopoeia of the United States.

Purpose*

Cocculus indicus 4C HPUS ... Relieves nausea or vomiting with dizziness and the need to lie down

Ipecacuanha 4C HPUS ... Relieves nausea and vomiting with hypersalivation

Nux Vomica 4C HPUS ... Relieves upset stomach, indigestion, nausea, and vomiting

Tabacum 4C HPUS ... Relieves dizziness, nausea and vomiting improved by fresh air

Petroleum 4C HPUS ... Relieves nausea improved by eating

Uses*

- for relief of nausea and vomiting associated with:
 - upset stomach
 - fullness due to overindulgence in food and drink
 - stomach flu
 - for relief of nausea, vomiting, or dizziness associated with motion from travel, amusement rides, or video games

Stop use and ask a doctor if nausea or vomiting is accompanied by
diarrhea

severe headache

fever

severe abdominal pain or cramping

Stop use and ask a doctor if symptoms persist for more than 2 days or worsen.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Adults and children 7 years of age and older - At the onset of symptoms, dissolve 2 tablets in the mouth. Repeat if necessary until symptoms improve, up to 3 times per day.

Children 2 to 6 years of age - Dissolve 2 tablets in 1 tablespoon of water for easier intake. Follow directions above.

Children under 2 years of age - Ask a doctor.

croscarmellose sodium, lactose, magnesium stearate

do not use if glued carton end flaps are open or if the blister seal is broken

store below 86° F (30° C)

Nausea Relief*

No Aspirin or Caffeine

60 Meltaway Tablets unflavored

No Known Drug Interactions

*CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE, NOT ACCEPTED MEDICAL EVIDENCE. NOT FDA EVALUATED.

**C,K,CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.

Questions or Comments?

BoironUSA.com

Info@Boiron.com

1-800-BOIRON-1

(1-800-264-7661)

Made in France

Distributed by Boiron Inc.

Newtown Square, PA 19073

Drug Facts

Active ingredients* (in each tablet)	Purpose*
Cocculus indicus 4C HPUS (0.5 mg) (contains less than 10 ⁻⁶ mg picrotoxinin)	Relieves nausea or vomiting with dizziness and the need to lie down
Ipecacuanha 4C HPUS (0.5 mg) (contains less than 10 ⁻⁶ mg emetine alkaloids)	Relieves nausea and vomiting with hypersalivation
Nux vomica 4C HPUS (0.5 mg) (contains less than 10 ⁻⁶ mg strychnine alkaloids)	Relieves upset stomach, indigestion, nausea, and vomiting
Tabacum 4C HPUS (0.5 mg) (contains less than 10 ⁻⁶ mg nicotine alkaloids)	Relieves dizziness, nausea and vomiting improved by fresh air
Petroleum 4C HPUS (0.5 mg) (contains less than 10 ⁻⁶ mg aromatic hydrocarbons)	Relieves nausea improved by eating

The letters "HPUS" indicate that the components in this product are officially monographed in the Homeopathic Pharmacopoeia of the United States.

Uses*

- for relief of nausea and vomiting associated with:
 - upset stomach
 - fullness due to overindulgence in food and drink
 - stomach flu
- for relief of nausea, vomiting, or dizziness associated with motion from travel, amusement rides, or video games

Warnings

- Ask a doctor before use if nausea or vomiting is accompanied by
- severe headache
 - fever
 - severe abdominal pain or cramping
 - diarrhea

Stop use and ask a doctor if symptoms persist for more than 2 days or worsen.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Adults and children 7 years of age and older	At the onset of symptoms, dissolve 2 tablets in the mouth. Repeat if necessary until symptoms improve, up to 3 times per day.
----------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------

Drug Facts (continued)

Directions (continued)

Children 2 to 6 years of age	Dissolve 2 tablets in 1 tablespoon of water for easier intake. Follow directions above.
Children under 2 years of age	Ask a doctor.

Other Information

- do not use if glued carton end flaps are open or if the blister seal is broken
- store below 86°F (30°C)

Inactive ingredients

croscarmellose sodium, lactose, magnesium stearate

Questions or comments?

BoironUSA.com
Info@Boiron.com
1-800-808-0801
(1-800-264-7661)

Made in France
Distributed by Boiron Inc.
Newtown Square, PA 19073
please recycle

NAUSEA RELIEF*

**Upset Stomach • Vomiting
Stomach Flu***



NauseaCalm[®]
MELTAWAY TABLETS

HOMEOPATHIC MEDICINE

- No Known Drug Interactions
- No Aspirin or Caffeine



BOIRON[®]

**World Leader in
Homeopathic Medicines**

60
MELTAWAY
TABLETS
Unflavored

*CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE, NOT ACCEPTED MEDICAL EVIDENCE. NOT FDA EVALUATED.
*C, K, CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.

NAUSEA RELIEF

NauseaCalm[®]

Adults and children 7 years of age and older: At the onset of symptoms, dissolve 2 tablets in the mouth. Repeat if necessary until symptoms improve, up to 3 times per day.
Children 2 to 6 years of age: Dissolve 2 tablets in 1 tablespoon of water for easier intake. Follow directions above. Children under 2 years of age: Ask a doctor.



NAUSEA RELIEF

NauseaCalm[™]

Adults and children 7 years of age and older: At the onset of symptoms, dissolve 2 tablets in the mouth. Repeat if necessary until symptoms improve, up to 3 times per day.
Children 2 to 6 years of age: Dissolve 2 tablets in 1 tablespoon of water for easier intake. Follow directions above. Children under 2 years of age: Ask a doctor.



Drug Facts

Active Ingredients* (in each tablet) **Purpose***

Cocculus indicus 4C HPUS (0.5 mg) Relieves nausea or vomiting with dizziness and the need to lie down (contains less than 10⁻¹⁰ mg picrotoxinin)

Ipecacuanha 4C HPUS (0.5 mg) Relieves nausea and vomiting (contains less than 10⁻¹⁰ mg emetine alkaloids) with hypersalivation

Nux vomica 4C HPUS (0.5 mg) Relieves upset stomach, indigestion, nausea, and vomiting (contains less than 10⁻¹⁰ mg strychnine alkaloids)

Tobacum 4C HPUS (0.5 mg) Relieves dizziness, nausea and vomiting improved by fresh air (contains less than 10⁻¹⁰ mg nicotine alkaloids)

Petroleum 4C HPUS (0.5 mg) Relieves nausea improved by eating (contains less than 10⁻¹⁰ mg aromatic hydrocarbons)

The letters "HPUS" indicate that the components in this product are officially monographed in the Homeopathic Pharmacopoeia of the United States.

Uses*

- for relief of nausea and vomiting associated with:
 - upset stomach
 - fullness due to overindulgence in food and drink
 - stomach flu
- for relief of nausea, vomiting, or dizziness associated with motion from travel, amusement rides, or video games

Warnings

Stop use and ask a doctor if nausea or vomiting is accompanied by ■ diarrhea
■ severe headache ■ fever ■ severe abdominal pain or cramping

Stop use and ask a doctor if symptoms persist for more than 2 days or worsen.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Adults and children 7 years of age and older	At the onset of symptoms, dissolve 2 tablets in the mouth. Repeat if necessary until symptoms improve, up to 3 times per day.
----------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------

Drug Facts (continued)

Directions (continued)

Children 2 to 6 years of age	Dissolve 2 tablets in 1 tablespoon of water for easier intake. Follow directions above.
Children under 2 years of age	Ask a doctor.

Other information

- do not use if glued carton end flaps are open or if the blister seal is broken
- store below 86°F (30°C)

Inactive ingredients
 croscarmellose sodium, lactose, magnesium stearate

Questions or comments?
 BoironUSA.com
 Info@Boiron.com
 1-800-BOIRON-1
 (1-800-264-7661)

Made in France
 Distributed by Boiron Inc.
 Newtown Square, PA 19073

♻️ please recycle

NDC 0220-9244-04

NAUSEA RELIEF*

Upset Stomach Stomach Flu*

NauseaCalm™

MELTAWAY TABLETS

HOMEOPATHIC MEDICINE

- Relieves Nausea & Vomiting*
- No Aspirin or Caffeine

World Leader in
Homeopathic Medicines

*CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE, NOT ACCEPTED MEDICAL EVIDENCE. NOT FDA EVALUATED.
 *C, X, GK, and X are homeopathic dilutions; see BoironUSA.com/info for details.

NAUSEACALM

anamirta cocculus seed, ipecac, strychnos nux-vomica seed, tobacco leaf, kerosene tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0220-9244
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ANAMIRTA COCCULUS SEED (UNII: 810258W28U) (ANAMIRTA COCCULUS SEED - UNII:810258W28U)	ANAMIRTA COCCULUS SEED	4 [hp_C]
KEROSENE (UNII: 1C89KKC04E) (KEROSENE - UNII:1C89KKC04E)	KEROSENE	4 [hp_C]
TOBACCO LEAF (UNII: 6YR2608RSU) (TOBACCO LEAF - UNII:6YR2608RSU)	TOBACCO LEAF	4 [hp_C]
IPECAC (UNII: 62I3C8233L) (IPECAC - UNII:62I3C8233L)	IPECAC	4 [hp_C]
STRYCHNOS NUX-VOMICA SEED (UNII: 269XH13919) (STRYCHNOS NUX-VOMICA SEED - UNII:269XH13919)	STRYCHNOS NUX-VOMICA SEED	4 [hp_C]

Inactive Ingredients

Ingredient Name	Strength
LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	9mm
Flavor		Imprint Code	Boiron
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0220-9244-04	1 in 1 CARTON	07/29/2021	
1		60 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved homeopathic		07/29/2021	

Labeler - Boiron (282560473)

Registrant - Boiron Inc. (014892269)

Establishment

Name	Address	ID/FEI	Business Operations
Boiron		282560473	manufacture(0220-9244)

Revised: 5/2023

Boiron