

**DAYTIME NIGHTTIME COLD FLU RELIEF- acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl
Family Dollar (FAMILY WELLNESS)**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

DRUG FACTS

Active ingredients for Daytime (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Active ingredients for Nighttime (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

Purpose for Daytime

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Purpose for Nighttime

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

DAYTIME

- **temporarily relieves common cold and flu symptoms**
 - **cough due to minor throat and bronchial irritation**
 - **nasal congestion**
 - **headache**
 - **minor aches and pains**

- **fever**
- **sore throat**

NIGHTTIME

- **temporarily relieves common cold and flu symptoms**
 - **cough due to minor throat and bronchial irritation**
 - **sore throat**
 - **headache**
 - **minor aches and pains**
 - **fever**
 - **runny nose and sneezing**

Warnings

DAYTIME

Liver warning: These products contain acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using these products

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

NIGHTTIME

Liver warning: This product contain acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or following by fever, headache, rash, nausea, vomiting, consult a doctor promptly.

Do not use

DAYTIME

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

NIGHTTIME

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

DAYTIME

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

NIGHTTIME

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

DAYTIME

taking the blood thinning drug warfarin

NIGHTTIME

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product,

DAYTIME

do not use more than directed

NIGHTTIME

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

DAYTIME

- nervousness, dizziness or sleeplessness occur
- pain, cough, and nasal congestion gets worse or lasts more than 7 days
- redness or swelling is present
- new symptoms occur
- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

NIGHTTIME

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash headache that lasts.

These could be a signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

DAYTIME

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over; take 2 softgels with water every 4 hours.
- children under 12 years: do not use

NIGHTTIME

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- children under 12 years: do not use

Other information

- store between 15°-30°C (59°-86°F)
- avoid excessive heat

Inactive ingredients

DAYTIME

butylated hydroxyanisole, butylated hydroxytoluene, carminic acid*, D&C yellow #10*, edible white ink, FD&C red #40*, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sodium metabisulfite*, sorbitan*, sorbitol, *contain one or more of these ingredient

NIGHTTIME

D&C yellow #10, edible white ink, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol *may contain this ingredient

Questions or comments?

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

Principal Display Panel

DAYTIME

MULTI-SYMPTOM

Cold & Flu Relief

DayTime / Non-Drowsy

Acetaminophen Pain Reliever/Fever Reducer

Dextromethorphan HBr Cough Suppressant

Phenylephrine HCl Nasal Decongestant

SOFTGELS

NIGHTTIME

Acetaminophen Pain Reliever/Fever Reducer

Dextromethorphan HBr Cough Suppressant

Doxylamine succinate Antihistamine

SOFTGELS

†This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® , DayQuil, NyQuil®, and LiquiCaps® are registered trademarks of the Procter and Gamble Company.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOW SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

When using Daytime and Nighttime products, carefully read the labeling to ensure correct dosing

DISTRIBUTED BY: MIDWOOD BRANDS,, llc

10611 MONROE RD, MATTHEWS, NC 28105 USA

Product Label

Keep out of reach of children. Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

If pregnant or breast-feeding, ask a health professional before use.

Signs of a serious condition:

- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present ■ new symptoms occur
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- nervousness, dizziness, or sleeplessness occur

Stop use and ask a doctor if:

- you have ■ liver disease ■ heart disease
- cough that occurs with too much phlegm (mucus)
- diabetes ■ thyroid disease ■ high blood pressure
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- trouble urinating due to an enlarged prostate gland
- asthma, chronic bronchitis, or emphysema
- trouble urinating due to an enlarged prostate gland

When using this product, do not use more than directed.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking the product.

Ask a doctor before use if you have ■ liver disease ■ heart disease ■ cough that occurs with too much phlegm (mucus) ■ diabetes ■ thyroid disease ■ high blood pressure ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ trouble urinating due to an enlarged prostate gland ■ asthma, chronic bronchitis, or emphysema ■ trouble urinating due to an enlarged prostate gland

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking the product.

Ask a doctor before use if you have ■ liver disease ■ heart disease ■ cough that occurs with too much phlegm (mucus) ■ diabetes ■ thyroid disease ■ high blood pressure ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ trouble urinating due to an enlarged prostate gland ■ asthma, chronic bronchitis, or emphysema ■ trouble urinating due to an enlarged prostate gland

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking the product.

Ask a doctor before use if you have ■ liver disease ■ heart disease ■ cough that occurs with too much phlegm (mucus) ■ diabetes ■ thyroid disease ■ high blood pressure ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ trouble urinating due to an enlarged prostate gland ■ asthma, chronic bronchitis, or emphysema ■ trouble urinating due to an enlarged prostate gland

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking the product.

DayTime Cold & Flu Softgel

Active Ingredients (in each softgel)

- Acetaminophen 325 mg
- Pain reliever/fever reducer
- Cough suppressant
- Dextromethorphan HBr 10 mg
- Nasal decongestant
- Phenylephrine HCl 5 mg
- Antihistamine

Purposes

- temporarily relieves common cold and flu symptoms
- cough due to minor throat and bronchial irritation
- sore throat
- runny nose & sneezing
- fever
- minor aches and pains
- headache
- minor aches and pains ■ fever ■ sore throat

Warnings

- Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours
- or more other drugs containing acetaminophen
- with other drugs containing acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking the product.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking the product.

Ask a doctor or pharmacist

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking the product.

Ask a doctor before use if you have ■ liver disease ■ heart disease ■ cough that occurs with too much phlegm (mucus) ■ diabetes ■ thyroid disease ■ high blood pressure ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ trouble urinating due to an enlarged prostate gland ■ asthma, chronic bronchitis, or emphysema ■ trouble urinating due to an enlarged prostate gland

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking the product.

Ask a doctor before use if you have ■ liver disease ■ heart disease ■ cough that occurs with too much phlegm (mucus) ■ diabetes ■ thyroid disease ■ high blood pressure ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ trouble urinating due to an enlarged prostate gland ■ asthma, chronic bronchitis, or emphysema ■ trouble urinating due to an enlarged prostate gland

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking the product.

NightTime Cold & Flu Softgel

Active Ingredients (in each softgel)

- Acetaminophen 325 mg
- Pain reliever/fever reducer
- Cough suppressant
- Dextromethorphan HBr 15 mg
- Nasal decongestant
- Phenylephrine HCl 5 mg
- Antihistamine

Purposes

- temporarily relieves common cold and flu symptoms
- cough due to minor throat and bronchial irritation
- sore throat
- runny nose & sneezing
- fever
- minor aches and pains
- headache
- minor aches and pains ■ fever ■ runny nose & sneezing ■ sore throat ■ cough due to minor throat and bronchial irritation

Warnings

- Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours
- or more other drugs containing acetaminophen
- with other drugs containing acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking the product.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking the product.

Ask a doctor or pharmacist

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking the product.

Ask a doctor before use if you have ■ liver disease ■ heart disease ■ cough that occurs with too much phlegm (mucus) ■ diabetes ■ thyroid disease ■ high blood pressure ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ trouble urinating due to an enlarged prostate gland ■ asthma, chronic bronchitis, or emphysema ■ trouble urinating due to an enlarged prostate gland

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking the product.

Ask a doctor before use if you have ■ liver disease ■ heart disease ■ cough that occurs with too much phlegm (mucus) ■ diabetes ■ thyroid disease ■ high blood pressure ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ trouble urinating due to an enlarged prostate gland ■ asthma, chronic bronchitis, or emphysema ■ trouble urinating due to an enlarged prostate gland

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking the product.

COMPARE TO THE ACTIVE INGREDIENTS IN VICKS® DAYQUIL® AND NYQUIL® COLD & FLU LIQUICAPS®

NDC 59319-472-24

MULTI-SYMP TOM

Cold & Flu Relief

DAYTIME/NON-DROWSY

Acetaminophen
Pain Reliever/Fever Reducer

Cough Suppressant
Dextromethorphan HBr
Nasal Decongestant
Phenylephrine HCl

16 SOFTGELS

NIGHT TIME

Acetaminophen
Pain Reliever/Fever Reducer

Cough Suppressant
Dextromethorphan HBr
Antihistamine
Doxylamine Succinate

8 SOFTGELS

OTC network

FAMILY wellness™

100% SATISFACTION OR YOUR MONEY BACK GUARANTEED

PARENTS: TAMPER EVIDENT DO NOT USE IF CARTON IS OPENED OR IF BUSTER UNIT IS DOWN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING. KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

www.StaysWellness.com

When using Daytime and Nighttime products, carefully read the labeling to ensure correct dosing.

Lot No:
Exp. Date:

DISTRIBUTED BY: MIDWOOD BRANDS, LLC
10611 MOUNOIE RD., MATTHEWS, NC 28105

NOT 100% SATISFIED? Return package and unused product within 30 days to any Family Dollar store for a refund (with receipt) or exchange.

PLU 065
1000000

SKU 090066

0 52251 11623 0

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over: take 2 softgels with water every 4 hours
- children under 12 years: do not use

Other Information

- store between 15-30°C (59-86°F) ■ avoid excessive heat

Inactive ingredients D&C yellow #10, white ink, FD&C blue #1, sorbitol, glycerin, polyethylene glycol, purified water, hydroxytoluene, carmine acid, D&C yellow #10, white ink, FD&C red #40, FD&C yellow #5, green, glycerin, polyethylene glycol, sorbitol, propylene glycol, purified water, sodium metabisulfite, sorbitan, sorbitol

Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- children under 12 years: do not use

Other Information

- store between 15-30°C (59-86°F) ■ avoid excessive heat

Inactive ingredients D&C yellow #10, white ink, FD&C blue #1, sorbitol, glycerin, polyethylene glycol, purified water, hydroxytoluene, carmine acid, D&C yellow #10, white ink, FD&C red #40, FD&C yellow #5, green, glycerin, polyethylene glycol, sorbitol, propylene glycol, purified water, sodium metabisulfite, sorbitan, sorbitol

Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

FAMILY WELLNESS DayTime NightTime Multi-symptom Cold & Flu

DAYTIME NIGHTTIME COLD FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55319-472
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-472-24	1 in 1 CARTON; Type 0: Not a Combination Product	03/31/2016	03/31/2025

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	8 BLISTER PACK	8
Part 2	16 BLISTER PACK	16

Part 1 of 2

NIGHTTIME COLD FLU RELIEF

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule

Product Information

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	green	Score	no score
Shape	CAPSULE	Size	21mm
Flavor		Imprint Code	P30;94A;SCU1;215;P120
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 CARTON		
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	03/31/2016	

Part 2 of 2

DAYTIME COLD FLU RELIEF

acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule

Product Information

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
FD&C RED NO. 40 (UNII: WZB9127XOA)
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)
GELATIN (UNII: 2G86QN327L)
GLYCERIN (UNII: PDC6A3C0OX)
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)
POVIDONE (UNII: FZ989GH94E)
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)
WATER (UNII: 059QF0KO0R)
SORBITAN (UNII: 6O92ICV9RU)
SORBITOL (UNII: 506T60A25R)
CARMINIC ACID (UNII: CID8Z8N95N)
SODIUM METABISULFITE (UNII: 4VON5FNS3C)
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)

Product Characteristics

Color	orange	Score	no score
Shape	CAPSULE	Size	21mm
Flavor		Imprint Code	P19;95A;512;P119
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		16 in 1 CARTON		
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	03/31/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	03/31/2016	03/31/2025

Labeler - Family Dollar (FAMILY WELLNESS) (024472631)