

PAIN RELIEF IBUPROFEN- ibuprofen tablet, film coated
Rite Aid Corporation

Rite Aid 44-292

Active ingredient (in each brown caplet)

Ibuprofen USP, 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - menstrual cramps
 - toothache
 - the common cold
 - backache
 - headache
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- skin reddening
- asthma (wheezing)
- rash
- facial swelling
- shock
- blisters
- hives

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - leg swelling
 - slurred speech
 - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20

weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 caplet, 2 caplets may be used
 - do not exceed 6 caplets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- see end flap for expiration date and lot number

Inactive ingredients

carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

NDC 11822-0292-4

Compare to the
active ingredient in

Advil® Caplets†

PAIN RELIEF

IBUPROFEN

IBUPROFEN TABLETS USP, 200 mg

PAIN RELIEVER/FEVER REDUCER
(NSAID)

ACTUAL SIZE

500

CAPLETS**

(**CAPSULE-SHAPED TABLETS)

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed by PF
Consumer Healthcare 1 LLC, owner of the registered trademark
Advil® Caplets. 50844 REV1221A29214

DISTRIBUTED BY:

RITE AID, 200 NEWBERRY COMMONS
ETTERS, PA 17319 www.riteaid.com

SATISFACTION GUARANTEE:

If you're not satisfied, we'll
happily refund your money.

NDC 11822-0292-4

Compare to the active ingredient in Advil® Caplets†

PAIN RELIEF

IBUPROFEN

IBUPROFEN TABLETS USP, 200 mg

PAIN RELIEVER/FEVER REDUCER (NSAID)

ACTUAL SIZE

500 CAPLETS**

(**CAPSULE-SHAPED TABLETS)

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each brown caplet)
Ibuprofen USP 200 mg (NSAID)*
*nonsteroidal anti-inflammatory drug

Purpose
Pain reliever/fever reducer

Uses
■ temporarily relieves minor aches and pains due to:
■ backache ■ toothache
■ menstrual cramps ■ headache
■ minor pain of arthritis
■ temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ facial swelling ■ shock ■ asthma (wheezing) ■ rash
■ skin reddening ■ blisters ■ hives
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ take more or for a longer time than directed ■ are age 60 or older
■ take a blood thinner (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have had stomach ulcers or bleeding problems
■ have 3 or more alcoholic drinks every day while using this product
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Directions
■ do not take more than directed
■ the smallest effective dose should be used
■ adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist
■ if pain or fever does not respond to 1 caplet, 2 caplets may be used
■ do not exceed 6 caplets in 24 hours, unless directed by a doctor
■ children under 12 years: ask a doctor

Other information
■ store between 20°-25° (68°-77°F)
■ avoid excessive heat 40°C (104°F)
■ use by expiration date on package

Inactive ingredients carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?
Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday
STOP PEELING

0 11822 42091 4

PEEL HERE FOR MORE DRUG FACTS

†This product is not manufactured or distributed by PF Consumer Healthcare 1 LLC, owner of the registered trademark Advil® Caplets. 50844 REV1221A29214

DISTRIBUTED BY:
RITE AID, 200 NEWBERRY COMMONS
ETTERS, PA 17319 www.riteaid.com

SATISFACTION GUARANTEE:
If you're not satisfied, we'll happily refund your money.

Drug Facts (continued)

Do not use
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
■ right before or after heart surgery

Ask a doctor before use if
■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic
■ you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are
■ under a doctor's care for any serious condition
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
■ taking any other drug

When using this product
■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood
■ have bloody or black stools
■ have stomach pain that does not get better
■ you have symptoms of heart problems or stroke:
■ chest pain ■ slurred speech
■ leg swelling ■ trouble breathing
■ weakness in one part or side of body
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present in the painful area
■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
■ do not take more than directed
■ the smallest effective dose should be used
■ adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist
■ if pain or fever does not respond to 1 caplet, 2 caplets may be used
■ do not exceed 6 caplets in 24 hours, unless directed by a doctor
■ children under 12 years: ask a doctor

Other information
■ store between 20°-25° (68°-77°F)
■ avoid excessive heat 40°C (104°F)
■ use by expiration date on package

Inactive ingredients carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?
Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday
STOP PEELING

0 11822 42091 4

PEEL HERE FOR MORE DRUG FACTS

†This product is not manufactured or distributed by PF Consumer Healthcare 1 LLC, owner of the registered trademark Advil® Caplets. 50844 REV1221A29214

DISTRIBUTED BY:
RITE AID, 200 NEWBERRY COMMONS
ETTERS, PA 17319 www.riteaid.com

SATISFACTION GUARANTEE:
If you're not satisfied, we'll happily refund your money.

Rite Aid 44-292

PAIN RELIEF IBUPROFEN

ibuprofen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-0292
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	brown	Score	no score
Shape	OVAL	Size	14mm
Flavor		Imprint Code	44;292
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0292-5	1 in 1 CARTON	05/24/1988	
1		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:11822-0292-2	1 in 1 CARTON	05/24/1988	
2		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:11822-0292-4	500 in 1 BOTTLE; Type 0: Not a Combination Product	05/24/1988	
	NDC:11822-			

4	NDC:11822-0292-8	1 in 1 BLISTER PACK	05/24/1988	03/12/2023
4		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		
5	NDC:11822-0292-6	1 in 1 CARTON	05/24/1988	10/08/2021
5		16 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
6	NDC:11822-0292-7	1 in 1 CARTON	05/24/1988	10/08/2021
6		125 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	05/24/1988	

Labeler - Rite Aid Corporation (014578892)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(11822-0292)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(11822-0292)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(11822-0292)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(11822-0292)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(11822-0292)