

BB TINTED TREATMENT 12-HOUR PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN FAIR- titanium dioxide and zinc oxide liquid
BB TINTED TREATMENT 12-HOUR PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN LIGHT- titanium dioxide and zinc oxide liquid
BB TINTED TREATMENT 12-HOUR PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN MEDIUM- titanium dioxide and zinc oxide liquid
BB TINTED TREATMENT 12-HOUR PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN TAN- titanium dioxide and zinc oxide liquid
BB TINTED TREATMENT 12-HOUR PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN DEEP- titanium dioxide and zinc oxide liquid
BB TINTED TREATMENT 12-HOUR PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN RICH- titanium dioxide and zinc oxide liquid
Tarte, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

BB Tinted Treatment 12-Hour Primer Broad Spectrum SPF 30 Sunscreen

Drug Facts

<i>Active Ingredients</i>	<i>Purpose</i>
Titanium Dioxide 4.55%	Sunscreen
Zinc Oxide 3.5%	Sunscreen

Uses

- Helps prevent sunburn.
- If used as directed with other sun protection measures (see ***Directions***), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Stop use and ask doctor if rash occurs.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Shake before use.
- Apply liberally 15 minutes before sun exposure.
- Use a water resistant sunscreen if swimming or sweating.

- Reapply at least every 2 hours.
- Children under 6 months: Ask a doctor.
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
 - Limit time in the sun, especially from 10 a.m.-2 p.m.
 - Wear long-sleeved shirts, pants, hats and sunglasses

Other information

- Protect the product in this container from excessive heat and direct sunlight.
- You may report a serious adverse reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainsboro, NJ 08536-0222.

Inactive Ingredients

Cyclopentasiloxane, isododecane, mica, polysilicone-11, polymethylsilsesquioxane, hexyl laurate, PEG-10 dimethicone, polyglyceryl-4 isostearate, stearic acid, cetyl PEG/PPG-10/1 dimethicone, alumina, triethoxycaprylylsilane, dipalmitoyl hydroxyproline, diamond powder, iron oxides.

Dist. by tarte, Inc.
New York, NY 10018.

PRINCIPAL DISPLAY PANEL - 10 mL Tube Carton - fair

tarte
high-performance naturals™
BB tinted treatment primer
Broad Spectrum
SPF 30 Sunscreen
net 10 mL / 0.33 Fl. Oz.

Drug Facts

Active Ingredients Purpose

Titanium Dioxide 4.55%.....Sunscreen
Zinc Oxide 3.5%.....Sunscreen

Uses

- Helps prevent sunburn.
- If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Stop use and ask doctor if rash occurs.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Shake before use.
- Apply liberally 15 minutes before sun exposure.
- Use a water resistant sunscreen if swimming or sweating.
- Reapply at least every 2 hours.
- Children under 6 months: Ask a doctor.

Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:

- Limit time in the sun, especially from 10 a.m.-2 p.m.
- Wear long-sleeved shirts, pants, hats and sunglasses

Drug Facts (continued)

Other information

- Protect the product in this container from excessive heat and direct sunlight.

- You may report a serious adverse reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainboro, NJ 08536-0222.

Inactive Ingredients

Cyclopentasiloxane, isododecane, mica, polyasilicone-11, polymethylsilsequioxane, hexyl laurate, PEG-10 dimethicone, polyglyceryl-4 isostearate, stearic acid, cetyl PEG/PPG-10/1 dimethicone, alumina, triethoxycaprylylsilane, dipalmitoyl hydroxyproline, diamond powder, iron oxides.

Made in USA. Dist. by tarte, Inc. New York, NY 10018. All rights reserved. BIORIUS: 7170 Manage, Belgium / London WC2H 9JQ, UK.

#claypowered

@tartecosmetics

www.tarte.com

fair
12-hour primer

tarte
high-performance naturals™

• triple-B complex: blend of natural diamond powder, plants & lipoamino acid

• pore minimizing, smoothing & brightening

• 12-hour primer & dermatologist tested

• wear alone for a sheer tint or under foundation for enhanced coverage

BB tinted treatment primer

Broad Spectrum
SPF 30 Sunscreen

net 10 mL / 0.33 FL. Oz.



crueltyfree
andvegan



FG08100-01



PRINCIPAL DISPLAY PANEL - 10 mL Tube Carton - light

tarte

high-performance naturals™

BB tinted treatment primer

Broad Spectrum

SPF 30 Sunscreen

net 10 mL / 0.33 Fl. Oz.

Drug Facts

Active Ingredients Purpose

Titanium Dioxide 4.55%.....Sunscreen
Zinc Oxide 3.5%.....Sunscreen

Uses

- Helps prevent sunburn.
- If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Stop use and ask doctor if rash occurs.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Shake before use.
- Apply liberally 15 minutes before sun exposure.
- Use a water resistant sunscreen if swimming or sweating.
- Reapply at least every 2 hours.
- Children under 6 months: Ask a doctor.
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
 - Limit time in the sun, especially from 10 a.m.-2 p.m.
 - Wear long-sleeved shirts, pants, hats and sunglasses

Drug Facts (continued)

Other information

- Protect the product in this container from excessive heat and direct sunlight.
- You may report a serious adverse reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainboro, NJ 08536-0222.

Inactive Ingredients

Cyclopentasiloxane, isododecane, polydimethylsiloxane, mica, polymethylsiloxane, hexyl laurate, PEG-10 dimethicone, polyglyceryl-4 isostearate, stearic acid, cetyl PEG/PPG-10/1 dimethicone, alumina, triethoxycaprylylsilane, dipalmitoyl hydroxyproline, diamond powder, iron oxides.

Made in USA. Dist. by tarte, Inc. New York, NY 10018. All rights reserved. BIORIUS: 7170 Manage, Belgium / London WC2H 9JQ, UK.

#claypowered

@tartecosmetics

www.tarte.com

light
12-hour primer

tarte
high-performance naturals™

• triple-B complex: blend of natural diamond powder, plants & lipoamino acid

• pore minimizing, smoothing & brightening

• 12-hour primer & dermatologist tested

• wear alone for a sheer tint or under foundation for enhanced coverage

BB tinted treatment primer

Broad Spectrum
SPF 30 Sunscreen

net 10 mL / 0.33 FL. Oz.



FG08101-01



PRINCIPAL DISPLAY PANEL - 10 mL Tube Carton - medium

tarte

high-performance naturals™

BB tinted treatment primer

Broad Spectrum

SPF 30 Sunscreen

net 10 mL / 0.33 Fl. Oz.

medium
12-hour primer

tarte
high-performance naturals

- triple-B complex: blend of natural diamond powder, plants & lipoamino acid

- pore minimizing, smoothing & brightening

- 12-hour primer & dermatologist tested

- wear alone for a sheer tint or under foundation for enhanced coverage

Drug Facts

Active Ingredients Purpose

Titanium Dioxide 4.55%.....Sunscreen
Zinc Oxide 3.5%.....Sunscreen

Uses

- Helps prevent sunburn.
- If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Stop use and ask doctor if rash occurs.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Shake before use.
- Apply liberally 15 minutes before sun exposure.
- Use a water resistant sunscreen if swimming or sweating.
- Reapply at least every 2 hours.
- Children under 6 months: Ask a doctor.
- Sun Protection Measures: Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
 - Limit time in the sun, especially from 10 a.m.-2 p.m.
 - Wear long-sleeved shirts, pants, hats and sunglasses

Drug Facts (continued)

Other information

- Protect the product in this container from excessive heat and direct sunlight.
- You may report a serious adverse reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainboro, NJ 08538-0222.

Inactive Ingredients

Cyclopentasiloxane, isododecane, polysilicone-11, mica, polymethylsilsequioxane, hexyl laurate, PEG-10 dimethicone, polyglyceryl-4 isostearate, stearic acid, cetyl PEG/PPG-10/1 dimethicone, alumina, triethoxycaprylylsilane, dipalmitoyl hydroxyproline, diamond powder, iron oxides.

Made in USA. Dist. by tarte, Inc. New York, NY 10018. All rights reserved. BIORIUS: 7170 Manage, Belgium / London WC2H 9JQ, UK.

#claypowered
@tartecosmetics
www.tarte.com

BB tinted treatment primer
Broad Spectrum
SPF 30 Sunscreen
net 10 mL / 0.33 FL Oz.



FG08102-01



PRINCIPAL DISPLAY PANEL - 10 mL Tube Carton - tan

tarte

high-performance naturals™

BB tinted treatment primer

Broad Spectrum

SPF 30 Sunscreen

net 10 mL / 0.33 Fl. Oz.

Drug Facts

Active Ingredients Purpose

Titanium Dioxide 4.55%.....Sunscreen
Zinc Oxide 3.5%.....Sunscreen

Uses

- Helps prevent sunburn.
- If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Stop use and ask doctor if rash occurs.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Shake before use.
- Apply liberally 15 minutes before sun exposure.
- Use a water resistant sunscreen if swimming or sweating.
- Reapply at least every 2 hours.
- Children under 6 months: Ask a doctor.
- Sun Protection Measures: Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
 - Limit time in the sun, especially from 10 a.m.-2 p.m.
 - Wear long-sleeved shirts, pants, hats and sunglasses

Drug Facts (continued)

Other information

- Protect the product in this container from excessive heat and direct sunlight.
- You may report a serious adverse reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainboro, NJ 08536-0222.

Inactive Ingredients

Cyclopentasiloxane, isododecane, polysilicone-11, mica, polymethylsilsequioxane, hexyl laurate, PEG-10 dimethicone, polyglyceryl-4 isostearate, stearic acid, cetyl PEG/PPG-10/1 dimethicone, alumina, triethoxycaprylylsilane, dipalmitoyl hydroxyproline, diamond powder, iron oxides.

Made in USA. Dist. by tarte, Inc. New York, NY 10018. All rights reserved. BIORIUS: 7170 Manage, Belgium / London WC2H 9JQ, UK.

#claypowered
@tartecosmetics
www.tarte.com

tan
12-hour primer

tarte
high-performance naturals

- triple-B complex: blend of natural diamond powder, plants & lipoamino acid

- pore minimizing, smoothing & brightening

- 12-hour primer & dermatologist tested

- wear alone for a sheer tint or under foundation for enhanced coverage

BB tinted treatment primer
Broad Spectrum
SPF 30 Sunscreen
net 10 mL / 0.33 FL Oz.



FG08103-01



PRINCIPAL DISPLAY PANEL - 10 mL Tube Carton - deep

tarte

high-performance naturals™

BB tinted treatment primer

Broad Spectrum

SPF 30 Sunscreen

net 10 mL / 0.33 Fl. Oz.

deep
12-hour primer

tarte
high-performance naturals™

- triple-B complex: blend of natural diamond powder, plants & lipoamino acid

- pore minimizing, smoothing & brightening

- 12-hour primer & dermatologist tested

- wear alone for a sheer tint or under foundation for enhanced coverage

Drug Facts

Active Ingredients Purpose

Titanium Dioxide 4.55%.....Sunscreen
Zinc Oxide 3.5%.....Sunscreen

Uses

- Helps prevent sunburn.
- If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Stop use and ask doctor if rash occurs.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Shake before use.
- Apply liberally 15 minutes before sun exposure.
- Use a water resistant sunscreen if swimming or sweating.
- Reapply at least every 2 hours.
- Children under 6 months: Ask a doctor.

Sun Protection Measures.
Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:

- Limit time in the sun, especially from 10 a.m.-2 p.m.
- Wear long-sleeved shirts, pants, hats and sunglasses

Drug Facts (continued)

Other information

- Protect the product in this container from excessive heat and direct sunlight.

- You may report a serious adverse reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainboro, NJ 08536-0222.

Inactive Ingredients

Cyclopentasiloxane, isododecane, polyisilicone-11, polymethylsiloxane, mica, hexyl laurate, PEG-10 dimethicone, polyglyceryl-4 isostearate, stearic acid, cetyl PEG/PPG-10/1 dimethicone, alumina, triethoxycaprylylsilane, dipalmitoyl hydroxyproline, diamond powder, iron oxides.

Made in USA. Dist. by tarte, Inc. New York, NY 10018. All rights reserved. BIORIUS: 7170 Manage, Belgium / London WC2H 9JQ, UK.

#claypowered

@tartecosmetics

www.tarte.com

BB tinted treatment primer

Broad Spectrum
SPF 30 Sunscreen

net 10 mL / 0.33 FL Oz.



FG08104-01



PRINCIPAL DISPLAY PANEL - 10 mL Tube Carton - rich

tarte

high-performance naturals™

BB tinted treatment primer

Broad Spectrum

SPF 30 Sunscreen

net 10 mL / 0.33 Fl. Oz.

Drug Facts

Active Ingredients Purpose

Titanium Dioxide 4.55%.....Sunscreen
Zinc Oxide 3.5%.....Sunscreen

Uses

- Helps prevent sunburn.
- If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Stop use and ask doctor if rash occurs.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Shake before use.
- Apply liberally 15 minutes before sun exposure.
- Use a water resistant sunscreen if swimming or sweating.
- Reapply at least every 2 hours.
- Children under 6 months: Ask a doctor.

Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:

- Limit time in the sun, especially from 10 a.m.-2 p.m.
- Wear long-sleeved shirts, pants, hats and sunglasses

Drug Facts (continued)

Other information

- Protect the product in this container from excessive heat and direct sunlight.
- You may report a serious adverse reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainboro, NJ 08536-0222.

Inactive Ingredients

Cyclopentasiloxane, isododecane, polyisilicone-11, polymethylsiloxane, hexyl laurate, PEG-10 dimethicone, polyglyceryl-4 isostearate, mica, stearic acid, cetyl PEG/PPG-10/1 dimethicone, alumina, triethoxycaprylylsilane, dipalmitoyl hydroxyproline, diamond powder, iron oxides.

Made in USA. Dist. by tarte, Inc. New York, NY 10018. All rights reserved. BIORIUS: 7170 Manage, Belgium / London WC2H 9JQ, UK.

#claypowered
@tartecosmetics
www.tarte.com

rich
12-hour primer

tarte
high-performance naturals

• triple-B complex: blend of natural diamond powder, plants & lipoamino acid

• pore minimizing, smoothing & brightening

• 12-hour primer & dermatologist tested

• wear alone for a sheer tint or under foundation for enhanced coverage

BB tinted treatment primer
Broad Spectrum
SPF 30 Sunscreen
net 10 mL / 0.33 FL Oz.



FG08105-01

BB TINTED TREATMENT 12-HOUR PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN FAIR

titanium dioxide and zinc oxide liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51060-324
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	45.5 mg in 1 mL
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	35 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
ISODODECANE (UNII: A8289P68Y2)	
MICA (UNII: V8A1AW0880)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
POLYMETHYLSILSESQUIOXANE (11 MICRONS) (UNII: Z570VEV8XK)	
HEXYL LAURATE (UNII: 4CG9F9W01Q)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)	
BENZIMIDAZOLE (UNII: E24GX49LD8)	
DIAMOND (UNII: 6GRV67N0U2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CETYL PEG/PPG-10/1 DIMETHICONE (HLB 2) (UNII: V2W71V8T0X)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
DIPALMITOYL HYDROXYPROLINE (UNII: E6AHA53N1H)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51060-324-01	1 in 1 CARTON	09/01/2012	
1		10 mL in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part352	09/01/2012	

BB TINTED TREATMENT 12-HOUR PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN LIGHT

titanium dioxide and zinc oxide liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51060-325
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	45.5 mg in 1 mL
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	35 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
MICA (UNII: V8A1AW0880)	
ISODODECANE (UNII: A8289P68Y2)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
POLYMETHYLSILSESQUIOXANE (11 MICRONS) (UNII: Z570VEV8XK)	
HEXYL LAURATE (UNII: 4CG9F9W01Q)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)	
BENZIMIDAZOLE (UNII: E24GX49LD8)	
DIAMOND (UNII: 6GRV67N0U2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CETYL PEG/PPG-10/1 DIMETHICONE (HLB 2) (UNII: V2W71V8TOX)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
DIPALMITOYL HYDROXYPROLINE (UNII: E6AHA53N1H)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:51060-325-01	1 in 1 CARTON	09/01/2012	
1		10 mL in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part352	09/01/2012	

BB TINTED TREATMENT 12-HOUR PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN MEDIUM

titanium dioxide and zinc oxide liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51060-326
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	45.5 mg in 1 mL
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	35 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
ISODODECANE (UNII: A8289P68Y2)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
MICA (UNII: V8A1AW0880)	
POLYMETHYLSILSESQUIOXANE (11 MICRONS) (UNII: Z570VEV8XK)	
HEXYL LAURATE (UNII: 4CG9F9W01Q)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CETYL PEG/PPG-10/1 DIMETHICONE (HLB 2) (UNII: V2W71V8T0X)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
DIPALMITOYL HYDROXYPROLINE (UNII: E6AHA53N1H)	
DIAMOND (UNII: 6GRV67N0U2)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51060-326-01	1 in 1 CARTON	09/01/2012	
1		10 mL in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part352	09/01/2012	

BB TINTED TREATMENT 12-HOUR PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN TAN

titanium dioxide and zinc oxide liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51060-327
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	45.5 mg in 1 mL
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	35 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
ISODODECANE (UNII: A8289P68Y2)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
POLYMETHYLSILSESQUIOXANE (11 MICRONS) (UNII: Z570VEV8XK)	
HEXYL LAURATE (UNII: 4CG9F9W01Q)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V15VM0)	
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)	
CYCLOMETHICONE 6 (UNII: XHK3U310BA)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CETYL PEG/PPG-10/1 DIMETHICONE (HLB 2) (UNII: V2W71V8T0X)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
DIPALMITOYL HYDROXYPROLINE (UNII: E6AHA53N1H)	
DIAMOND (UNII: 6GRV67N0U2)	

MICA (UNII: V8A1AW0880)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51060-327-01	1 in 1 CARTON	09/01/2012	
1		10 mL in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part352	09/01/2012	

BB TINTED TREATMENT 12-HOUR PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN DEEP

titanium dioxide and zinc oxide liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51060-328
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	45.5 mg in 1 mL
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	35 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
ISODODECANE (UNII: A8289P68Y2)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
POLYMETHYLSILSESQUIOXANE (11 MICRONS) (UNII: Z570VEV8XK)	
HEXYL LAURATE (UNII: 4CG9F9W01Q)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)	
DIAMOND (UNII: 6GRV67N0U2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

CETYL PEG/PPG-10/1 DIMETHICONE (HLB 2) (UNII: V2W71V8T0X)
ALUMINUM OXIDE (UNII: LMI26O6933)
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)
DIPALMITOYL HYDROXYPROLINE (UNII: E6AHA53N1H)
MICA (UNII: V8A1AW0880)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51060-328-01	1 in 1 CARTON	09/01/2012	
1		10 mL in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part352	09/01/2012	

BB TINTED TREATMENT 12-HOUR PRIMER BROAD SPECTRUM SPF 30 SUNSCREEN RICH
titanium dioxide and zinc oxide liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51060-329
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	45.5 mg in 1 mL
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	35 mg in 1 mL

Inactive Ingredients	
Ingredient Name	Strength
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
ISODODECANE (UNII: A8289P68Y2)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
POLYMETHYLSILSESQUIOXANE (11 MICRONS) (UNII: Z570VEV8XK)	
HEXYL LAURATE (UNII: 4CG9F9W01Q)	

PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)
CYCLOMETHICONE 6 (UNII: XHK3U310BA)
STEARIC ACID (UNII: 4ELV7Z65AP)
CETYL PEG/PPG-10/1 DIMETHICONE (HLB 2) (UNII: V2W71V8T0X)
ALUMINUM OXIDE (UNII: LMI26O6933)
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)
DIPALMITOYL HYDROXYPROLINE (UNII: E6AHA53N1H)
DIAMOND (UNII: 6GRV67N0U2)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
MICA (UNII: V8A1AW0880)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERROSO FERRIC OXIDE (UNII: XM0M87F357)

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51060-329-01	1 in 1 CARTON	09/01/2012	
1		10 mL in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part352	09/01/2012	

Labeler - Tarte, Inc. (027905186)