

ASPIRIN LOW DOSE- aspirin tablet, delayed release
COSTCO WHOLESALE CORPORATION

Kirkland 44-600A

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains or as recommended by your doctor.
Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- take more or for a longer time than directed
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have asthma
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - feel faint
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

Kirkland

Signature

COMPARE TO BAYER®

LOW DOSE ASPIRIN

active ingredient†

NDC 63981-563-96

ITM. / ART. 310715

LOW DOSE

ASPIRIN

81 mg

Pain Reliever (NSAID)

*Aspirin Regimen***

Enteric Coated

2 x 365

Tablets

Actual Size

**Talk to your doctor or other healthcare provider before using this product for your heart.

†This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Low Dose Aspirin. 50844 REV0122E60051

Manufactured by: LNK INTERNATIONAL, INC.

60 Arkay Drive, Hauppauge, NY 11788 USA

For: Costco Wholesale Corporation

P.O. Box 34535, Seattle, WA 98124-1535 USA

1-800-774-2678 • www.costco.com

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

KIRKLAND
Signature

COMPARE TO BAYER®
LOW DOSE ASPIRIN
active ingredient†

NDC 63981-563-96
ITM. / ART. 310715

LOW DOSE
ASPIRIN
81 mg

Pain Reliever (NSAID)

- Aspirin Regimen**
- Enteric Coated

2 x 365
Tablets



Actual Size

**Talk to your doctor or other healthcare provider before using this product for your heart.

TAMPER EVIDENT: DO NOT USE IF
IMPRINTED SAFETY SEAL UNDER
CAP IS BROKEN OR MISSING



ITM. / ART. 310715
1-800-774-2678 • www.costco.com
P.O. Box 34535, Seattle, WA 98124-1535 USA
For Costco Wholesale Corporation
60 Arkay Drive, Hauppauge, NY 11788 USA

Manufactured by: LNK INTERNATIONAL, INC. REV0122E60051
 This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Low Dose Aspirin, 50844

<p>Questions or comments? 1-800-426-9391</p>	<p>■ you are taking a diuretic disease, liver cirrhosis, or kidney disease</p> <p>■ you have high blood pressure, heart such as heartburn ■ you have asthma</p> <p>■ you have a history of stomach problems, stomach bleeding warning applies to you</p> <p>Ask a doctor before use if</p> <p>■ if you have ever had an allergic reaction to this product or any of its ingredients</p> <p>■ if you have ever had an allergic reaction to pain reliever/ever reducer</p> <p>■ if you are allergic to aspirin or any other</p>
<p>Inactive ingredients com starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, stibic wax, silica, simethicone, sodium bicarbonate, triacetin, triethyl citrate</p>	<p>Do not use</p> <p>■ if you are allergic to aspirin or any other steroid drug ■ are age 60 or older</p> <p>■ take a blood thinning (anticoagulant) or while using this product</p> <p>■ have 3 or more alcoholic drinks every day</p> <p>■ have had stomach ulcers or bleeding problems</p> <p>■ take more or for a longer time than directed</p> <p>■ naproxen, naproxen, or others] nonprescription NSAIDs [aspirin, take other drugs containing prescription or you</p>
<p>Other information</p> <p>■ use by expiration date on package</p> <p>■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)</p>	<p>stomach bleeding. The chance is higher if you contains an NSAID, which may cause severe stomach bleeding. This product</p> <p>■ shock ■ asthma (wheezing) ■ hives ■ facial swelling</p> <p>allergic reaction, which may include:</p> <p>Allergy alert: Aspirin may cause a severe rare but serious illness.</p> <p>could be an early sign of Reye's syndrome, a consult a doctor because these symptoms occur, in behavior with nausea and vomiting occur, product. When using this product, if changes or flu-like symptoms should not use this who have or are recovering from chicken pox</p>
<p>Directions</p> <p>■ do not take more than directed</p> <p>■ drink a full glass of water with each dose</p> <p>■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor</p> <p>■ children under 12 years: do not use unless directed by a doctor</p>	<p>Warnings</p> <p>Reye's syndrome: Children and teenagers or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ shock ■ asthma (wheezing) ■ hives ■ facial swelling</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]</p>
<p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>■ do not take more than directed</p> <p>■ drink a full glass of water with each dose</p> <p>■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor</p> <p>■ children under 12 years: do not use unless directed by a doctor</p>	<p>Uses for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.</p> <p>Warnings</p> <p>Reye's syndrome: Children and teenagers or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ shock ■ asthma (wheezing) ■ hives ■ facial swelling</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]</p>
<p>Drug Facts (continued)</p> <p>Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis</p> <p>Stop use and ask a doctor if</p> <p>■ an allergic reaction occurs. Seek medical help right away.</p> <p>■ you experience any of the following signs of stomach bleeding: ■ vomit blood ■ have bloody or black stools ■ feel faint ■ have stomach pain that does not get better ■ ringing in the ears or a loss of hearing occurs</p> <p>■ pain gets worse or lasts more than 10 days</p> <p>■ fever gets worse or lasts more than 3 days</p> <p>■ redness or swelling is present</p> <p>■ new symptoms occur</p> <p>These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p>	<p>Active ingredient Aspirin 81 mg (NSAID) Pain reliever, nonsteroidal anti-inflammatory drug</p>

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

ASPIRIN
 LOW DOSE
 81 mg
 Pain Reliever (NSAID)

Kirkland 44-600A

ASPIRIN LOW DOSE

aspirin tablet, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63981-563
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
METHACRYLIC ACID (UNII: 1CS02G8656)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SHELLAC (UNII: 46N107B71O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	L
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63981-563-96	2 in 1 PACKAGE	05/01/2011	
1		365 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:63981-563-51	2 in 1 CARTON	05/01/2011	05/28/2021
2		365 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:63981-563-62	1 in 1 PACKAGE	05/01/2011	11/08/2021
3		15 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/01/2011	

Labeler - COSTCO WHOLESALE CORPORATION (103391843)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(63981-563)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(63981-563)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(63981-563) , pack(63981-563)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(63981-563)

Revised: 4/2025

COSTCO WHOLESALE CORPORATION