

**COLD AND FLU RELIEF NIGHTTIME- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled
Safeway, Inc.**

Safeway Nighttime Cold & Flu

Active Ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine Succinate 6.25 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Purpose

Pain reliever / fever reducer

Cough suppressant

Antihistamine

Uses

- temporarily relieves common cold and flu symptoms:
 - sore throat
 - headache
 - minor aches and pains
 - fever
 - runny nose and sneezing
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough that such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- **when using other Daytime or Nighttime products, carefully read each label to insure correct dosing**

Other Information

- store between 15-30°C (59-86°F)
- avoid excessive heat

Inactive Ingredients

D&C yellow #10, edible white ink, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or Comments?

Call **1-888-723-3929 Monday-Friday 7AM-6PM PST**

Principal Display Panel

COMPARE TO Vick® NyQuil® LiquiCaps® active ingredients†

Nighttime Cold & Flu Relief

Acetaminophen 325 mg - Pain Reliever / Fever Reducer

Dextromethorphan HBr 15 mg - Cough Suppressant

Doxylamine Succinate 6.25 mg - Antihistamine

SOFTGELS**

(**Liquid-filled capsules)

†This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, NyQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS

TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY BETTER LIVING BRANDS LLC

P.O. BOX 99, PLEASANTON, CA 94566-0009

www.betterlivingbrandsLLC.com

Package Label



Nighttime Cold & Flu Relief

**BONUS
PACK
50%
MORE**



Nighttime Cold & Flu Relief

Acetaminophen 325 mg - Pain Reliever/Fever Reducer
Dextromethorphan HBr 15 mg - Cough Suppressant
Doxylamine succinate 6.25 mg - Antihistamine

36 SOFTGELS**
 (**Liquid-filled capsules)



Actual Size

**24 36
SOFTGELS**

COMPARE TO
Vicks® NyQuil® LiquiCaps®
 active ingredients†

NDC 21130-071-36

**BONUS
PACK
50%
MORE**

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.
KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.



Lot / Exp.:
Product of:

Drug Facts (continued) Active ingredients (in each softgel) Acetaminophen 325 mg Dextromethorphan HBr 15 mg Doxylamine succinate 6.25 mg Purposes Pain reliever/fever reducer Cough suppressant Antihistamine Uses Temporarily relieves common cold and flu symptoms ■ sore throat ■ headache ■ minor aches and pains ■ fever ■ cough due to minor throat and bronchial irritation Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. Safe throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Directions ■ do not take more than 4 doses in 24 hours ■ do not take more than directed (see Overdose warning) Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Keep out of reach of children. Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Stop use and ask a doctor if: ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. When using this product ■ do not exceed recommended dosage ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin Ask a doctor or pharmacist before use if you are: ■ a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema ■ cough that occurs with too much phlegm (mucus) ■ trouble urinating due to an enlarged prostate gland Ask a doctor before use if you have: ■ liver disease ■ glaucoma	
---	--

3 21130 77220 0



RD 17124

S2801

OUR PROMISE
QUALITY & SATISFACTION
100% GUARANTEED
ON YOUR MONEY BACK.

PROCTER & GAMBLE COMPANY
Company, Vicks®, NyQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.

Call 1-888-723-3929 Monday-Friday 7AM-6PM PST

Other Information
■ store between 15-30°C (59-86°F) ■ avoid excessive heat
■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing
■ children under 12 years: do not use
■ swallow whole; do not crush, chew, or dissolve
■ adults and children 12 years and over: take 2 softgels with water every 6 hours

Drug Facts (continued)
■ Inactive Ingredients D&C yellow #10, edible white ink, FD&C blue #1, sorbitan, glycerin, polyethylene glycol, povidone, purified water, gelatin, glycerin, sorbitan, sorbitan, sorbitan
■ Questions or comments? Call 1-888-723-3929 Monday-Friday 7AM-6PM PST

1-888-723-3929
PLEASANTON, CA 94566-0009
P.O. BOX 99
BETTER LIVING BRANDS LLC
DISTRIBUTED BY
www.betterlivingbrandsllc.com

03/14/14

SIGNATURE CARE Nighttime Cold and Flu Relief

COLD AND FLU RELIEF NIGHTTIME

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:21130-071
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	green	Score	no score
Shape	OVAL	Size	21mm
Flavor		Imprint Code	P30;94A;SCU1;215;P120

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-071-24	24 in 1 CARTON	02/29/2016	02/28/2025
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:21130-071-36	36 in 1 CARTON	02/29/2016	02/28/2025
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	02/29/2016	02/28/2025

Labeler - Safeway, Inc. (009137209)

Revised: 5/2024

Safeway, Inc.