

EZETIMIBE AND SIMVASTATIN - ezetimibe and simvastatin tablet

Ascend Laboratories, LLC

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use EZETIMIBE AND SIMVASTATIN TABLETS safely and effectively. See full prescribing information for EZETIMIBE AND SIMVASTATIN TABLETS.

EZETIMIBE and SIMVASTATIN tablets, for oral use

Initial U.S. Approval: 2004

RECENT MAJOR CHANGES

Indications and Usage (1)	2/2024
Dosage and Administration (2.1, 2.2, 2.3, 2.4, 2.5)	2/2024
Contraindications, Pregnancy and Lactation (4) Removed	2/2024
Warnings and Precautions (5.1, 5.2, 5.3)	2/2024

INDICATIONS AND USAGE

Ezetimibe and simvastatin tablets is a combination of ezetimibe, a dietary cholesterol absorption inhibitor, and simvastatin, an HMG-CoA reductase inhibitor (statin) indicated: (1)

- As an adjunct to diet to reduce elevated low density lipoprotein cholesterol (LDL-C):
 - In adults with primary hyperlipidemia.
 - In adults and pediatric patients aged 10 years and older with heterozygous familial hypercholesterolemia (HeFH).
- As an adjunct to other LDL-C lowering therapies to reduce LDL-C in adults with homozygous familial hypercholesterolemia (HoFH).

Simvastatin

Simvastatin, when used as a component of ezetimibe and simvastatin, is indicated to reduce the risk of total mortality by reducing risk of coronary heart disease death, non-fatal myocardial infarction and stroke, and the need for coronary and non-coronary revascularization procedures in adults with established coronary heart disease, cerebrovascular disease, peripheral vascular disease, and/or diabetes, who are at high risk of coronary heart disease events.

DOSAGE AND ADMINISTRATION

- *Important Dosage and Administration Information: (2.1)*
 - Take ezetimibe and simvastatin orally once daily in the evening with or without food.
 - Maximum recommended dosage is ezetimibe and simvastatin 10/40 mg once daily. Ezetimibe and simvastatin 10/80 mg daily dosage is restricted to patients who have been taking ezetimibe and simvastatin 10/80 mg daily chronically (e.g., for 12 months or more) without evidence of muscle toxicity.
 - For patients that require a high-intensity statin or are unable to achieve their LDL-C goal receiving ezetimibe and simvastatin 10/40 mg daily, prescribe alternative LDL-C-lowering treatment.
 - If a dose is missed, take the missed dose as soon as possible. Do not double the next dose.
 - Assess LDL-C when clinically appropriate, as early as 2 weeks after initiating ezetimibe and simvastatin, and adjust the dosage if necessary.
- *Adults:* Recommended dosage range of 10/10 mg to 10/40 mg once daily. (2.2)
- See full prescribing information for ezetimibe and simvastatin dosage modifications due to drug interactions. (2.3)
- *Patients with Renal Impairment:* Doses exceeding 10/20 mg should be used with caution and close monitoring in patients with moderate to severe renal impairment. (2.4)

DOSAGE FORMS AND STRENGTHS

- Tablets (ezetimibe, USP mg/simvastatin, USP mg): 10/10, 10/20, 10/40, 10/80 (3)

CONTRAINDICATIONS

- Concomitant use of strong CYP3A4 inhibitors (select azole anti-fungals, macrolide antibiotics, anti-viral

- medications and nefazodone). (4)
- Concomitant use of cyclosporine, danazol or gemfibrozil. (4)
 - Acute liver failure or decompensated cirrhosis. (4)
 - Hypersensitivity to simvastatin, ezetimibe or any excipient of ezetimibe and simvastatin tablets. (4)

-----**WARNINGS AND PRECAUTIONS**-----

- *Myopathy and Rhabdomyolysis*: Risk factors include age 65 years or greater, uncontrolled hypothyroidism, renal impairment, concomitant use with certain other drugs, and higher ezetimibe and simvastatin dosage. Chinese patients may be at higher risk for myopathy. Discontinue ezetimibe and simvastatin if markedly elevated CK levels occur or myopathy is diagnosed or suspected. Temporarily discontinue ezetimibe and simvastatin in patients experiencing an acute or serious condition at high risk of developing renal failure secondary to rhabdomyolysis. Inform patients of the risk of myopathy and rhabdomyolysis when starting or increasing ezetimibe and simvastatin dosage. Instruct patients to promptly report unexplained muscle pain, tenderness, or weakness, particularly if accompanied by malaise or fever. (5.1)
- *Immune-Mediated Necrotizing Myopathy (IMNM)*: Rare reports of IMNM, an autoimmune myopathy, have been reported. Discontinue ezetimibe and simvastatin if IMNM is suspected. (5.2)
- *Hepatic Dysfunction*: Increases in serum transaminases have occurred, some persistent. Rare reports of fatal and non-fatal hepatic failure have occurred. Consider testing liver enzyme before initiating therapy and as clinically indicated thereafter. If serious hepatic injury with clinical symptoms and/or hyperbilirubinemia or jaundice occurs, promptly discontinue ezetimibe and simvastatin tablets. (5.3)

-----**ADVERSE REACTIONS**-----

- Common (incidence $\geq 2\%$ and greater than placebo) adverse reactions in clinical trials: headache, increased ALT, myalgia, upper respiratory tract infection, and diarrhea. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Ascend Laboratories, LLC at 1-877-272-7901 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

-----**DRUG INTERACTIONS**-----

- See full prescribing information for details regarding concomitant use of ezetimibe and simvastatin with other drugs or grapefruit juice that increase the risk of myopathy and rhabdomyolysis. (2.3, 7.1)
- *Cholestyramine*: Combination decreases exposure of ezetimibe. (2.3, 7.2)
- *Coumarin Anticoagulants*: Obtain INR before ezetimibe and simvastatin initiation and monitor INR during ezetimibe and simvastatin dosage initiation or adjustment. (7.3)
- *Digoxin*: During ezetimibe and simvastatin initiation, monitor digoxin levels. (7.3)
- *Fenofibrates*: Combination increases exposure of ezetimibe. If cholelithiasis is suspected in a patient receiving ezetimibe and a fenofibrate, gallbladder studies are indicated and alternative lipid-lowering therapy should be considered. (7.3, 12.3)

-----**USE IN SPECIFIC POPULATIONS**-----

- *Pregnancy*: May cause fetal harm (8.1)
- *Lactation*: Breastfeeding not recommended during treatment with ezetimibe and simvastatin. (8.2)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 5/2024

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 Important Dosage and Administration Information

2.2 Recommended Dosage in Adult Patients

2.3 Recommended Dosage in Pediatric Patients 10 Years of Age and Older with HeFH

2.4 Recommended Dosage in Patients with Renal Impairment

2.5 Dosage Modifications Due to Drug Interactions

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

5.1 Myopathy and Rhabdomyolysis

5.2 Immune-Mediated Necrotizing Myopathy

5.3 Hepatic Dysfunction

5.4 Increases in HbA1c and Fasting Serum Glucose Levels

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

6.2 Postmarketing Experience

7 DRUG INTERACTIONS

7.1 Drug Interactions that Increase the Risk of Myopathy and Rhabdomyolysis with ezetimibe and simvastatin tablets

7.2 Drug Interactions that Decrease the Efficacy of ezetimibe and simvastatin tablets

7.3 Ezetimibe and simvastatin tablets Effect on Other Drugs

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Renal Impairment

8.7 Hepatic Impairment

8.8 Chinese Patients

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Ezetimibe and Simvastatin

Ezetimibe and simvastatin is a combination of simvastatin and ezetimibe indicated:

- As an adjunct to diet to reduce elevated low density lipoprotein cholesterol (LDL-C):
 - In adults with primary hyperlipidemia.
 - In adults and pediatric patients aged 10 years and older with heterozygous familial hypercholesterolemia (HeFH).
- As an adjunct to other LDL-C-lowering therapies to reduce elevated LDL-C in adults with homozygous familial hypercholesterolemia (HoFH).

Simvastatin

Simvastatin, when used as a component of ezetimibe and simvastatin, is indicated to reduce the risk of total mortality by reducing risk of coronary heart disease death, non-fatal myocardial infarction and stroke, and the need for coronary and non-coronary revascularization procedures in adults with established coronary heart disease, cerebrovascular disease, peripheral vascular disease, and/or diabetes, who are at high risk of coronary heart disease events.

2 DOSAGE AND ADMINISTRATION

2.1 Important Dosage and Administration Information

- Take ezetimibe and simvastatin orally once daily in the evening with or without food.
- The maximum recommended dosage is ezetimibe and simvastatin 10/40 mg once daily. The ezetimibe and simvastatin 10/80 mg daily dosage is restricted to adult patients who have been taking ezetimibe and simvastatin 10/80 mg daily chronically (e.g., for 12 months or more) without evidence of muscle toxicity [*see Warnings and Precautions (5.1)*].
- For patients that require a high-intensity statin or are unable to achieve their LDL-C goal receiving ezetimibe and simvastatin 10/40 mg daily, prescribe alternative LDL-C-lowering treatment.
- If a dose is missed, take the missed dose as soon as possible. Do not double the next dose.
- Assess LDL-C when clinically appropriate, as early as 4 weeks after initiating ezetimibe and simvastatin, and adjust the dosage if necessary.

2.2 Recommended Dosage in Adult Patients

The recommended dosage range of ezetimibe and simvastatin 10/10 mg to 10/40 mg once a day.

2.3 Recommended Dosage in Pediatric Patients 10 Years of Age and Older with HeFH

The recommended dosage range of ezetimibe and simvastatin tablets 10/10 mg to 10/40 mg once a day.

2.4 Recommended Dosage in Patients with Renal Impairment

Renal impairment is a risk factor for statin-associated myopathy. Doses of ezetimibe and simvastatin tablets exceeding 10/20 mg should be used with caution and close monitoring in patients with moderate to severe renal impairment [*see Warnings and*

Precautions (5.1) and Use in Specific Populations (8.6)].

There are no dosage adjustment recommendations for patients with mild renal impairment.

2.5 Dosage Modifications Due to Drug Interactions

Concomitant use of ezetimibe and simvastatin tablets with the following drugs requires dosage modification of **ezetimibe and simvastatin tablets**[*see Warnings and Precautions (5.1) and Drug Interactions (7.1)*].

Patients taking Lomitapide

Reduce the dosage of ezetimibe and simvastatin tablets by 50%. Do not exceed ezetimibe and simvastatin tablets 10/20 mg once daily (or 10/40 mg once daily for patients who have previously taken ezetimibe and simvastatin tablets 10/80 mg daily chronically while taking lomitapide) [*see Dosage and Administration (2.1)*].

Patients taking Verapamil, Diltiazem, or Dronedarone

Do not exceed ezetimibe and simvastatin tablets 10/10 mg once daily.

Patients taking Amiodarone, Amlodipine, or Ranolazine

Do not exceed ezetimibe and simvastatin tablets 10/20 mg once daily.

Patients taking Bile Acid Sequestrants

In patients taking a bile acid sequestrant, administer ezetimibe and simvastatin tablets at least 2 hours before or 4 hours after the bile acid sequestrant.

3 DOSAGE FORMS AND STRENGTHS

Ezetimibe and simvastatin tablets

- 10/10, (ezetimibe, USP 10 mg/simvastatin, USP 10 mg tablets) are white to off-white, capsule shaped tablets debossed with “A029” on one side and plain on other side.
- 10/20, (ezetimibe, USP 10 mg/simvastatin, USP 20 mg tablets) are white to off-white, capsule shaped tablets debossed with “A028” on one side and plain on other side.
- 10/40, (ezetimibe, USP 10 mg/simvastatin, USP 40 mg tablets) are white to off-white, capsule shaped tablets and debossed with “A027” on one side and plain on other side.
- 10/80, (ezetimibe, USP 10 mg/simvastatin, USP 80 mg tablets) are white to off-white, capsule shaped tablets debossed with “A026” on one side and plain on other side.

4 CONTRAINDICATIONS

Ezetimibe and simvastatin tablets is contraindicated in the following conditions:

- Concomitant use of strong CYP3A4 inhibitors (select azole anti-fungals, macrolide antibiotics, antiviral medications, and nefazodone) [see *Drug Interactions (7.1)*].
- Concomitant use of cyclosporine, danazol, or danazol [see *Drug Interactions (7.1)*].
- Acute liver failure or decompensated cirrhosis [see *Warnings and Precautions (5.3)*].
- Hypersensitivity to simvastatin, ezetimibe, or any excipients in ezetimibe and simvastatin tablets. Hypersensitivity reactions, including anaphylaxis, angioedema, and Stevens-Johnson syndrome, have been reported [see *Adverse Reactions (6.2)*].

5 WARNINGS AND PRECAUTIONS

5.1 Myopathy and Rhabdomyolysis

Ezetimibe and simvastatin tablets may cause myopathy and rhabdomyolysis. Acute kidney injury secondary to myoglobinuria and rare fatalities have occurred as a result of rhabdomyolysis in patients treated with statins, including ezetimibe and simvastatin tablets.

In clinical trials of 24,747 simvastatin-treated patients with a median follow-up of 4 years, the incidence of myopathy, defined as unexplained muscle weakness, pain, or tenderness accompanied by creatinine kinase (CK) increases greater than ten times the upper limit of normal (10 X ULN), were approximately 0.03%, 0.08%, and 0.61% in patients treated with simvastatin 20 mg, 40 mg, and 80 mg daily, respectively. In another clinical trial of 12,064 simvastatin-treated patients (with a history of myocardial infarction) with a mean follow-up of 6.7 years, the incidences of myopathy in patients taking simvastatin 20 mg and 80 mg daily were approximately 0.02% and 0.9%, respectively. The incidences of rhabdomyolysis (defined as myopathy with a CK >40 X ULN) in patients taking simvastatin 20 mg and 80 mg daily were approximately 0% and 0.4%, respectively [see *Adverse Reactions (6.1)*].

In the Trial of Heart and Renal Protection (SHARP), 9,270 patients with chronic kidney disease were allocated to receive ezetimibe and simvastatin tablets 10/20 mg daily (n=4650) or placebo (n=4620). During a median follow-up period of 4.9 years, the incidence of myopathy (defined as unexplained muscle weakness or pain with a serum creatine kinase [CK] >10 times upper limit of normal [ULN]) was 0.2% for ezetimibe and simvastatin tablets and 0.1% for placebo: the incidence of rhabdomyolysis (defined as myopathy with a CK > 40 times ULN) was 0.09% for ezetimibe and simvastatin tablets and 0.02% for placebo.

In postmarketing experience with ezetimibe, cases of myopathy and rhabdomyolysis have been reported. Most patients who developed rhabdomyolysis were taking a statin prior to initiating ezetimibe. However, rhabdomyolysis has been reported with ezetimibe monotherapy and with the addition of ezetimibe to agents known to be associated with increased risk of rhabdomyolysis, such as fibric acid derivatives. Ezetimibe and simvastatin tablets and a fenofibrate, if taking concomitantly, should both be immediately discontinued if myopathy is diagnosed or suspected.

Risk Factors for Myopathy

Risk factors for myopathy include age 65 years or greater, uncontrolled hypothyroidism, renal impairment, concomitant use with certain other drugs (including other lipid-lowering therapies), and higher ezetimibe and simvastatin tablets dosage; Chinese patients on ezetimibe and simvastatin tablets may be at higher risk for myopathy [see *Contraindications (4)*, *Drug Interactions (7.1)*, and *Use in Specific Populations (8.8)*]. The risk of myopathy is increased by elevated plasma levels of simvastatin and simvastatin acid. The risk is also greater in patients taking ezetimibe and simvastatin tablets 80 mg daily compared with patients taking lower ezetimibe and simvastatin tablets dosages and compared with patients using other statins with similar or greater LDL-C-lowering efficacy [see *Adverse Reactions (6.1)*].

Steps to Prevent or Reduce the Risk of Myopathy and Rhabdomyolysis

The concomitant use of strong CYP3A4 inhibitors with ezetimibe and simvastatin tablets is contraindicated. If short-term treatment with strong CYP3A4 inhibitors is required, temporarily suspend ezetimibe and simvastatin tablets during the duration of strong CYP3A4 inhibitor treatment. The concomitant use of ezetimibe and simvastatin tablets with gemfibrozil, cyclosporine, or danazol is also contraindicated [see *Contraindications (4)* and *Drug Interactions (7.1)*].

Ezetimibe and simvastatin tablets dosage modifications are recommended for patients taking lomitapide, verapamil, diltiazem, dronedarone, amiodarone, amlodipine or ranolazine [see *Dosage and Administration (2.5)*]. Ezetimibe and simvastatin tablets use should be temporarily suspended in patients taking daptomycin. Lipid modifying doses (≥ 1 gram/day) of niacin, fibrates, colchicine, and grapefruit juice may also increase the risk of myopathy and rhabdomyolysis [see *Drug Interactions (7.1)*].

Use the 80 mg daily dosage of ezetimibe and simvastatin tablets only in patients who have been taking simvastatin 80 mg daily chronically without evidence of muscle toxicity [see *Dosage and Administration (2.1)*]. If patients treated with ezetimibe and simvastatin tablets 80 mg are prescribed an interacting drug that increases the risk for myopathy and rhabdomyolysis, switch to an alternate statin [see *Drug Interactions (7.1)*].

Discontinue ezetimibe and simvastatin tablets if markedly elevated CK levels occur or if myopathy is either diagnosed or suspected. Muscle symptoms and CK increases may resolve if ezetimibe and simvastatin tablets is discontinued. Temporarily discontinue ezetimibe and simvastatin tablets in patients experiencing an acute or serious condition at high risk of developing renal failure secondary to rhabdomyolysis, e.g., sepsis; shock; severe hypovolemia; major surgery; trauma; severe metabolic, endocrine, or electrolyte disorders; or uncontrolled epilepsy.

Inform patients of the risk of myopathy and rhabdomyolysis when starting or increasing the ezetimibe and simvastatin tablets dosage and advise patients receiving ezetimibe and simvastatin tablets 80 mg of the increased risk of myopathy and rhabdomyolysis. Instruct patients to promptly report any unexplained muscle pain, tenderness or weakness, particularly if accompanied by malaise or fever.

5.2 Immune-Mediated Necrotizing Myopathy

There have been rare reports of immune-mediated necrotizing myopathy (IMNM), an autoimmune myopathy, associated with statin use, including reports of recurrence when the same or a different statin was administered. IMNM is characterized by proximal muscle weakness and elevated serum creatine kinase that persist despite discontinuation of statin treatment; positive anti-HMG CoA reductase antibody; muscle biopsy showing necrotizing myopathy without significant inflammation; and improvement with immunosuppressive agents. Additional neuromuscular and serologic testing may be necessary. Treatment with immunosuppressive agents may be required. Discontinue ezetimibe and simvastatin tablets if IMNM is suspected.

5.3 Hepatic Dysfunction

Increases in serum transaminases have been reported with use of ezetimibe and simvastatin tablets [see *Adverse Reactions (6.1)*]. In most cases, these changes appeared soon after initiation, were transient, were not accompanied by symptoms, and resolved or improved on continued therapy or after a brief interruption in therapy. Marked persistent increases of hepatic transaminases have also occurred with ezetimibe and simvastatin tablets. There have been rare postmarketing reports of fatal and non-fatal hepatic failure in patients taking statins, including ezetimibe and simvastatin tablets.

In three placebo-controlled, 12-week trials, the incidence of consecutive elevations (≥ 3 X ULN) in serum transaminases was 1.7% overall for patients treated with ezetimibe and simvastatin tablets and appeared to be dose-related with an incidence of 2.6% for patients treated with ezetimibe and simvastatin tablets 10/80. In controlled long-term (48-week) extensions, which included both newly-treated and previously-treated patients, the incidence of consecutive elevations (≥ 3 X ULN) in serum transaminases was 1.8% overall and 3.6% for patients treated with ezetimibe and simvastatin tablets 10/80. These elevations in transaminases were generally asymptomatic, not associated with cholestasis, and returned to baseline after discontinuation of therapy or with continued treatment.

In SHARP, 9,270 patients with chronic kidney disease were allocated to receive ezetimibe and simvastatin tablets 10/20 mg daily (n=4650), or placebo (n=4620). During a median follow-up period of 4.9 years, the incidence of consecutive elevations of transaminases (>3 X ULN) was 0.7% for ezetimibe and simvastatin tablets and 0.6% for placebo.

Patients who consume substantial quantities of alcohol and/or have a history of liver disease may be at increased risk for hepatic injury.

Consider liver enzyme testing before ezetimibe and simvastatin tablets initiation and when clinically indicated thereafter. Ezetimibe and simvastatin tablets is contraindicated in patients with acute liver failure or decompensated cirrhosis [see *Contraindications (4)*]. If serious hepatic injury with clinical symptoms and/or hyperbilirubinemia or jaundice occurs, promptly discontinue ezetimibe and simvastatin tablets.

5.4 Increases in HbA1c and Fasting Serum Glucose Levels

Increases in HbA1c and fasting serum glucose levels have been reported with statins, including ezetimibe and simvastatin tablets. Optimize lifestyle measures, including regular exercise, maintaining a healthy body weight, and making healthy food choices.

6 ADVERSE REACTIONS

The following serious adverse reactions are discussed in greater detail in other sections of the label:

- Myopathy and Rhabdomyolysis [*see Warnings and Precautions (5.1)*]
- Immune-Mediated Necrotizing Myopathy [*see Warnings and Precautions (5.2)*]
- Hepatic Dysfunction [*see Warnings and Precautions (5.3)*]
- Increases in HbA1c and Fasting Serum Glucose Levels [*see Warnings and Precautions (5.4)*]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Ezetimibe and Simvastatin Tablets

In the ezetimibe and simvastatin tablets (ezetimibe and simvastatin) placebo-controlled clinical trials database of 1,420 patients (age range 20 to 83 years, 52% female, 87% White, 3% Black or African American, 3% Asians, 5% other races identified as Hispanic or Latino ethnicity) with a median treatment duration of 27 weeks, 5% of patients on ezetimibe and simvastatin tablets and 2.2% of patients on placebo discontinued due to adverse reactions.

The most commonly reported adverse reactions (incidence $\geq 2\%$ and greater than placebo) in controlled clinical trials were: headache (5.8%), increased ALT (3.7%), myalgia (3.6%), upper respiratory tract infection (3.6%), and diarrhea (2.8%). The most common adverse reactions in the group treated with ezetimibe and simvastatin tablets that led to treatment discontinuation and occurred at a rate greater than placebo were: increased ALT (0.9%), myalgia (0.6%), increased AST (0.4%), and back pain (0.4%). Ezetimibe and simvastatin tablets has been evaluated for safety in more than 10,189 patients in clinical trials.

Table 1 summarizes the frequency of clinical adverse reactions reported in $\geq 2\%$ of patients treated with ezetimibe and simvastatin tablets (n=1420) and at an incidence greater than placebo from four placebo-controlled trials.

Table 1*: Adverse Reactions Reported $\geq 2\%$ of Patients Treated with Ezetimibe and Simvastatin Tablets at an Incidence Greater than Placebo Regardless of Causality

	% Placebo N = 371	% Ezetimibe 10 mg N = 302	% Simvastatin† N = 1234	% Ezetimibe and Simvastatin Tablets † N = 1420
Headache	5.4	6.0	5.9	5.8
Upper respiratory tract infection	2.7	5.0	5.0	3.6
Myalgia	2.4	2.3	2.6	3.6
Diarrhea	2.2	5.0	3.7	2.8
Pain in extremity	1.3	3.0	2.0	2.3
Influenza	0.8	1.0	1.9	2.3

*Includes two placebo-controlled combination studies in which the active ingredients equivalent to ezetimibe and simvastatin tablets were coadministered and two placebo-controlled studies in which ezetimibe and simvastatin tablets was administered.

†All doses.

Study of Heart and Renal Protection

In SHARP, 9,270 patients were allocated to ezetimibe and simvastatin tablets 10/20 mg daily (n=4650) or placebo (n=4620) for a median follow-up period of 4.9 years. The proportion of patients who permanently discontinued trial treatment as a result of either an adverse event or abnormal safety blood result was 10.4% vs. 9.8% among patients allocated to ezetimibe and simvastatin tablets and placebo, respectively. Comparing those allocated to ezetimibe and simvastatin tablets vs. placebo, the incidence of myopathy (defined as unexplained muscle weakness or pain with a serum CK >10 times ULN) was 0.2% vs. 0.1% and the incidence of rhabdomyolysis (defined as myopathy with a CK >40 times ULN) was 0.09% vs. 0.02%, respectively. Consecutive elevations of transaminases (>3 X ULN) occurred in 0.7% vs. 0.6%, respectively. Patients were asked about the occurrence of unexplained muscle pain or weakness at each trial visit: 21.5% vs. 20.9% patients ever reported muscle symptoms in the ezetimibe and simvastatin tablets and placebo groups, respectively. Cancer was diagnosed during the trial in 9.4% vs. 9.5% of patients assigned to ezetimibe and simvastatin tablets and placebo, respectively.

Ezetimibe

Other adverse reactions reported with ezetimibe in placebo-controlled studies, regardless of causality assessment: *Musculoskeletal system disorders*: arthralgia; *Infections and infestations*: sinusitis; *Body as a whole – general disorders*: fatigue.

Simvastatin

In a clinical outcome trial in which 12,064 adult patients with a history of myocardial infarction were treated with simvastatin (mean follow-up 6.7 years), the incidence of myopathy (defined as unexplained muscle weakness or pain with a serum creatine kinase [CK] >10 times (1200 U/L) upper limit of normal [ULN]) in patients taking simvastatin 20 mg and 80 mg daily was approximately 0.02% and 0.9% respectively. The

incidence of rhabdomyolysis (defined as myopathy with a CK >40 times ULN) in patients taking simvastatin 20 mg and 80 mg daily was approximately 0% and 0.4%. The incidence of myopathy and rhabdomyolysis, was highest during the first year and then notably decreased during the subsequent years of treatment. In this trial, patients were carefully monitored and some interacting medicinal products were excluded. Other adverse reactions reported with simvastatin in placebo-controlled clinical trials: atrial fibrillation; vertigo; abdominal pain, constipation, dyspepsia, flatulence, gastritis; eczema, rash; diabetes mellitus; bronchitis, sinusitis, urinary tract infections; asthenia, edema/swelling; and insomnia.

Laboratory Tests

Marked persistent increases of hepatic serum transaminases have been noted [see *Warnings and Precautions (5.3)*]. Elevated alkaline phosphatase and γ -glutamyl transpeptidase have been reported. About 5% of patients taking simvastatin had elevations of CK levels of 3 or more times the normal value on one or more occasions. This was attributable to the noncardiac fraction of CK [see *Warnings and Precautions (5.1)*].

6.2 Postmarketing Experience

The following adverse reactions have been identified during post-approval use of ezetimibe and simvastatin tablets. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Body as Whole: fever, chills, malaise, asthenia

Blood and Lymphatic System Disorders: anemia, thrombocytopenia, leukopenia, hemolytic anemia, positive ANA, ESR increase, eosinophilia

Gastrointestinal Disorders: pancreatitis, nausea, vomiting

Hepatobiliary Disorders: cholelithiasis, cholecystitis, elevations in liver transaminases including elevations more than 5 X ULN, hepatitis/jaundice, fatal and non-fatal hepatic failure

Immune System Disorders: hypersensitivity syndrome including: anaphylaxis, angioedema, lupus erythematosus-like syndrome, dermatomyositis, vasculitis

Musculoskeletal and Connective Tissue Disorders: muscle cramps, immune-mediated necrotizing myopathy, rhabdomyolysis, myalgia, arthralgia, polymyalgia rheumatica, arthritis, elevated creatine phosphokinase

Nervous System Disorders: dizziness, depression, paresthesia, peripheral neuropathy, rare reports of cognitive impairment (e.g., memory loss, forgetfulness, amnesia, memory impairment, confusion) associated with statin use. Cognitive impairment was generally nonserious, and reversible upon statin discontinuation, with variable times to symptom onset (1 day to years) and symptom resolution (median of 3 weeks). There have been rare reports of new-onset or exacerbation of myasthenia gravis, including ocular myasthenia, and reports of recurrence when the same or a different statin was administered.

Skin and Subcutaneous Tissue Disorders: rash, pruritus, alopecia, a variety of skin changes (e.g., nodules, discoloration, dryness of skin/mucous membranes, changes to hair/nails), purpura, lichen planus, urticaria, photosensitivity, flushing, toxic epidermal necrolysis, erythema multiforme, including Stevens-Johnson syndrome.

Respiratory and Thoracic: interstitial lung disease, dyspnea

Reproductive System Disorders: erectile dysfunction

7 DRUG INTERACTIONS

Ezetimibe and Simvastatin Tablets

7.1 Drug Interactions that Increase the Risk of Myopathy and Rhabdomyolysis with ezetimibe and simvastatin tablets

Ezetimibe and simvastatin tablets is a substrate of CYP3A4 and of the transport protein OATP1B1. Ezetimibe and simvastatin tablets plasma levels can be significantly increased with concomitant administration of inhibitors of CYP3A4 and OATP1B1.

Table 2 includes a list of drugs that increase the risk of myopathy and rhabdomyolysis when used concomitantly with ezetimibe and simvastatin tablets and instructions for preventing or managing them [see *Warnings and Precautions (5.1) and Clinical Pharmacology (12.3)*].

Table 2: Drug Interactions that Increase the Risk of Myopathy and Rhabdomyolysis with Ezetimibe and Simvastatin Tablets

Strong CYP3A4 inhibitors	
<i>Clinical Impact:</i>	Simvastatin is a substrate of CYP3A4. Concomitant use of strong CYP3A4 inhibitors with ezetimibe and simvastatin tablets increases simvastatin exposure and increases the risk of myopathy and rhabdomyolysis, particularly with higher ezetimibe and simvastatin tablets dosages.
<i>Intervention:</i>	Concomitant use of strong CYP3A4 inhibitors with ezetimibe and simvastatin tablets is contraindicated [see <i>Contraindications (4)</i>]. If treatment with a CYP3A4 inhibitor is unavoidable, suspend ezetimibe and simvastatin tablets during the course of strong CYP3A4 inhibitor treatment.
<i>Examples:</i>	Select azole anti-fungals (e.g., itraconazole, ketoconazole, posaconazole, and voriconazole), select macrolide antibiotics (e.g., erythromycin and clarithromycin, telithromycin), select HIV protease inhibitors (e.g., nelfinavir, ritonavir, and darunavir/ritonavir), select HCV protease inhibitors (e.g., boceprevir and telaprevir), cobicistat-containing products, and nefazodone.
Cyclosporine, Danazol, or Gemfibrozil	
<i>Clinical Impact:</i>	The risk of myopathy and rhabdomyolysis is increased with

	concomitant use of cyclosporine, danazol, or gemfibrozil with ezetimibe and simvastatin tablets. Gemfibrozil may cause myopathy when given alone.
<i>Intervention:</i>	Concomitant use of cyclosporine, danazol, or gemfibrozil with ezetimibe and simvastatin tablets is contraindicated [see <i>Contraindications (4)</i>].
Amiodarone, Dronedarone, Ranolazine, or Calcium Channel Blockers	
<i>Clinical Impact:</i>	The risk of myopathy and rhabdomyolysis is increased by concomitant use of amiodarone, dronedarone, ranolazine, or calcium channel blockers with ezetimibe and simvastatin tablets.
<i>Intervention:</i>	For patients taking verapamil, diltiazem, or dronedarone, do not exceed ezetimibe and simvastatin tablets 10/10 mg daily. For patients taking amiodarone, amlodipine, or ranolazine, do not exceed ezetimibe and simvastatin tablets 10/20 mg daily [see <i>Dosage and Administration (2.3)</i>].
Lomitapide	
<i>Clinical Impact:</i>	Simvastatin exposure is approximately doubled with concomitant use of lomitapide and the risk of myopathy and rhabdomyolysis is increased.
<i>Intervention:</i>	Reduce the dose of ezetimibe and simvastatin tablets by 50% if initiating lomitapide. Do not exceed ezetimibe and simvastatin tablets 10/20 mg daily (or ezetimibe and simvastatin tablets 10/40 mg daily for patients who have previously taken ezetimibe and simvastatin tablets 10/80 mg daily chronically) while taking lomitapide [see <i>Dosage and Administration (2.1, 2.3)</i>].
Daptomycin	
<i>Clinical Impact:</i>	Cases of rhabdomyolysis have been reported with simvastatin administered with daptomycin. Both ezetimibe and simvastatin tablets and daptomycin can cause myopathy and rhabdomyolysis when given alone and the risk of myopathy and rhabdomyolysis may be increased by coadministration.
<i>Intervention:</i>	If treatment with daptomycin is required, consider temporarily suspending ezetimibe and simvastatin tablets during the course of daptomycin treatment.
Niacin	
<i>Clinical Impact:</i>	Cases of myopathy and rhabdomyolysis have been observed with concomitant use of lipid modifying dosages of niacin-containing products (≥ 1 gram/day niacin) with ezetimibe and simvastatin tablets. The risk of myopathy is greater in Chinese patients. In a clinical trial (median follow-up 3.9 years) of patients at high risk of CVD and with well-controlled LDL-C levels on simvastatin 40 mg/day with or without ezetimibe 10 mg/day, there was no incremental benefit on cardiovascular outcomes with the addition of lipid-modifying doses of niacin
<i>Intervention:</i>	Concomitant use of ezetimibe and simvastatin tablets with lipid-modifying dosages of niacin is not recommended in Chinese patients [see <i>Use in Specific Populations (8.8)</i>]. For non-Chinese patients, consider if the benefit of using lipid-modifying doses of niacin concomitantly with ezetimibe and simvastatin tablets outweighs the increased risk of myopathy and rhabdomyolysis. If

	concomitant use is decided, monitor patients for signs and symptoms of myopathy, particularly during initiation of therapy and during upward dose titration of either drug.
Fibrates (other than Gemfibrozil)	
<i>Clinical Impact:</i>	Fibrates may cause myopathy when given alone. The risk of myopathy and rhabdomyolysis is increased with concomitant use of fibrates with ezetimibe and simvastatin tablets.
<i>Intervention:</i>	Consider if the benefit of using fibrates concomitantly with ezetimibe and simvastatin tablets outweighs the increased risk of myopathy and rhabdomyolysis. If concomitant use is decided, monitor patients for signs and symptoms of myopathy, particularly during initiation of therapy and during upward dose titration of either drug.
Colchicine	
<i>Clinical Impact:</i>	Cases of myopathy and rhabdomyolysis have been reported with concomitant use of colchicine with ezetimibe and simvastatin tablets.
<i>Intervention:</i>	Consider if the benefit of using colchicine concomitantly with ezetimibe and simvastatin tablets outweighs the increased risk of myopathy and rhabdomyolysis. If concomitant use is decided, monitor patients for signs and symptoms of myopathy, particularly during initiation of therapy and during upward dose titration of either drug.
Grapefruit Juice	
<i>Clinical Impact:</i>	Grapefruit juice can raise the plasma levels of simvastatin and may increase the risk of myopathy and rhabdomyolysis.
<i>Intervention:</i>	Avoid grapefruit juice when taking ezetimibe and simvastatin tablets.

7.2 Drug Interactions that Decrease the Efficacy of ezetimibe and simvastatin tablets

Table 3 presents drug interactions that may decrease the efficacy of ezetimibe and simvastatin tablets and instructions for preventing or managing them.

Table 3: Drug Interactions that Decrease the Efficacy of Ezetimibe and Simvastatin Tablets

Bile Acid Sequestrants	
<i>Clinical Impact:</i>	Concomitant cholestyramine administration decreased the mean exposure of total ezetimibe approximately 55%. The incremental LDL-C reduction due to adding ezetimibe and simvastatin tablets to cholestyramine may be reduced by this interaction [see <i>Clinical Pharmacology (12.3)</i>].
<i>Intervention:</i>	In patients taking a bile acid sequestrant, administer ezetimibe and simvastatin tablets at least 2 hours before or at least 4 hours after cholestyramine [see <i>Dosage and</i>

7.3 Ezetimibe and simvastatin tablets Effect on Other Drugs

Table 4 presents ezetimibe and simvastatin tablets effect on other drugs and instructions for preventing or managing them.

Table 4: Ezetimibe and Simvastatin Tablets Effects on Other Drugs

Coumarin Anticoagulants	
<i>Clinical Impact:</i>	Ezetimibe and simvastatin tablets may potentiate the effect of coumarin anticoagulants and increase the INR. The concomitant use of simvastatin (20 to 40 mg) and coumarin anticoagulants increased the INR from a baseline of 1.7 to 1.8 in healthy subjects and from 2.6 to 3.4 in patients with hyperlipidemia. There are postmarketing reports of clinically evident bleeding and/or increased INR in patients taking concomitant statins (with or without ezetimibe) and coumarin anticoagulants.
<i>Intervention:</i>	In patients taking coumarin anticoagulants, obtain an INR before starting ezetimibe and simvastatin tablets and frequently enough after initiation, dose titration, or discontinuation to ensure that no significant alteration in INR occurs. Once the INR is stable, monitor INR at regularly recommended intervals.
Digoxin	
<i>Clinical Impact:</i>	Concomitant use of digoxin with ezetimibe and simvastatin tablets may result in elevated plasma digoxin concentrations [see <i>Clinical Pharmacology (12.3)</i>].
<i>Intervention:</i>	Monitor digoxin levels in patients taking digoxin when ezetimibe and simvastatin tablets is initiated.
Fenofibrates	
<i>Clinical Impact:</i>	Both ezetimibe and fenofibrates may increase cholesterol excretion into the bile, leading to cholelithiasis.
<i>Intervention:</i>	If cholelithiasis is suspected in a patient receiving ezetimibe and simvastatin tablets and a fenofibrate, gallbladder studies are indicated and alternative lipid-lowering therapy should be considered [see the product labeling for fenofibrate and fenofibric acid].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Ezetimibe and simvastatin tablets

Risk Summary

Discontinue ezetimibe and simvastatin tablets when pregnancy is recognized. Alternatively, consider the ongoing therapeutic needs of the individual patient.

Ezetimibe and simvastatin tablets decreases synthesis of cholesterol and possibly other biologically active substances derived from cholesterol; therefore, ezetimibe and simvastatin tablets may cause fetal harm when administered to pregnant patients based on the mechanism of action [see *Clinical Pharmacology (12.1)*]. In addition, treatment of hyperlipidemia is not generally necessary during pregnancy. Atherosclerosis is a chronic process and the discontinuation of lipid-lowering drugs during pregnancy should have little impact on the outcome of long-term therapy of primary hyperlipidemia for most patients.

Available data from case series and prospective and retrospective observational cohort studies over decades of use with statins in pregnant women have not identified a drug-associated risk of major congenital malformations. Published data from prospective and retrospective observational cohort studies with ezetimibe and simvastatin tablets use in pregnant women are insufficient to determine if there is a drug-associated risk of miscarriage (see *Data*).

In animal reproduction studies, no adverse developmental effects were observed in pregnant rats or rabbits orally administered simvastatin during the period of organogenesis at doses that resulted in 2.5 and 2 times, respectively, the human exposure at the maximum recommended human dosage of 80 mg/day, based on body surface area (mg/m²). In animal reproduction studies, no adverse developmental effects were observed in pregnant rats and rabbits orally administered ezetimibe during the period of organogenesis at doses that resulted in up to 10 and 150 times, respectively, the human exposure at the MRHD, based on AUC (see *Data*).

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Data

Human Data

Ezetimibe

There are insufficient data on ezetimibe use in pregnant women to evaluate for a drug-associated risk of major birth defects, miscarriage, or adverse maternal or fetal outcomes.

Simvastatin

A Medicaid cohort linkage trial of 1,152 statin-exposed pregnant women compared to 886,996 controls did not find a significant teratogenic effect from maternal use of statins in the first trimester of pregnancy, after adjusting for potential confounders –

including maternal age, diabetes mellitus, hypertension, obesity, and alcohol and tobacco use – using propensity score-based methods. The relative risk of congenital malformations between the group with statin use and the group with no statin use in the first trimester was 1.07 (95% confidence interval 0.85 to 1.37) after controlling for confounders, particularly pre-existing diabetes mellitus. There were also no statistically significant increases in any of the organ-specific malformations assessed after accounting for confounders. In the majority of pregnancies, statin treatment was initiated prior to pregnancy and was discontinued at some point in the first trimester when pregnancy was identified. Trial limitations include reliance on physician coding to define the presence of a malformation, lack of control for certain confounders such as body mass index, use of prescription dispensing as verification for the use of a statin, and lack of information on non-live births.

Animal Data

Ezetimibe

In oral (gavage) embryo-fetal development studies of ezetimibe conducted in rats (gestation days 6 to 15) and rabbits (gestation days 7 to 19), there was no evidence of maternal toxicity or embryoletality at any dose tested (250, 500, 1,000 mg/kg/day) at exposure equivalent to 10 to 150 times the clinical exposure, based on AUC, in rats and rabbits. In rats, increased incidences of common fetal skeletal findings (extra pair of thoracic ribs, unossified cervical vertebral centra, shortened ribs) were observed at 1,000 mg/kg/day (~10 times the human exposure at 10 mg daily based on AUC_{0-24hr} for total ezetimibe). In rabbits treated with ezetimibe, an increased incidence of extra thoracic ribs was observed at 1,000 mg/kg/day (150 times the human exposure at 10 mg daily based on AUC_{0-24hr} for total ezetimibe).

The animal-to-human exposure multiple for total ezetimibe at the no-observed effect level was 6 times for rat and 134 times for rabbit. Fetal exposure to ezetimibe (conjugated and unconjugated) was confirmed in subsequent placental transfer studies conducted using a maternal dose of 1,000 mg/kg/day. The fetal maternal plasma exposure ratio (total ezetimibe) was 1.5 for rats on gestation day 20 and 0.03 for rabbits on gestation day 22.

The effect of ezetimibe on prenatal and postnatal development and maternal function was evaluated in pregnant rats at doses of 100, 300 or 1,000 mg/kg/day (gestation day 6 through lactation day 21). No maternal toxicity or adverse developmental outcomes were observed up to and including the highest dose tested (17 times the human exposure at 10 mg daily based on AUC_{0-24hr} for total ezetimibe).

Multiple-dose studies of ezetimibe given in combination with statins in rats and rabbits during organogenesis resulted in higher ezetimibe and statin exposures. Reproductive findings occurred at lower doses in combination therapy compared to monotherapy.

Simvastatin

Simvastatin was given to pregnant rats at doses of 6.25, 12.5 and 25 mg/kg/day (0.6 times, 1.3 times, and 2.5 times, respectively, the maximum recommended dosage of 80 mg/day when normalized to body surface area) from gestation days 6 to 17 and to pregnant rabbits from gestation days 6 to 18 at doses of 2.5, 5, and 10 mg/kg/day (0.5 times, 1 times, and 2 times, respectively, the maximum recommended dosage of 80

mg/day when normalized to body surface area). For both species, there was no evidence of maternal toxicity, or embryoletality. In rats, mean fetal body weights in the 25 mg/kg/day group were decreased 5.4%. Similar fetal body weight effects were not observed in rabbits.

Simvastatin doses of 6.25, 12.5 and 25 mg/kg/day (0.6 times, 1.3 times, and 2.5 times, respectively, the maximum recommended dosage of 80 mg/day when normalized to body surface area) were given to pregnant rats from gestation day 15 to lactation day 21. Slight decreases in maternal body weight gain and pup postnatal day 0 weight were observed in the 25 mg/kg/day dose group. Mean body weight gain of pups during lactation was slightly decreased at doses ≥ 12.5 mg/kg/day. Post weaning weight, behavior, reproductive performance and fertility of the offspring were not affected at any dose tested.

Placental transfer of simvastatin was not evaluated in rats or rabbits. However, it has been shown that other drugs in this class cross the placenta.

8.2 Lactation

Risk Summary There is no information about the presence of ezetimibe or simvastatin in human breast milk, the effects of the drug on the breastfed infant or the effect of the drug on milk production. However, it has been shown that other statins pass into human milk. Statins, including ezetimibe and simvastatin tablets, decrease cholesterol synthesis and possibly the synthesis of other biologically active substances derived from cholesterol and may cause harm to the breast fed infant.

Because of the potential for serious adverse reactions in a breastfed infant, based on the mechanism of action, advise patients that breastfeeding is not recommended during treatment with ezetimibe and simvastatin tablets [see *Use in Specific Populations (8.2) and Clinical Pharmacology (12.1)*].

Data

Animal Data

Ezetimibe was present in the milk of lactating rats. The pup to maternal plasma ratio for total ezetimibe was 0.5 on lactation day 12.

8.4 Pediatric Use

The safety and effectiveness of ezetimibe in combination with a statin as an adjunct to diet to reduce LDL-C have been established in pediatric patients 10 years of age and older with HeFH. Use of ezetimibe and simvastatin tablets for this indication is based on a double-blind, placebo-controlled clinical trial in 248 pediatric patients (142 males and 106 postmenarchal females) 10 years of age and older with HeFH [see *Clinical Studies (14)*]. In this limited controlled trial, there was no significant effect on growth or sexual maturation in the adolescent males or females, or on menstrual cycle length in females.

The safety and effectiveness of ezetimibe and simvastatin tablets have not been established in pediatric patients younger than 10 years of age with HeFH, or in pediatric patients with other types of hyperlipidemia.

8.5 Geriatric Use

Advanced age (≥ 65 years) is a risk factor for ezetimibe and simvastatin tablets - associated myopathy and rhabdomyolysis. Dose selection for an elderly patient should be cautious, recognizing the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy and the higher risk of myopathy. Monitor geriatric patients receiving ezetimibe and simvastatin tablets for the increased risk of myopathy [see *Warnings and Precautions (5.1)*].

Of the 10,189 patients who received ezetimibe and simvastatin tablets in clinical studies, 3,242 (32%) were 65 and older (this included 844 (8%) who were 75 and older). No overall differences in safety or effectiveness were observed between these subjects and younger subjects, and other reported clinical experience has not identified differences in responses between the elderly and younger patients but greater sensitivity of some older individuals cannot be ruled out.

8.6 Renal Impairment

Renal impairment is a risk factor for myopathy and rhabdomyolysis. Monitor all patients with renal impairment for development of myopathy. Doses of ezetimibe and simvastatin tablets exceeding 10/20 mg should be used with caution and close monitoring in patients with moderate to severe renal impairment [see *Dosage and Administration (2.4) and Warnings and Precautions (5.1)*].

In the SHARP trial of 9,270 patients with moderate to severe renal impairment (6,247 non-dialysis patients with median serum creatinine 2.5 mg/dL and median estimated glomerular filtration rate 25.6 mL/min/1.73 m², and 3,023 dialysis patients), the incidence of serious adverse events, adverse events leading to discontinuation of trial treatment, or adverse events of special interest (musculoskeletal adverse events, liver enzyme abnormalities, incident cancer) was similar between patients ever assigned to ezetimibe and simvastatin tablets 10/20 mg (n=4650) or placebo (n=4620) during a median follow-up of 4.9 years.

8.7 Hepatic Impairment

Ezetimibe and simvastatin tablets is contraindicated in patients with acute liver failure or decompensated cirrhosis. [See *Contraindications (4) and Warnings and Precautions (5.3)*.]

8.8 Chinese Patients

In a clinical trial in which patients at high risk of CVD were treated with simvastatin 40 mg/day (median follow-up 3.9 years), the incidence of myopathy was approximately 0.05% for non-Chinese patients (n=7367) compared with 0.24% for Chinese patients (n=5468). In this trial the incidence of myopathy for Chinese patients on simvastatin 40 mg/day or ezetimibe and simvastatin 10/40 mg/day coadministered with extended-release niacin 2 g/day was 1.24%.

Chinese patients may be at higher risk for myopathy, monitor these patients appropriately. Coadministration of ezetimibe and simvastatin tablets with lipid-modifying doses of niacin-containing products (≥ 1 g/day niacin) is not recommended in Chinese patients [see *Warnings and Precautions (5.1) and Drug Interactions (7.1)*].

10 OVERDOSAGE


No specific antidotes for ezetimibe and simvastatin tablets are known. In the event of an overdose with ezetimibe and simvastatin tablets, consider contacting the Poison Help Line (1-800-222-1222) or a medical toxicologist for overdosage management recommendations.

11 DESCRIPTION

Ezetimibe and simvastatin tablets contains ezetimibe, USP, a dietary cholesterol absorption inhibitor, and simvastatin, USP, an HMG-CoA reductase inhibitor.


The chemical name of ezetimibe, USP is 1-(4-fluorophenyl)-3(R)-[3-(4-fluorophenyl)-3(S)-hydroxypropyl]-4(S)-(4-hydroxyphenyl)-2-azetidinone. The molecular formula is $C_{24}H_{21}F_2NO_3$ and its molecular weight is 409.43.

Ezetimibe, USP is a white to off white, crystalline powder, hygroscopic that is soluble in methanol. Its structural formula is:



Simvastatin, USP, an inactive lactone, is hydrolyzed to the corresponding β -hydroxyacid form, which is an inhibitor of HMG-CoA reductase. Simvastatin, USP is butanoic acid, 2, 2-dimethyl-,1,2,3,7,8,8a-hexahydro-3,7dimethyl-8-[2-(tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)-ethyl]-1-naphthalenyl ester, [1S-[1 α ,3 α ,7 β ,8 β (2S*,4S*),-8a β]]. The molecular formula of simvastatin, USP is $C_{25}H_{38}O_5$ and its molecular weight is 418.57.

Simvastatin, USP is a white to off-white powder that is freely soluble in chloroform, methanol and alcohol. Sparingly soluble in Propylene glycol, very slightly soluble in Hexane. Practically insoluble in water. Its structural formula is:



Ezetimibe and simvastatin tablets is available for oral use as tablets containing 10 mg of ezetimibe, USP, and 10 mg of simvastatin, USP (ezetimibe and simvastatin tablets

10/10), 20 mg of simvastatin, USP (ezetimibe and simvastatin tablets 10/20), 40 mg of simvastatin, USP (ezetimibe and simvastatin tablets 10/40), or 80 mg of simvastatin, USP (ezetimibe and simvastatin tablets 10/80). Each tablet contains the following inactive ingredients: Lactose Monohydrate, Microcrystalline Cellulose, Croscarmellose Sodium, Sodium Lauryl Sulfate, Hypromellose, Citric Acid Monohydrate, Propyl Gallate, Butylated Hydroxy Anisole and Magnesium Stearate.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Ezetimibe and simvastatin tablets

Plasma cholesterol is derived from intestinal absorption and endogenous synthesis. Ezetimibe and simvastatin tablets contains ezetimibe and simvastatin, two lipid-lowering compounds with complementary mechanisms of action.

Ezetimibe

Ezetimibe reduces blood cholesterol by inhibiting the absorption of cholesterol by the small intestine. The molecular target of ezetimibe has been shown to be the sterol transporter, Niemann-Pick C1-Like 1 (NPC1L1), which is involved in the intestinal uptake of cholesterol and phytosterols.

Ezetimibe localizes at the brush border of the small intestine and inhibits the absorption of cholesterol, leading to a decrease in the delivery of intestinal cholesterol to the liver. This causes a reduction of hepatic cholesterol stores and an increase in clearance of cholesterol from the blood.

Simvastatin

Simvastatin is a prodrug and is hydrolyzed to its active β -hydroxyacid form, simvastatin acid, after administration. Simvastatin acid and its metabolites are inhibitors of HMG-CoA reductase, the rate-limiting enzyme converts HMG-CoA to mevalonate, a precursor of cholesterol.

12.2 Pharmacodynamics

Ezetimibe and simvastatin tablets reduces total cholesterol (total-C), LDL-C, apolipoprotein (Apo) B, and non-high-density lipoprotein cholesterol (non-HDL-C) in patients with hyperlipidemia.

Ezetimibe

In a 2-week clinical trial in 18 hypercholesterolemic patients, ezetimibe inhibited intestinal cholesterol absorption by 54%, compared with placebo. Ezetimibe had no clinically meaningful effect on the plasma concentrations of the fat-soluble vitamins A, D, and E and did not impair adrenocortical steroid hormone production.

Simvastatin

Inhibition of HMG-CoA reductase by simvastatin acid accelerates the expression of LDL-receptors, followed by the uptake of LDL-C from blood to the liver, leading to a decrease in plasma LDL-C and total cholesterol. Sustained inhibition of cholesterol synthesis in the liver also decreases levels of very-low density lipoproteins. The maximum LDL-C reduction of ZOCOR is usually achieved by 4 weeks and is maintained after that.

12.3 Pharmacokinetics

The results of a bioequivalence trial in healthy subjects demonstrated that the ezetimibe and simvastatin tablets (ezetimibe and simvastatin) 10 mg/10 mg to 10 mg/80 mg combination tablets are bioequivalent to coadministration of corresponding doses of ezetimibe and simvastatin as individual tablets.

Absorption

Ezetimibe

After oral administration, ezetimibe is absorbed and extensively conjugated to a pharmacologically active phenolic glucuronide (ezetimibe-glucuronide). Mean maximum plasma concentrations (C_{max}) occur within 1 to 2 hours for ezetimibe-glucuronide and 4 to 12 hours for ezetimibe. The absolute bioavailability of ezetimibe cannot be determined as the compound is virtually insoluble in aqueous media suitable for injection.

Simvastatin

The availability of the β -hydroxyacid to the systemic circulation following an oral dose of simvastatin was found to be less than 5% of the dose, consistent with extensive hepatic first-pass extraction.

Effect of Food on Oral Absorption

Ezetimibe

Concomitant food administration (high-fat or non-fat meals) had no effect on the extent of absorption of ezetimibe when administered as 10-mg tablets. The C_{max} value of ezetimibe was increased by 38% with consumption of high-fat meals.

Simvastatin

Relative to the fasting state, the plasma profiles of both active and total inhibitors of HMG-CoA reductase were not affected when simvastatin was administered immediately before an American Heart Association recommended low-fat meal.

Distribution

Ezetimibe

Ezetimibe and ezetimibe-glucuronide are highly bound (>90%) to human plasma proteins.

Simvastatin

Both simvastatin and its β -hydroxyacid metabolite are highly bound (approximately 95%) to human plasma proteins. When radiolabeled simvastatin was administered to rats, simvastatin-derived radioactivity crossed the blood-brain barrier.

Elimination

Metabolism

Ezetimibe

Ezetimibe is primarily metabolized in the small intestine and liver via glucuronide conjugation with subsequent biliary and renal excretion. Minimal oxidative metabolism has been observed in all species evaluated.

In humans, ezetimibe is rapidly metabolized to ezetimibe-glucuronide. Ezetimibe and ezetimibe-glucuronide are the major drug-derived compounds detected in plasma, constituting approximately 10 to 20% and 80 to 90% of the total drug in plasma, respectively. Both ezetimibe and ezetimibe-glucuronide are eliminated from plasma with a half-life of approximately 22 hours for both ezetimibe and ezetimibe-glucuronide. Plasma concentration-time profiles exhibit multiple peaks, suggesting enterohepatic recycling.

Simvastatin

Simvastatin is a lactone that is readily hydrolyzed *in vivo* to the corresponding β -hydroxyacid, a potent inhibitor of HMG-CoA reductase. Inhibition of HMG-CoA reductase is a basis for an assay in pharmacokinetic studies of the β -hydroxyacid metabolites (active inhibitors) and, following base hydrolysis, active plus latent inhibitors (total inhibitors) in plasma following administration of simvastatin. The major active metabolites of simvastatin present in human plasma are the β -hydroxyacid of simvastatin and its 6'-hydroxy, 6'-hydroxymethyl, and 6'-exomethylene derivatives.

Excretion

Ezetimibe

Following oral administration of ^{14}C -ezetimibe (20 mg) to human subjects, total ezetimibe (ezetimibe + ezetimibe-glucuronide) accounted for approximately 93% of the total radioactivity in plasma. After 48 hours, there were no detectable levels of radioactivity in the plasma.

Approximately 78% and 11% of the administered radioactivity were recovered in the feces and urine,

respectively, over a 10-day collection period. Ezetimibe was the major component in feces and accounted for 69% of the administered dose, while ezetimibe-glucuronide was the major component in urine and accounted for 9% of the administered dose.

Simvastatin

Following an oral dose of ^{14}C -labeled simvastatin in man, 13% of the dose was excreted

in urine and 60% in feces. Plasma concentrations of total radioactivity (simvastatin plus ¹⁴C-metabolites) peaked at 4 hours and declined rapidly to about 10% of peak by 12 hours postdose.

Specific Populations

Geriatric Patients

Ezetimibe

In a multiple-dose trial with ezetimibe given 10 mg once daily for 10 days, plasma concentrations for total ezetimibe were about 2-fold higher in older (≥65 years) healthy subjects compared to younger subjects.

Simvastatin

In a trial including 16 geriatric patients between 70 and 78 years of age who received simvastatin 40 mg/day, the mean plasma level of total inhibitors activity was increased approximately 45% compared with 18 patients between 18 to 30 years of age. [See Use in Specific Populations (8.5).]

Gender

Ezetimibe

In a multiple-dose trial with ezetimibe given 10 mg once daily for 10 days, plasma concentrations for total ezetimibe were slightly higher (<20%) in females than in males.

Race

Ezetimibe

Based on a meta-analysis of multiple-dose pharmacokinetic studies, there were no pharmacokinetic differences between Black or African American and White subjects. Studies in Asian subjects indicated that the pharmacokinetics of ezetimibe was similar to those seen in White subjects.

Hepatic Impairment

Ezetimibe

After a single 10-mg dose of ezetimibe, the mean exposure (based on area under the curve [AUC]) to total ezetimibe was increased approximately 1.7-fold in patients with mild hepatic impairment (Child-Pugh score 5 to 6), compared to healthy subjects. The mean AUC values for total ezetimibe and ezetimibe increased approximately 3- to 4-fold and 5- to 6-fold, respectively, in patients with moderate (Child-Pugh score 7 to 9) or severe hepatic impairment (Child-Pugh score 10 to 15). In a 14-day, multiple-dose trial (10 mg daily) in patients with moderate hepatic impairment, the mean AUC for total ezetimibe and ezetimibe increased approximately 4-fold compared to healthy subjects.

Renal Impairment

Ezetimibe

After a single 10-mg dose of ezetimibe in patients with severe renal disease (n=8; mean CrCl \leq 30 mL/min/1.73 m²), the mean AUC for total ezetimibe and ezetimibe increased approximately 1.5-fold, compared to healthy subjects (n=9).

Simvastatin

Pharmacokinetic studies with another statin having a similar principal route of elimination to that of simvastatin have suggested that for a given dose level higher systemic exposure may be achieved in patients with severe renal impairment (as measured by creatinine clearance).

Drug Interactions [See also Drug Interactions (7).]

No clinically significant pharmacokinetic interaction was seen when ezetimibe was coadministered with simvastatin. No specific pharmacokinetic drug interaction studies with ezetimibe and simvastatin tablets have been conducted other than the following trial with NIASPAN (Niacin extended-release tablets).

Niacin: The effect of ezetimibe and simvastatin tablets (10/20 mg daily for 7 days) on the pharmacokinetics of NIASPAN extended-release tablets (1,000 mg for 2 days and 2,000 mg for 5 days following a low-fat breakfast) was studied in healthy subjects. The mean C_{max} and AUC of niacin increased 9% and 22%, respectively. The mean C_{max} and AUC of nicotinuric acid increased 10% and 19%, respectively (N=13). In the same trial, the effect of NIASPAN on the pharmacokinetics of ezetimibe and simvastatin tablets were evaluated (N=15). While concomitant NIASPAN decreased the mean C_{max} of total ezetimibe (1%), and simvastatin (2%), it increased the mean C_{max} of simvastatin acid (18%). In addition, concomitant NIASPAN increased the mean AUC of total ezetimibe (26%), simvastatin (20%), and simvastatin acid (35%).

Cases of myopathy/rhabdomyolysis have been observed with simvastatin coadministered with lipid-modifying doses (\geq 1 g/day niacin) of niacin-containing products. [*See Warnings and Precautions (5.1) and Drug Interactions (7.1).*]

Cytochrome P450: Ezetimibe had no significant effect on a series of probe drugs (caffeine, dextromethorphan, tolbutamide, and IV midazolam) known to be metabolized by cytochrome P450 (1A2, 2D6, 2C8/9 and 3A4) in a “cocktail” trial of twelve healthy adult males. This indicates that ezetimibe is neither an inhibitor nor an inducer of these cytochrome P450 isozymes, and it is unlikely that ezetimibe will affect the metabolism of drugs that are metabolized by these enzymes.

In a trial of 12 healthy volunteers, simvastatin at the 80-mg dose had no effect on the metabolism of the probe cytochrome P450 isoform 3A4 (CYP3A4) substrates midazolam and erythromycin. This indicates that simvastatin is not an inhibitor of CYP3A4 and, therefore, is not expected to affect the plasma levels of other drugs metabolized by CYP3A4.

Simvastatin acid is a substrate of the transport protein OATP1B1. Concomitant administration of medicinal products that are inhibitors of the transport protein OATP1B1 may lead to increased plasma concentrations of simvastatin acid and an

increased risk of myopathy. For example, cyclosporine has been shown to increase the AUC of statins; although the mechanism is not fully understood, the increase in AUC for simvastatin acid is presumably due, in part, to inhibition of CYP3A4 and/or OATP1B1[see *Drug Interactions (7)*].

Simvastatin is a substrate for CYP3A4. Inhibitors of CYP3A4 can raise the plasma levels of HMG-CoA reductase inhibitory activity and increase the risk of myopathy. [See *Warnings and Precautions (5.1)* and *Drug Interactions (7.1)*.]

Ezetimibe

Table 5 displays the effect of coadministered drugs on total ezetimibe.

Table 5: Effect of Coadministered Drugs on Total Ezetimibe

Coadministered Drug and Dosing Regimen	Total Ezetimibe*	
	Change in AUC	Change in C _{max}
Cyclosporine-stable dose required (75 to 150 mg BID) ^{†, ‡}	↑ 240%	↑ 290%
Fenofibrate, 200 mg QD, 14 days [‡]	↑ 48%	↑ 64%
Gemfibrozil, 600 mg BID, 7 days [‡]	↑ 64%	↑ 91%
Cholestyramine, 4 g BID, 14 days [‡]	↓ 55%	↓ 4%
Aluminum & magnesium hydroxide combination antacid, single dose [§]	↓ 4%	↓ 30%
Cimetidine, 400 mg BID, 7 days	↑ 6%	↑ 22%
Glipizide, 10 mg, single dose	↑ 4%	↓ 8%
Statins		
Lovastatin 20 mg QD, 7 days	↑ 9%	↑ 3%
Pravastatin 20 mg QD, 14 days	↑ 7%	↑ 23%
Atorvastatin 10 mg QD, 14 days	↓ 2%	↑ 12%
Rosuvastatin 10 mg QD, 14 days	↑ 13%	↑ 18%
Fluvastatin 20 mg QD, 14 days	↓ 19%	↑ 7%

* Based on 10 mg-dose of ezetimibe.

[†] Post-renal transplant patients with mild impaired or normal renal function. In a different trial, a renal transplant patient with severe renal insufficiency (creatinine clearance of 13.2 mL/min/1.73 m²) who was receiving multiple medications, including cyclosporine, demonstrated a 12-fold greater exposure to total ezetimibe compared to healthy subjects.

[‡] See 7. Drug Interactions.

[§] Supralox, 20 mL.

Table 6 displays the effects of ezetimibe coadministration on systemic exposure to other drugs.

Table 6: Effect of Ezetimibe Coadministration on Systemic Exposure to Other Drugs

Coadministered Drug and its Dosage Regimen	Ezetimibe Dosage Regimen	Change in AUC of Coadministered Drug	Change in C_{max} of Coadministered Drug
Warfarin, 25 mg single dose on Day 7	10 mg QD, 11 days	↓2% (R-warfarin) ↓4% (S-warfarin)	↑3% (R-warfarin) ↑1% (S-warfarin)
Digoxin, 0.5 mg single dose	10 mg QD, 8 days	↑2%	↓7%
Gemfibrozil, 600 mg BID, 7 days*	10 mg QD, 7 days	↓1%	↓11%
Ethinyl estradiol & Levonorgestrel, QD, 21 days	10 mg QD, Days 8 to 14 of 21 day oral contraceptive cycle	Ethinyl estradiol 0% Levonorgestrel 0%	Ethinyl estradiol ↓9% Levonorgestrel ↓5%
Glipizide, 10 mg on Days 1 and 9	10 mg QD, Days 2 to 9	↓3%	↓5%
Fenofibrate, 200 mg QD, 14 days*	10 mg QD, 14 days	↑11%	↑7%
Cyclosporine, 100 mg single dose Day 7*	20 mg QD, 8 days	↑15%	↑10%
Statins			
Lovastatin 20 mg QD, 7 days	10 mg QD, 7 days	↑19%	↑3%
Pravastatin 20 mg QD, 14 days	10 mg QD, 14 days	↓20%	↓24%
Atorvastatin 10 mg QD, 14 days	10 mg QD, 14 days	↓4%	↑7%
Rosuvastatin 10 mg QD, 14 days	10 mg QD, 14 days	↑19%	↑17%
Fluvastatin 20 mg QD, 14 days	10 mg QD, 14 days	↓39%	↓27%

* See 7. Drug Interactions.

Simvastatin

Table 7 displays the effects of coadministration drugs or grapefruit juice on simvastatin systemic exposure [see *Drug Interactions (7)*].

Table 7: Effect of Coadministered Drugs or Grapefruit Juice on Simvastatin Systemic Exposure

Coadministered Drug or Grapefruit Juice	Dosing of Coadministered Drug or Grapefruit Juice	Dosing of Simvastatin	Geometric Mean Ratio (Ratio* with / without coadministered drug) No Effect = 1.00		
				AUC	C_{max}
Telithromycin [†]	200 mg QD for 4 days	80 mg	simvastatin acid [‡]	12 8.9	15 5.3

			simvastatin		
Nelfinavir †	1250 mg BID for 14 days	20 mg QD for 28 days	simvastatin acid‡		
			simvastatin	6	6.2
Itraconazole†	200 mg QD for 4 days	80 mg	simvastatin acid‡		13.1
			simvastatin		13.1
Posaconazole	100 mg (oral suspension) QD for 13 days	40 mg	simvastatin acid‡	7.3	9.2
			simvastatin	10.3	9.4
	200 mg (oral suspension) QD for 13 days	40 mg	simvastatin acid‡	8.5	9.5
			simvastatin	10.6	11.4
Gemfibrozil	600 mg BID for 3 days	40 mg	simvastatin acid‡	2.85	2.18
			simvastatin	1.35	0.91
Grapefruit Juice§ (high dose)	200 mL of double-strength TID¶	60 mg single dose	simvastatin acid	7	
			simvastatin	16	
Grapefruit Juice§ (low dose)	8 oz (about 237 mL) of single-strength#	20 mg single dose	simvastatin acid	1.3	
			simvastatin	1.9	
Verapamil SR	240 mg QD Days 1 to 7 then 240 mg BID on Days 8 to 10	80 mg on Day 10	simvastatin acid	2.3	2.4
			simvastatin	2.5	2.1
Diltiazem	120 mg BID for 10 days	80 mg on Day 10	simvastatin acid	2.69	2.69
			simvastatin	3.10	2.88
Diltiazem	120 mg BID for 14 days	20 mg on Days 14	simvastatin	4.6	3.6
Dronedarone	400 mg BID for 14 days	40 mg QD for 14 days	simvastatin acid	1.96	2.14
			simvastatin	3.90	3.75
Amiodarone	400 mg QD for 3 days	40 mg on Day 3	simvastatin acid	1.75	1.72
			simvastatin	1.76	1.79
Amlodipine	10 mg QD for 10 days	80 mg on Days	simvastatin	1.58	1.56

	days	10	acid	1.77	1.47
			simvastatin		
Ranolazine SR	1000 mg BID for 7 days	80 mg on Day 1 and Days 6 to 9	simvastatin acid	2.26	2.28
			simvastatin acid	1.86	1.75
			simvastatin acid		
Lomitapide	60 mg QD for 7 days	40 mg single dose	simvastatin acid	1.7	1.6
			simvastatin acid	2	2
			simvastatin acid		
Lomitapide	10 mg QD for 7 days	20 mg single dose	simvastatin acid	1.4	1.4
			simvastatin acid	1.6	1.7
			simvastatin acid		
Fenofibrate	160 mg QD for 14 days	80 mg QD on Days 8 to 14	simvastatin acid	0.64	0.89
			simvastatin acid	0.89	0.83
			simvastatin acid		
Propranolol	80 mg single dose	80 mg single dose	total inhibitor	0.79	↓ from 33.6 to 21.1 ng·eq/mL
			active inhibitor	0.79	↓ from 7.0 to 4.7 ng·eq/mL

* Results based on a chemical assay except results with propranolol as indicated.

† Results could be representative of the following CYP3A4 inhibitors: ketoconazole, erythromycin, clarithromycin, HIV protease inhibitors, and nefazodone.

‡ Simvastatin acid refers to the β -hydroxyacid of simvastatin.

§ The effect of amounts of grapefruit juice between those used in these two studies on simvastatin pharmacokinetics has not been studied.

¶ Double-strength: one can of frozen concentrate diluted with one can of water. Grapefruit juice was administered TID for 2 days, and 200 mL together with single dose simvastatin and 30 and 90 minutes following single dose simvastatin on Day 3.

Single-strength: one can of frozen concentrate diluted with 3 cans of water. Grapefruit juice was administered with breakfast for 3 days, and simvastatin was administered in the evening on Day 3.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Ezetimibe and Simvastatin Tablets

No animal carcinogenicity or fertility studies have been conducted with the combination of ezetimibe and simvastatin. The combination of ezetimibe with simvastatin did not show evidence of mutagenicity *in vitro* in a microbial mutagenicity (Ames) test with *Salmonella typhimurium* and *Escherichia coli* with or without metabolic activation. No evidence of clastogenicity was observed *in vitro* in a chromosomal aberration assay in human peripheral blood lymphocytes with ezetimibe and simvastatin with or without metabolic activation. There was no evidence of genotoxicity at doses up to 600 mg/kg with the combination of ezetimibe and simvastatin (1:1) in the *in vivo* mouse micronucleus test.

Ezetimibe

A 104-week dietary carcinogenicity trial with ezetimibe was conducted in rats at doses up to 1,500 mg/kg/day (males) and 500 mg/kg/day (females) (~20 times the human exposure at 10 mg daily based on AUC_{0-24hr} for total ezetimibe). A 104-week dietary carcinogenicity trial with ezetimibe was also conducted in mice at doses up to 500 mg/kg/day (>150 times the human exposure at 10 mg daily based on AUC_{0-24hr} for total ezetimibe). There were no statistically significant increases in tumor incidences in drug-treated rats or mice.

No evidence of mutagenicity was observed *in vitro* in a microbial mutagenicity (Ames) test with *Salmonella typhimurium* and *Escherichia coli* with or without metabolic activation. No evidence of clastogenicity was observed *in vitro* in a chromosomal aberration assay in human peripheral blood lymphocytes with or without metabolic activation. In addition, there was no evidence of genotoxicity in the *in vivo* mouse micronucleus test.

In oral (gavage) fertility studies of ezetimibe conducted in rats, there was no evidence of reproductive toxicity at doses up to 1,000 mg/kg/day in male or female rats (~7 times the human exposure at 10 mg daily based on AUC_{0-24hr} for total ezetimibe).

Simvastatin

In a 72-week carcinogenicity trial, mice were administered daily doses of simvastatin of 25, 100, and 400 mg/kg body weight, which resulted in mean plasma drug levels approximately 1, 4, and 8 times higher than the mean human plasma drug level, respectively, (as total inhibitory activity based on AUC) after an 80-mg oral dose. Liver carcinomas were significantly increased in high-dose females and mid- and high- dose males with a maximum incidence of 90% in males. The incidence of adenomas of the liver was significantly increased in mid- and high-dose females. Drug treatment also significantly increased the incidence of lung adenomas in mid- and high-dose males and females. Adenomas of the Harderian gland (a gland of the eye of rodents) were significantly higher in high-dose mice than in controls. No evidence of a tumorigenic effect was observed at 25 mg/kg/day.

In a separate 92-week carcinogenicity trial in mice at doses up to 25 mg/kg/day, no evidence of a tumorigenic effect was observed (mean plasma drug levels were 1 times higher than humans given 80 mg simvastatin as measured by AUC).

In a two-year trial in rats at 25 mg/kg/day, there was a statistically significant increase in the incidence of thyroid follicular adenomas in female rats exposed to approximately 11

times higher levels of simvastatin than in humans given 80 mg simvastatin (as measured by AUC).

A second two-year rat carcinogenicity trial with doses of 50 and 100 mg/kg/day produced hepatocellular adenomas and carcinomas (in female rats at both doses and in males at 100 mg/kg/day). Thyroid follicular cell adenomas were increased in males and females at both doses; thyroid follicular cell carcinomas were increased in females at 100 mg/kg/day. The increased incidence of thyroid neoplasms appears to be consistent with findings from other statins. These treatment levels represented plasma drug levels (AUC) of approximately 7 and 15 times (males) and 22 and 25 times (females) the mean human plasma drug exposure after an 80-mg daily dose.

No evidence of mutagenicity was observed in a microbial mutagenicity (Ames) test with or without rat or mouse liver metabolic activation. In addition, no evidence of damage to genetic material was noted in an *in vitro* alkaline elution assay using rat hepatocytes, a V-79 mammalian cell forward mutation trial, an *in vitro* chromosome aberration trial in CHO cells, or an *in vivo* chromosomal aberration assay in mouse bone marrow.

There was decreased fertility in male rats treated with simvastatin for 34 weeks at 25 mg/kg body weight (4 times the maximum human exposure level, based on AUC, in patients receiving 80 mg/day); however, this effect was not observed during a subsequent fertility trial in which simvastatin was administered at this same dose level to male rats for 11 weeks (the entire cycle of spermatogenesis including epididymal maturation). No microscopic changes were observed in the testes of rats from either trial. At 180 mg/kg/day (which produces exposure levels 22 times higher than those in humans taking 80 mg/day based on surface area, mg/m²), seminiferous tubule degeneration (necrosis and loss of spermatogenic epithelium) was observed. In dogs, there was drug-related testicular atrophy, decreased spermatogenesis, spermatocytic degeneration and giant cell formation at 10 mg/kg/day (approximately 2 times the human exposure, based on AUC, at 80 mg/day). The clinical significance of these findings is unclear.

14 CLINICAL STUDIES

Primary Hyperlipidemia in Adults

Ezetimibe and simvastatin tablets

Ezetimibe and simvastatin tablets reduces LDL-C in adult patients with primary hyperlipidemia. Maximal to near maximal response is generally achieved within 2 weeks and maintained during chronic therapy.

Ezetimibe and simvastatin tablets is effective in males and females with primary hyperlipidemia. There were insufficient numbers of patients who self-identified as Black or African American, Asian, or other races to determine if these patients responded differently than White patients.

Five multicenter, double-blind trials conducted with either ezetimibe and simvastatin tablets or coadministered ezetimibe and simvastatin equivalent to ezetimibe and simvastatin tablets in patients with primary hyperlipidemia are reported: two were comparisons with simvastatin, two were comparisons with atorvastatin, and one was a comparison with rosuvastatin.

In a multicenter, double-blind, placebo-controlled, 12-week trial, 1,528 patients with primary hyperlipidemia were randomized to one of ten treatment groups: placebo, ezetimibe (10 mg), simvastatin (10 mg, 20 mg, 40 mg, or 80 mg), or ezetimibe and simvastatin tablets (10/10, 10/20, 10/40, or 10/80).

When patients receiving ezetimibe and simvastatin tablets were compared to those receiving all doses of simvastatin, ezetimibe and simvastatin tablets significantly lowered total-C, LDL-C, Apo B, TG, and non-HDL-C. The effects of ezetimibe and simvastatin tablets on HDL-C were similar to the effects seen with simvastatin. Further analysis showed ezetimibe and simvastatin tablets significantly increased HDL-C compared with placebo. (See Table 8.) The lipid response to ezetimibe and simvastatin tablets was similar in patients with TG levels greater than or less than 200 mg/dL.

Table 8: Response to Ezetimibe and Simvastatin Tablets in Patients with Primary Hyperlipidemia (Mean* % Change from Untreated Baseline†)

Treatment (Daily Dose)	N	Total-C	LDL-C	Apo B	HDL-C	TG*	Non-HDL-C
Pooled data (All Ezetimibe and Simvastatin Tablets doses)‡	609	-38	-53	-42	+7	-24	-49
Pooled data (All simvastatin doses)‡	622	-28	-39	-32	+7	-21	-36
Ezetimibe 10 mg	149	-13	-19	-15	+5	-11	-18
Placebo	148	-1	-2	0	0	-2	-2
Ezetimibe and Simvastatin Tablets by dose							
10/10	152	-31	-45	-35	+8	-23	-41
10/20	156	-36	-52	-41	+10	-24	-47
10/40	147	-39	-55	-44	+6	-23	-51
10/80	154	-43	-60	-49	+6	-31	-56
Simvastatin by dose							
10 mg	158	-23	-33	-26	+5	-17	-30
20 mg	150	-24	-34	-28	+7	-18	-32
40 mg	156	-29	-41	-33	+8	-21	-38
80 mg	158	-35	-49	-39	+7	-27	-45

* For triglycerides, median % change from baseline.

† Baseline - on no lipid-lowering drug.

‡Ezetimibe and simvastatin tablets doses pooled (10/10 to 10/80) significantly reduced total-C, LDL-C, Apo B, TG, and non-HDL-C compared to simvastatin and significantly increased HDL-C compared to placebo.

In a multicenter, double-blind, controlled, 23-week trial, 710 patients with known CHD or

CHD risk equivalents, as defined by the NCEP ATP III guidelines, and an LDL-C \geq 130 mg/dL were randomized to one of four treatment groups: coadministered ezetimibe and simvastatin equivalent to ezetimibe and simvastatin tablets (10/10, 10/20, and 10/40) or simvastatin 20 mg. Patients not reaching an LDL-C $<$ 100 mg/dL had their simvastatin dose titrated at 6-week intervals to a maximal dose of 80 mg.

At Week 5, the LDL-C reductions with ezetimibe and simvastatin tablets 10/10, 10/20, or 10/40 were significantly larger than with simvastatin 20 mg (see Table 9).

Table 9: Response to Ezetimibe and Simvastatin Tablets after 5 Weeks in Patients with CHD or CHD Risk Equivalents and an LDL-C \geq 130 mg/dL

	Simvastatin 20 mg	Ezetimibe and Simvastatin Tablets 10/10	Ezetimibe and Simvastatin Tablets 10/20	Ezetimibe and Simvastatin Tablets 10/40
N	253	251	109	97
Mean baseline LDL-C	174	165	167	171
Percent change LDL-C	-38	-47	-53	-59

In a multicenter, double-blind, 6-week trial, 1902 patients with primary hyperlipidemia were randomized to one of eight treatment groups: Ezetimibe and simvastatin tablets (10/10, 10/20, 10/40, or 10/80) or atorvastatin (10 mg, 20 mg, 40 mg, or 80 mg).

Across the dosage range, when patients receiving ezetimibe and simvastatin tablets were compared to those receiving milligram-equivalent statin doses of atorvastatin, ezetimibe and simvastatin tablets lowered total-C, LDL-C, Apo B, and non-HDL-C significantly more than atorvastatin. Only the 10/40 mg and 10/80 mg ezetimibe and simvastatin tablets doses increased HDL-C significantly more than the corresponding milligram-equivalent statin dose of atorvastatin. The effects of ezetimibe and simvastatin tablets on TG were similar to the effects seen with atorvastatin. (See Table 10.)

Table 10: Response to Ezetimibe and Simvastatin Tablets and Atorvastatin in Patients with Primary Hyperlipidemia

(Mean* % Change from Untreated Baseline[†])

Treatment							
(Daily Dose)	N	Total-C	LDL-C[‡]	Apo B[‡]	HDL-C	TG*	Non-HDL-C[‡]
Ezetimibe and Simvastatin Tablets by dose							
10/10	230	-34 [§]	-47 [§]	-37 [§]	+8	-26	-43 [§]
10/20	233	-37 [§]	-51 [§]	-40 [§]	+7	-25	-46 [§]
10/40	236	-41 [§]	-57 [§]	-46 [§]	+9 [§]	-27	-52 [§]
10/80	224	-43 [§]	-59 [§]	-48 [§]	+8 [§]	-31	-54 [§]
Atorvastatin by dose							

10 mg	235	-27	-36	-31	+7	-21	-34
20 mg	230	-32	-44	-37	+5	-25	-41
40 mg	232	-36	-48	-40	+4	-24	-45
80 mg	230	-40	-53	-44	+1	-32	-50

* For triglycerides, median % change from baseline.

† Baseline - on no lipid-lowering drug.

‡Ezetimibe and simvastatin tablets doses pooled (10/10 to 10/80)

provided significantly greater reductions in total-C, LDL-C, Apo B, and non-HDL-C compared to atorvastatin doses pooled (10 to 80).

§ p<0.05 for difference with atorvastatin at equal mg doses of the simvastatin component.

In a multicenter, double-blind, 24-week, forced-titration trial, 788 patients with primary hyperlipidemia were randomized to receive coadministered ezetimibe and simvastatin equivalent to ezetimibe and simvastatin tablets (10/10 and 10/20) or atorvastatin 10 mg. For all three treatment groups, the dose of the statin was titrated at 6-week intervals to 80 mg. At each pre-specified dose comparison, ezetimibe and simvastatin tablets lowered LDL-C to a greater degree than atorvastatin (see Table 11).

Table 11: Response to Ezetimibe and Simvastatin Tablets and Atorvastatin in Patients with Primary Hyperlipidemia (Mean* % Change from Untreated Baseline†)

Treatment	N	Total-C	LDL-C	Apo B	HDL-C	CTG*	Non-HDL-C
Week 6							
Atorvastatin 10 mg†	262	-28	-37	-32	+5	-23	-35
Ezetimibe and Simvastatin Tablets 10/10§	263	-34¶	-46¶	-38¶	+8¶	-26	-43¶
Ezetimibe and Simvastatin Tablets 10/20#	263	-36¶	-50¶	-41¶	+10¶	-25	-46¶
Week 12							
Atorvastatin 20 mg	246	-33	-44	-38	+7	-28	-42
Ezetimibe and Simvastatin Tablets 10/20	250	-37¶	-50¶	-41¶	+9	-28	-46¶
Ezetimibe and Simvastatin Tablets 10/40	252	-39¶	-54¶	-45¶	+12¶	-31	-50¶
Week 18							
Atorvastatin 40 mg	237	-37	-49	-42	+8	-31	-47
Ezetimibe and Simvastatin Tablets 10/40 ^p	482	-40¶	-56¶	-45¶	+11¶	-32	-52¶
Week 24							
Atorvastatin 80 mg	228	-40	-53	-45	+6	-35	-50
Ezetimibe and Simvastatin	450	-42¶	-50¶	-40¶	+12¶	-35	-55¶

Tablets 10/80 ^P	+33	-43	-33	-43	-12	-33	-33
----------------------------	-----	-----	-----	-----	-----	-----	-----

* For triglycerides, median % change from baseline.

† Baseline - on no lipid-lowering drug.

‡ Atorvastatin: 10 mg start dose titrated to 20 mg, 40 mg, and 80 mg through Weeks 6, 12, 18, and 24.

§Ezetimibe and simvastatin tablets: 10/10 start dose titrated to 10/20, 10/40, and 10/80 through Weeks 6, 12, 18, and 24.

¶ $p \leq 0.05$ for difference with atorvastatin in the specified week.

Ezetimibe and simvastatin tablets: 10/20 start dose titrated to 10/40, 10/40, and 10/80 through Weeks 6, 12, 18, and 24.

^P Data pooled for common doses of ezetimibe and simvastatin tablets at Weeks 18 and 24.

In a multicenter, double-blind, 6-week trial, 2,959 patients with primary hyperlipidemia, were randomized to one of six treatment groups: ezetimibe and simvastatin tablets (10/20, 10/40, or 10/80) or rosuvastatin (10 mg, 20 mg, or 40 mg).

The effects of ezetimibe and simvastatin tablets and rosuvastatin on total-C, LDL-C, Apo B, TG, non-HDL-C and HDL-C are shown in Table 12.

Table 12: Response to Ezetimibe and Simvastatin Tablets and Rosuvastatin in Patients with Primary Hyperlipidemia (Mean*% Change from Untreated Baseline†)

Treatment							
(Daily Dose)	N	Total-C ‡	LDL-C‡	Apo B‡	HDL-C	TG*	Non-HDL-C‡
Ezetimibe and Simvastatin Tablets by dose							
10/20	476	-37 [§]	-52 [§]	-42 [§]	+7	-23 [§]	-47 [§]
10/40	477	-39 [¶]	-55 [¶]	-44 [¶]	+8	-27	-50 [¶]
10/80	474	-44 [#]	-61 [#]	-50 [#]	+8	-30 [#]	-56 [#]
Rosuvastatin by dose							
10 mg	475	-32	-46	-37	+7	-20	-42
20 mg	478	-37	-52	-43	+8	-26	-48
40 mg	475	-41	-57	-47	+8	-28	-52

* For triglycerides, median % change from baseline.

† Baseline - on no lipid-lowering drug.

‡Ezetimibe and simvastatin tablets doses pooled (10/20 to 10/80) provided significantly greater reductions in total-C, LDL-C, Apo B, and non-HDL-C compared to rosuvastatin doses pooled (10 to 40 mg).

§ $p < 0.05$ vs. rosuvastatin 10 mg.

¶ $p < 0.05$ vs. rosuvastatin 20 mg.

$p < 0.05$ vs. rosuvastatin 40 mg.

In a multicenter, double-blind, 24-week trial, 214 patients with type 2 diabetes mellitus treated with thiazolidinediones (rosiglitazone or pioglitazone) for a minimum of 3 months and simvastatin 20 mg for a minimum of 6 weeks were randomized to receive either simvastatin 40 mg or the coadministered active ingredients equivalent to ezetimibe and simvastatin tablets 10/20. The median LDL-C and HbA1c levels at baseline were 89 mg/dL and 7.1%, respectively.

Ezetimibe and simvastatin tablets 10/20 was significantly more effective than doubling the dose of simvastatin to 40 mg. The median percent changes from baseline for ezetimibe and simvastatin tablets vs. simvastatin were: LDL-C -25% and -5%; total-C -16% and -5%; Apo B -19% and -5%; and non-HDL-C -23% and -5%. Results for HDL-C and TG between the two treatment groups were not significantly different.

Ezetimibe

In two multicenter, double-blind, placebo-controlled, 12-week trials in 1719 patients with primary hyperlipidemia, ezetimibe significantly lowered total-C (-13%), LDL-C (-19%), Apo B (-14%), and TG (-8%), and increased HDL-C (+3%) compared to placebo. Reduction in LDL-C was consistent across age, sex, and baseline LDL-C.

Simvastatin

In two large, placebo-controlled clinical trials, the Scandinavian Simvastatin Survival Trial (N=4,444 patients) and the Heart Protection Trial (N=20,536 patients), the effects of treatment with simvastatin were assessed in patients at high risk of coronary events because of existing coronary heart disease, diabetes, peripheral vessel disease, history of stroke or other cerebrovascular disease. Simvastatin was proven to reduce: the risk of total mortality by reducing CHD deaths; the risk of non-fatal myocardial infarction and stroke; and the need for coronary and non-coronary revascularization procedures.

No incremental benefit of ezetimibe and simvastatin tablets on cardiovascular morbidity and mortality over and above that demonstrated for simvastatin has been established.

Heterozygous Familial Hypercholesterolemia (HeFH) in Pediatric Patients

The effects of ezetimibe coadministered with simvastatin (n=126) compared to simvastatin monotherapy (n=122) have been evaluated in males and females with HeFH. In a multicenter, double-blind, controlled trial followed by an open-label phase, 142 males and 106 postmenarchal females, 10 to 17 years of age (mean age 14.2 years, 43% females, 82% White, 4% Asian, 2% Black or African American, 13% multi-racial; 14% identified as Hispanic or Latino ethnicity) with HeFH were randomized to receive either ezetimibe coadministered with simvastatin or simvastatin monotherapy. Inclusion in the trial required 1) a baseline LDL-C level between 160 and 400 mg/dL and 2) a medical history and clinical presentation consistent with HeFH. The mean baseline LDL-C value was 225 mg/dL (range: 161 to 351 mg/dL) in the ezetimibe coadministered with simvastatin group compared to 219 mg/dL (range: 149 to 336 mg/dL) in the simvastatin monotherapy group. The patients received coadministered ezetimibe and simvastatin (10 mg, 20 mg, or 40 mg) or simvastatin monotherapy (10 mg, 20 mg, or 40 mg) for 6 weeks, coadministered ezetimibe and 40-mg simvastatin or 40-mg simvastatin

monotherapy for the next 27 weeks, and open-label coadministered ezetimibe and simvastatin (10 mg, 20 mg, or 40 mg) for 20 weeks thereafter.

The results of the trial at Week 6 are summarized in Table 13. Results at Week 33 were consistent with those at Week 6.

Table 13: Mean Percent Difference at Week 6 Between the Pooled Ezetimibe Coadministered with Simvastatin Group and the Pooled Simvastatin Monotherapy Group in Adolescent Patients with HeFH

	Total-C	LDL-C	Apo B	Non-HDL-C
Mean percent difference between treatment groups	-12%	-15%	-12%	-14%
95% Confidence Interval	(-15%,-9%)	(-18%,-12%)	(-15%,-9%)	(-17%,-11%)

Homozygous Familial Hypercholesterolemia (HoFH) in Adults

A double-blind, randomized, 12-week trial was performed in patients with a clinical and/or genotypic diagnosis of HoFH. Data were analyzed from a subgroup of patients (n=14) receiving simvastatin 40 mg at baseline. Increasing the dose of simvastatin from 40 to 80 mg (n=5) produced a reduction of LDL-C of 13% from baseline on simvastatin 40 mg. Coadministered ezetimibe and simvastatin equivalent to ezetimibe and simvastatin tablets (10/40 and 10/80 pooled, n=9), produced a reduction of LDL-C of 23% from baseline on simvastatin 40 mg. In those patients coadministered ezetimibe and simvastatin equivalent to ezetimibe and simvastatin tablets (10/80, n=5), a reduction of LDL-C of 29% from baseline on simvastatin 40 mg was produced.

Chronic Kidney Disease (CKD) in Adults

The Trial of Heart and Renal Protection (SHARP) was a multinational, randomized, placebo- controlled, double-blind trial that investigated the effect of ezetimibe and simvastatin tablets on the time to a first major vascular event (MVE) among 9,438 patients with moderate to severe chronic kidney disease (approximately one- third on dialysis at baseline) who did not have a history of myocardial infarction or coronary revascularization. An MVE was defined as nonfatal MI, cardiac death, stroke, or any revascularization procedure. Patients were allocated to treatment using a method that took into account the distribution of 8 important baseline characteristics of patients already enrolled and minimized the imbalance of those characteristics across the groups.

For the first year, 9,438 patients were allocated 4:4:1, to ezetimibe and simvastatin tablets 10/20, placebo, or simvastatin 20 mg daily, respectively. The 1-year simvastatin arm enabled the comparison of ezetimibe and simvastatin tablets to simvastatin with

regard to safety and effect on lipid levels. At 1 year the simvastatin-only arm was re-allocated 1:1 to ezetimibe and simvastatin tablets 10/20 or placebo. A total of 9,270 patients were ever allocated to ezetimibe and simvastatin tablets 10/20 (n=4650) or placebo (n=4620) during the trial. The median follow-up duration was 4.9 years. Patients had a mean age of 61 years; 63% were male, 72% were White, and 23% were diabetic; and, for those not on dialysis at baseline, the median serum creatinine was 2.5 mg/dL and the median estimated glomerular filtration rate (eGFR) was 25.6 mL/min/1.73 m², with 94% of patients having an eGFR < 45 mL/min/1.73 m². Eligibility did not depend on lipid levels. Mean LDL-C at baseline was 108 mg/dL. At 1 year, the mean LDL-C was 26% lower in the simvastatin arm and 38% lower in the ezetimibe and simvastatin tablets arm relative to placebo. At the midpoint of the trial (2.5 years), the mean LDL-C was 32% lower for ezetimibe and simvastatin tablets relative to placebo. Patients no longer taking trial medication were included in all lipid measurements.

In the primary intent-to-treat analysis, 639 (15.2%) of 4,193 patients initially allocated to ezetimibe and simvastatin tablets and 749 (17.9%) of 4,191 patients initially allocated to placebo experienced an MVE. This corresponded to a relative risk reduction of 16% (p=0.001) (see Figure 1). Similarly, 526 (11.3%) of 4650 patients ever allocated to ezetimibe and simvastatin tablets and 619 (13.4%) of 4,620 patients ever allocated to placebo experienced a major atherosclerotic event (MAE; a subset of the MVE composite that excluded non-coronary cardiac deaths and hemorrhagic stroke), corresponding to a relative risk reduction of 17% (p=0.002). The trial demonstrated that treatment with ezetimibe and simvastatin tablets 10/20 mg versus placebo reduced the risk for MVE and MAE in this CKD population. The trial design precluded drawing conclusions regarding the independent contribution of either ezetimibe or simvastatin to the observed effect.

The treatment effect of ezetimibe and simvastatin tablets on MVE was attenuated among patients on dialysis at baseline compared with those not on dialysis at baseline. Among 3023 patients on dialysis at baseline, ezetimibe and simvastatin tablets reduced the risk of MVE by 6% (RR 0.94: 95% CI 0.80 to 1.09) compared with 22% (RR 0.78: 95% CI 0.69 to 0.89) among 6247 patients not on dialysis at baseline (interaction P=0.08).

Figure 1: Effect of Ezetimibe and Simvastatin Tablets on the Primary Endpoint of Risk of Major Vascular Events

The individual components of MVE in all patients ever allocated to ezetimibe and simvastatin tablets or placebo are presented in Table 14.

Table 14: Number of First Events for Each Component of the Major Vascular Event Composite Endpoint in SHARP*

Outcome	Ezetimibe and Simvastatin Tablets	Placebo (N=4620)	Risk Ratio (95% CI)	P-value

	10/20 (N=4650)			
Major Vascular Events	701 (15.1%)	814 (17.6%)	0.85 (0.77 to 0.94)	0.001
Nonfatal MI	134 (2.9%)	159 (3.4%)	0.84 (0.66 to 1.05)	0.12
Cardiac Death	253 (5.4%)	272 (5.9%)	0.93 (0.78 to 1.10)	0.38
Any Stroke	171 (3.7%)	210 (4.5%)	0.81 (0.66 to 0.99)	0.038
Non-hemorrhagic Stroke	131 (2.8%)	174 (3.8%)	0.75 (0.60 to 0.94)	0.011
Hemorrhagic Stroke	45 (1.0%)	37 (0.8%)	1.21 (0.78 to 1.86)	0.40
Any Revascularization	284 (6.1%)	352 (7.6%)	0.79 (0.68 to 0.93)	0.004

*Intention-to-treat analysis on all SHARP patients ever allocated to ezetimibe and simvastatin tablets or placebo.

Among patients not on dialysis at baseline, ezetimibe and simvastatin tablets did not reduce the risk of progressing to end-stage renal disease compared with placebo (RR 0.97: 95% CI 0.89 to 1.05).

Simvastatin Cardiovascular Outcome Trials in Adults at High Risk of Coronary Heart Disease Events

In a randomized, double-blind, placebo-controlled, multi-centered trial [the Scandinavian Simvastatin Survival Trial (Trial 4S)], the effect of therapy with simvastatin on total mortality was assessed in 4,444 adult patients with CHD (history of angina and/or a previous myocardial infarction) and baseline total cholesterol (total-C) between 212 and 309 mg/dL who were on a lipid-lowering diet. In Trial 4S, patients were treated with standard care, including lipid-lowering diet, and randomized to either simvastatin 20-40 mg/day (n=2,221) or placebo (n=2,223) for a median duration of 5.4 years.

- Simvastatin significantly reduced the risk of mortality by 30% (p=0.0003, 182 deaths in the simvastatin group vs 256 deaths in the placebo group). The risk of CHD mortality was significantly reduced by 42% (p=0.00001, 111 deaths in the simvastatin group vs 189 deaths in the placebo group). There was no statistically significant difference between groups in non-cardiovascular mortality.
- Simvastatin significantly reduced the risk for the secondary composite endpoint (time to first occurrence of CHD death, definite or probable hospital verified non-fatal MI, silent MI verified by ECG, or resuscitated cardiac arrest) by 34% (p<0.00001, 431 vs 622 patients with one or more events). Simvastatin reduced the risk of major coronary events to a similar extent across the range of baseline total and LDL cholesterol levels. The risk of having a hospital-verified non-fatal MI was reduced by 37%.
- Simvastatin significantly reduced the risk for undergoing myocardial revascularization procedures (coronary artery bypass grafting or percutaneous transluminal coronary angioplasty) by 37% (p<0.00001, 252 vs 383 patients).
- Simvastatin significantly reduced the risk of fatal plus non-fatal cerebrovascular events (combined stroke and transient ischemic attacks) by 28% (p=0.033, 75 vs

102 patients).

- Over the course of the trial, treatment with simvastatin led to mean reductions in total-C, LDL-C and triglycerides (TG) of 25%, 35%, and 10%, respectively, and a mean increase in high-density lipoprotein cholesterol (HDL-C) of 8%. In contrast, treatment with placebo led to increases in total-C, LDL-C and TG of 1%, 1%, and 7%, respectively.
- Because there were only 53 female deaths (approximately 18% of the trial population was female), the effect of simvastatin on mortality in females could not be adequately assessed. However, simvastatin significantly reduced the risk of having major coronary events in females by 34% (60 vs 91 women with one or more event).
- Simvastatin resulted in similar decreases in relative risk for total mortality, CHD mortality, and major coronary events in geriatric patients (≥ 65 years) compared with younger adults.

The Heart Protection Trial (Trial HPS) was a randomized, placebo-controlled, double-blind, multi-centered trial with a mean duration of 5 years conducted in 10,269 patients on simvastatin 40 mg and 10,267 on placebo. Patients had a mean age of 64 years (range 40 to 80 years old), 97% were White, and were at high risk of developing a major coronary event because of existing CHD (65%), diabetes (Type 2, 26%; Type 1, 3%), history of stroke or other cerebrovascular disease (16%), peripheral vascular disease (33%), or they were males ≥ 65 years with hypertension in (6%). At baseline:

- 3,421 patients (17%) had LDL-C levels below 100 mg/dL, including 953 (5%) below 80 mg/dL; and
- 10,047 patients (49%) had levels greater than 130 mg/dL.

Patients were randomized to simvastatin or placebo using a covariate adaptive method which considered the distribution of 10 important baseline characteristics of patients already enrolled.

The Trial HPS results showed that simvastatin 40 mg/day significantly reduced: total and CHD mortality; and non-fatal MI, stroke, and revascularization procedures (coronary and non-coronary) (see Table 15).

Table 15: CHD Mortality and Cardiovascular Events in Adult Patients with High Risk of Developing a Major Coronary Event in Trial HPS

Endpoint	Simvastatin (N=10,269) n (%)*	Placebo (N=10,267) n (%)*	Risk Reduction (%) (95% CI)	p-Value
Primary Mortality	1,328 (12.9%)	1,507 (14.7%)	13% (6 to 19%)	p=0.0003
CHD mortality	587 (5.7%)	707 (6.9%)	18% (8 to 26%)	p=0.0005
Secondary Non-fatal MI	357 (3.5%)	574 (5.6%)	38% (30 to 46%)	p<0.0001

Stroke	444 (4.3%)	585 (5.7%)	25% (15 to 34%)	p<0.0001
Tertiary Coronary revascularization	513 (5%)	725 (7.1%)	30% (22 to 38%)	p<0.0001
Peripheral and other non- coronary revascularization	450 (4.4%)	532 (5.2%)	16% (5 to 26%)	p=0.006

* n = number of patients with indicated event

Two composite endpoints were defined to have enough events to assess relative risk reductions across a range of baseline characteristics:

- Major coronary events (MCE) was comprised of CHD mortality and non-fatal MI. Analyzed by time-to-first event; 898 patients (8.7%) treated with simvastatin had events and 1,212 patients (11.8%) treated with placebo had events.
- Major vascular events (MVE) was comprised of MCE, stroke, and revascularization procedures including coronary, peripheral and other non-coronary procedures. Analyzed by time-to-first event; 2,033 patients (19.8%) treated with simvastatin had events and 2,585 patients (25.2%) on placebo had events.

Simvastatin use led to significant relative risk reductions for both composite endpoints (27% for MCE and 24% for MVE, p<0.0001) and for all components of the composite endpoints. The risk reductions produced by simvastatin in both MCE and MVE were evident and consistent regardless of cardiovascular disease related medical history at trial entry (i.e., CHD alone; or peripheral vascular disease, cerebrovascular disease, diabetes or treated hypertension, with or without CHD), gender, age, baseline levels of LDL-C, baseline concomitant cardiovascular medications (i.e., aspirin, beta blockers, or calcium channel blockers), smoking status, or obesity. Patients with diabetes showed risk reductions for MCE and MVE due to simvastatin treatment regardless of baseline HbA1c levels or obesity.

16 HOW SUPPLIED/STORAGE AND HANDLING

- Ezetimibe and simvastatin tablets 10/10, (ezetimibe, USP 10 mg/simvastatin, USP 10 mg tablets) are white to off-white, capsule shaped tablets debossed with "A029" on one side and plain on other side.

They are supplied as follows:

NDC 67877-507-30 bottles of 30

NDC 67877-507-90 bottles of 90

NDC 67877-507-01 bottles of 1000 (If repackaged in blisters, then opaque or light-resistant blisters should be used.)

NDC 67877-507-38 carton of 100 (10 x 10) unit-dose tablets

- Ezetimibe and simvastatin tablets 10/20, (ezetimibe, USP 10 mg/simvastatin, USP 20 mg tablets) are white to off-white, capsule shaped tablets debossed with “A028” on one side and plain on other side.

They are supplied as follows:

NDC 67877-508-30 bottles of 30

NDC 67877-508-90 bottles of 90

NDC 67877-508-01 bottles of 1000 (If repackaged in blisters, then opaque or light-resistant blisters should be used.)

NDC 67877-508-38 carton of 100 (10 x 10) unit-dose tablets

- Ezetimibe and simvastatin tablets 10/40, (ezetimibe, USP 10 mg/simvastatin, USP 40 mg tablets) are white to off-white, capsule shaped tablets debossed with “A027” on one side and plain on other side.

They are supplied as follows:

NDC 67877-509-30 bottles of 30

NDC 67877-509-90 bottles of 90

NDC 67877-509-05 bottles of 500 (If repackaged in blisters, then opaque or light-resistant blisters should be used.)

NDC 67877-509-37 carton of 50 (5 x 10) unit-dose tablets

- Ezetimibe and simvastatin tablets 10/80, (ezetimibe, USP 10 mg/simvastatin, USP 80 mg tablets) are white to off-white, capsule shaped tablets debossed with “A026” on one side and plain on other side.

They are supplied as follows:

NDC 67877-510-30 bottles of 30

NDC 67877-510-90 bottles of 90

NDC 67877-510-05 bottles of 500 (If repackaged in blisters, then opaque or light-resistant blisters should be used.)

NDC 67877-510-37 carton of 50 (5 x 10) unit-dose tablets

Storage

Store at 20 to 25° C (68 to 77 °F). [See USP Controlled Room Temperature.] Keep container tightly closed.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).

Myopathy and Rhabdomyolysis

Advise patients that ezetimibe and simvastatin tablets may cause myopathy and rhabdomyolysis. Inform patients taking the 80 mg daily dose of simvastatin that they are at an increased risk. Inform patients that the risk is also increased when taking certain types of medication or consuming grapefruit juice and they should discuss all medication, both prescription and over the counter, with their healthcare provider. Instruct patients to inform other healthcare providers prescribing a new medication or increasing the dose of an existing medication that they are taking ezetimibe and simvastatin tablets. Instruct patients to promptly report any unexplained muscle pain,

tenderness or weakness particularly if accompanied by malaise or fever [see *Contraindications (4), Warnings and Precautions (5.1), and Drug Interactions (7.1)*].

Hepatic Dysfunction

Inform patients that ezetimibe and simvastatin tablets may cause liver enzyme elevations and possibly liver failure. Advise patients to promptly report fatigue, anorexia, right upper abdominal discomfort, dark urine or jaundice [see *Warnings and Precautions (5.3)*].

Increases in HbA1c and Fasting Serum Glucose Levels

Inform patients that increases in HbA1c and fasting serum glucose levels may occur with ezetimibe and simvastatin tablets. Encourage patients to optimize lifestyle measures, including regular exercise, maintaining a healthy body weight, and making healthy food choices [see *Warnings and Precautions (5.4)*].

Pregnancy

Advise pregnant patients and patients who can become pregnant of the potential risk to a fetus. Advise patients to inform their healthcare provider of a known or suspected pregnancy to discuss if ezetimibe and simvastatin tablets should be discontinued [see *Use in Specific Populations (8.1)*].

Lactation

Advise patients that breastfeeding is not recommended during treatment with ezetimibe and simvastatin tablets [see *Use in Specific Populations (8.2)*].

Missed Dose

Instruct patients to take ezetimibe and simvastatin tablets only as prescribed. If a dose is missed, it should be taken as soon as possible. Advise patients not to double their next dose.

For Patient Information, please visit:
http://www.ascendlaboratories.com/pti/ezetim_simvatablets.pdf

Manufactured by:

Alkem Laboratories Ltd.
Mumbai - 400 013, INDIA.

Distributed by:

Ascend Laboratories, LLC
Parsippany, NJ 07054

Revised: May, 2024

Patient Information

Ezetimibe and Simvastatin Tablets

(e-ZET-i-mibe / SIM-va-STAT-in)

Read this Information carefully before you start taking ezetimibe and simvastatin tablets and each time you get a ezetimibe and simvastatin tablets.

There may be new information. This information does not take the place of talking with your healthcare provider about your medical condition or your treatment. If you have any questions about ezetimibe and simvastatin tablets, ask your healthcare provider. Only your healthcare provider can determine if ezetimibe and simvastatin tablets is right for you.

What is ezetimibe and simvastatin tablets?

Ezetimibe and simvastatin tablets is a prescription medicine that contains the cholesterol lowering medicines, simvastatin and ezetimibe:

- Ezetimibe and simvastatin tablets is used along with diet to lower elevated low-density lipoprotein cholesterol (LDL-C) or bad cholesterol in:
 - o adults with primary hyperlipidemia (high level of fats in your blood).
 - o adults and children 10 years of age and older with heterozygous familial hypercholesterolemia (HeFH). HeFH is an inherited condition that causes high levels of bad cholesterol.
- Ezetimibe and simvastatin tablets is also used with other cholesterol lowering treatments to lower elevated LDL-C levels in adults with homozygous familial hypercholesterolemia (HoFH). HoFH is an inherited condition that causes high levels of bad cholesterol.
- Simvastatin when used as a component of ezetimibe and simvastatin tablets is used to lower:
 - o the risk of death by lowering the risk of heart disease death.
 - o the risk of heart attacks and strokes.
 - o the need for certain types of heart and blood vessel procedures to improve blood flow called arterial revascularization in people with known heart, cerebrovascular disease (conditions that affect blood flow and the blood vessels in the brain), peripheral vascular disease (a blood circulation disorder that causes the blood vessels outside of your heart and brain to narrow, block, or spasm), and diabetes, who are at high risk for heart disease problems.

The safety and effectiveness of ezetimibe and simvastatin tablets has not been established in children younger than 10 years of age with inherited heterozygous familial

hypercholesterolemia (HeFH) or other types of hyperlipidemia.

Do not take ezetimibe and simvastatin tablets if you:

- take certain medicines called CYP3A4 inhibitors such as:
 - o certain antifungal medicines (such as itraconazole, ketoconazole, posaconazole, voriconazole).
 - o certain antibiotics (including erythromycin, clarithromycin, telithromycin).
 - o HIV protease inhibitors (such as indinavir, nelfinavir, ritonavir, saquinavir, tipranavir, or atazanavir and cobicistat-containing products such as (elvitegravir/cobicistat/emtricitabine/tenofovir disoproxil fumarate).
 - o certain hepatitis C virus protease inhibitors (such as boceprevir or telaprevir).
 - o the antidepressant nefazodone.
- take medicines called cyclosporine, danazol, or gemfibrozil.
- have liver problems.
- are allergic to simvastatin, ezetimibe, or any of the ingredients in ezetimibe and simvastatin tablets. See the end of this Patient Information leaflet for a complete list of ingredients in ezetimibe and simvastatin tablets.

Ask your healthcare provider or pharmacist if you are not sure if your medicine is listed above.

Before you take ezetimibe and simvastatin tablets, tell your healthcare provider about all of your medical conditions, including if you:

- have unexplained muscle aches or weakness.
- have or have had myasthenia gravis (a disease causing general muscle weakness including in some cases muscles used for breathing), ocular myasthenia (a disease causing eye muscle weakness).
 - have kidney problems.
- have liver problems or drink more than 2 glasses of alcohol daily.
- have thyroid problems.
- are 65 years of age or older.
- are of Chinese descent.
- are pregnant or plan to become pregnant. If you become pregnant while taking ezetimibe and simvastatin tablets, call your healthcare provider right away to discuss stopping ezetimibe and simvastatin tablets.
- are breastfeeding or plan to breastfeed. It is not known if ezetimibe and simvastatin tablets passes into your breast milk.

Do not breastfeed while taking ezetimibe and simvastatin tablets.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Talk to your healthcare provider before you start taking any new medicines.

Tell your healthcare provider who prescribes ezetimibe and simvastatin tablets if another healthcare provider increases the dose of another medicine you are taking.

Ezetimibe and simvastatin tablets may affect the way other medicines work, and other medicines may affect how ezetimibe and simvastatin tablets works. Especially tell your healthcare provider if you take:

- digoxin (a drug used to treat irregular heartbeat).
- coumarin anticoagulants (drugs that prevent blood clots, such as warfarin).
- fibric acid derivatives (such as fenofibrate).

Taking ezetimibe and simvastatin tablets with certain substances can also increase the risk of muscle problems. Especially tell your healthcare provider if you take:

- amiodarone or dronedarone (medicines used to treat an irregular heartbeat).
- verapamil, diltiazem, amlodipine, or ranolazine (medicines used to treat high blood pressure, chest pain associated with heart disease, or other heart conditions).
- lomitapide (a medicine used to treat a serious and rare genetic cholesterol condition).
- daptomycin (a drug used to treat complicated skin and bloodstream infections).
- large doses of niacin or nicotinic acid, especially if you are of Chinese descent.
- colchicine (a medicine used to treat gout).
- grapefruit juice.

Ask your healthcare provider or pharmacist for a list of medicines if you are not sure. Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

How should I take ezetimibe and simvastatin tablets?

- Take ezetimibe and simvastatin tablets exactly as your healthcare provider tells you to take it.
- **Do not** change your dose or stop taking ezetimibe and simvastatin tablets without talking to your healthcare provider.
- Take ezetimibe and simvastatin tablets 1 time each day in the evening.
- If you miss a dose, take it as soon as you remember. If you do not remember until it is time for your next dose, skip the missed dose and go back to your regular schedule. Do not take 2 doses of ezetimibe and simvastatin tablets at the same time. Talk with your healthcare provider if you have questions about a missed dose.
- While taking ezetimibe and simvastatin tablets, continue to follow your cholesterol-lowering diet and to exercise as your healthcare provider told you to.
- Your healthcare provider may do blood tests to check your cholesterol while you take

ezetimibe and simvastatin tablets. Your healthcare provider may change your dose of ezetimibe and simvastatin tablets if needed.

If you take too much ezetimibe and simvastatin tablets, call your healthcare provider or Poison Help Line at 1-800-222-1222 or go to the nearest hospital emergency room right away.

What are the possible side effects of ezetimibe and simvastatin tablets?

Ezetimibe and simvastatin tablets may cause serious side effects including:

- **Muscle pain, tenderness, and weakness (myopathy).** Muscle problems, including muscle breakdown, can be serious in some people and rarely cause kidney damage that can lead to death.

Tell your healthcare provider right away if:

- o **you have unexplained muscle pain, tenderness, or weakness, especially if you have a fever or feel more tired than usual, while you take ezetimibe and simvastatin tablets.**

- o you have muscle problems that do not go away even after your healthcare provider has advised you to stop taking ezetimibe and simvastatin tablets. Your healthcare provider may do further tests to diagnose the cause of your muscle problems.

Your chances of getting muscle problems are higher if you:

- o are taking certain other medicines while you take ezetimibe and simvastatin tablets.
- o are 65 years of age or older.
- o are female.
- o have thyroid problems (hypothyroidism) that are not controlled.
- o have kidney problems.
- o are taking higher doses of ezetimibe and simvastatin tablets.
- o are Chinese.

- **Liver problems.** Your healthcare provider should do blood tests to check your liver before you start taking ezetimibe and simvastatin tablets and if you have any symptoms of liver problems while you take ezetimibe and simvastatin tablets.

Call your healthcare provider right away if you have the following symptoms of liver problems:

- o feeling tired or weak
- o loss of appetite
- o right-sided upper belly pain
- o dark urine
- o yellowing of your skin or the whites of your eyes

- **increase in blood sugar (glucose) levels.** Ezetimibe and simvastatin tablets may cause an increase in your blood sugar levels.

The most common side effects of ezetimibe and simvastatin tablets include:

- headache
- increased liver enzyme levels
- muscle pain
- upper respiratory infection
- diarrhea

Tell your healthcare provider if you have any side effect that bothers you or does not go away.

These are not all the possible side effects of ezetimibe and simvastatin tablets.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store ezetimibe and simvastatin tablets?

- Store ezetimibe and simvastatin tablets at room temperature between 68°F to 77°F (20°C to 25°C).
- Keep ezetimibe and simvastatin tablets in its original container until you use it.
- Keep ezetimibe and simvastatin tablets in a tightly closed container.

Keep ezetimibe and simvastatin tablets and all medicines out of the reach of children.

General information about safe and effective use of ezetimibe and simvastatin tablets.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use ezetimibe and simvastatin tablets for a condition for which it was not prescribed. Do not give ezetimibe and simvastatin tablets to other people, even if they have the same symptoms that you have. It may harm them.

You can ask your pharmacist or healthcare provider for information about ezetimibe and simvastatin tablets that is written for health professionals.

For more information, call 1-877-272-7901.

What are the ingredients in ezetimibe and simvastatin tablets?

Active Ingredients: ezetimibe, USP and simvastatin, USP

Inactive Ingredients: Lactose Monohydrate, Microcrystalline Cellulose, Croscarmellose Sodium,

Sodium Lauryl Sulfate, Hypromellose, Citric Acid Monohydrate, Propyl Gallate, Butylated Hydroxy Anisole

and Magnesium Stearate.

This Patient Information has been approved by the U.S. Food and

Drug Administration.

For Patient Information, please visit:

http://www.ascendlaboratories.com/pti/ezetim_simvatablets.pdf

Manufactured by:

Alkem Laboratories Ltd.

Mumbai - 400 013, INDIA.

Distributed by:

Ascend Laboratories, LLC
Parsippany, NJ 07054

Revised: May, 2024

PT 2626-07

PACKAGE LABEL. PRINCIPAL DISPLAY PANEL

ASCEND Laboratories, LLC

NDC 67877-507-30

Ezetimibe and Simvastatin Tablets 10 mg/10 mg

Rx only

30 Tablets

Each tablet contains 10 mg ezetimibe, USP and 10 mg simvastatin, USP.
Store at 20 to 25°C (68 to 77°F).
[See USP Controlled Room Temperature.]
Keep container tightly closed.

USUAL DOSAGE: See package insert.

Manufactured by:
ALKEM Alkem Laboratories Ltd., INDIA.
Distributed by: Ascend Laboratories, LLC
Parsippany, NJ 07054

Code No.: HP/Drugs/MNB/05/105
PL9897-02

NDC 67877-507-30
Ezetimibe and Simvastatin Tablets
10 mg/10 mg
Rx Only 30 Tablets

ASCEND
Laboratories, LLC

Unvarnished area
45 x 20 mm (LXH)
Rest label should be
with UV Varnish

For Patient Information, please visit:
http://www.ascendlaboratories.com/pti/ezetim_simvatablets.pdf

ASCEND Laboratories, LLC

NDC 67877-508-01

Ezetimibe and Simvastatin Tablets 10 mg/20 mg

Rx only
1,000 Tablets

Each tablet contains 10 mg ezetimibe, USP and 20 mg simvastatin, USP.

Store at 20 to 25°C (68 to 77°F).
[See USP Controlled Room Temperature.]
Dispense in a tightly-closed container.
This is a bulk package and not intended for dispensing. Package is not child-resistant.

USUAL DOSAGE: See package insert.

For Patient Information, please visit:
http://www.ascendlaboratories.com/pti/ezetim_simvastatblts.pdf

Manufactured by:
ALKEM Alkem Laboratories Ltd., INDIA.

Distributed by: Ascend Laboratories, LLC
Parsippany, NJ 07054

NDC 67877-508-01

Ezetimibe and Simvastatin Tablets

10 mg/20 mg

Rx Only 1,000 Tablets

ASCEND
laboratories, LLC

Unvarnished area
45 x 30 mm (LXH)
Rest label should be
with UV Varnish

N
3 67877 150801 6

Code No.:
HP/Drugs/MNB/05/105 PL9923-02

ASCEND Laboratories, LLC
NDC 67877-509-90

Ezetimibe and Simvastatin Tablets 10 mg/40 mg
Rx only
90 Tablets

Each tablet contains 10 mg ezetimibe, USP and 40 mg simvastatin, USP.

Store at 20 to 25°C (68 to 77°F).
[See USP Controlled Room Temperature.]
Keep container tightly closed.

USUAL DOSAGE: See package insert.

For Patient Information, please visit:
http://www.ascendlaboratories.com/pti/ezetim_simvastatblts.pdf

Manufactured by:
ALKEM Alkem Laboratories Ltd., INDIA.

Distributed by: Ascend Laboratories, LLC
Parsippany, NJ 07054

NDC 67877-509-90

Ezetimibe and Simvastatin Tablets

10 mg/40 mg

Rx Only 90 Tablets

ASCEND
laboratories, LLC

Unvarnished area
45 x 28 mm (LXH)
Rest label should be
with UV Varnish

Code No.:
HP/Drugs/MNB/05/105 PL9925-02

N
3 67877 150990 7

ASCEND Laboratories, LLC
NDC 67877-510-05

Ezetimibe and Simvastatin Tablets 10 mg/80 mg
 Rx only
 500 Tablets

Each tablet contains 10 mg ezetimibe, USP and 80 mg simvastatin, USP.

Store at 20 to 25°C (68 to 77°F).
 [See USP Controlled Room Temperature.]
 Dispense in a tightly-closed container.
 This is a bulk package and not intended for dispensing. Package is not child-resistant.

USUAL DOSAGE: See package insert.

For Patient Information, please visit:
http://www.ascendlaboratories.com/pti/ezetim_simvatablets.pdf

Manufactured by:
 ALKEM Alkem Laboratories Ltd., INDIA.

Distributed by: Ascend Laboratories, LLC
 Parsippany, NJ 07054

NDC 67877-510-05

Ezetimibe and Simvastatin Tablets

10 mg/80 mg

Rx Only 500 Tablets

ASCEND
 Laboratories, LLC

Unvarnished area
 45 x 30 mm (LXH)
 Rest label should be
 with UV Varnish

N
 3 67877 51005 7

Code No.:
 HPIDrugs/MNB/05105 PL9829-02

EZETIMIBE AND SIMVASTATIN

ezetimibe and simvastatin tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:67877-507
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
EZETIMIBE (UNII: EOR26LQQ24) (EZETIMIBE - UNII:EOR26LQQ24)	EZETIMIBE	10 mg
SIMVASTATIN (UNII: AGG2FN16EV) (SIMVASTATIN - UNII:AGG2FN16EV)	SIMVASTATIN	10 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	

ISOPROPYL ALCOHOL (UNII: ND2M416302)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	WHITE (White to off white)	Score	no score
Shape	CAPSULE	Size	8mm
Flavor		Imprint Code	A029
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67877-507-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	12/24/2017	
2	NDC:67877-507-90	90 in 1 BOTTLE; Type 0: Not a Combination Product	12/24/2017	
3	NDC:67877-507-01	1000 in 1 BOTTLE; Type 0: Not a Combination Product	12/24/2017	
4	NDC:67877-507-38	10 in 1 CARTON	12/24/2017	
4	NDC:67877-507-33	10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209222	12/24/2017	

EZETIMIBE AND SIMVASTATIN

ezetimibe and simvastatin tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:67877-508
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
EZETIMIBE (UNII: EOR26LQQ24) (EZ ETIMIBE - UNII:EOR26LQQ24)	EZETIMIBE	10 mg
SIMVASTATIN (UNII: AGG2FN16EV) (SIMVASTATIN - UNII:AGG2FN16EV)	SIMVASTATIN	20 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	WHITE (White to off white)	Score	no score
Shape	CAPSULE	Size	11mm
Flavor		Imprint Code	A028
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67877-508-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	12/24/2017	
2	NDC:67877-508-90	90 in 1 BOTTLE; Type 0: Not a Combination Product	12/24/2017	
3	NDC:67877-508-01	1000 in 1 BOTTLE; Type 0: Not a Combination Product	12/24/2017	
4	NDC:67877-508-38	10 in 1 CARTON	12/24/2017	
4	NDC:67877-508-33	10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209222	12/24/2017	

EZETIMIBE AND SIMVASTATIN

ezetimibe and simvastatin tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:67877-509
---------------------	-------------------------	---------------------------	---------------

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
EZETIMIBE (UNII: EOR26LQQ24) (EZ ETIMIBE - UNII:EOR26LQQ24)	EZETIMIBE	10 mg
SIMVASTATIN (UNII: AGG2FN16EV) (SIMVASTATIN - UNII:AGG2FN16EV)	SIMVASTATIN	40 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
WATER (UNII: 059QF0K00R)	

Product Characteristics

Color	WHITE (White to off white)	Score	no score
Shape	CAPSULE	Size	14mm
Flavor		Imprint Code	A027
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67877-509-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	12/24/2017	
2	NDC:67877-509-90	90 in 1 BOTTLE; Type 0: Not a Combination Product	12/24/2017	
3	NDC:67877-509-05	500 in 1 BOTTLE; Type 0: Not a Combination Product	12/24/2017	
4	NDC:67877-509-37	05 in 1 CARTON	12/24/2017	
4	NDC:67877-509-33	10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209222	12/24/2017	

EZETIMIBE AND SIMVASTATIN

ezetimibe and simvastatin tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:67877-510
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
EZETIMIBE (UNII: EOR26LQQ24) (EZETIMIBE - UNII:EOR26LQQ24)	EZETIMIBE	10 mg
SIMVASTATIN (UNII: AGG2FN16EV) (SIMVASTATIN - UNII:AGG2FN16EV)	SIMVASTATIN	80 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
PROPYL GALLATE (UNII: 8D45NN7V92)	
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	WHITE (White to off white)	Score	no score
Shape	CAPSULE	Size	18mm
Flavor		Imprint Code	A026
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67877-510-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	12/24/2017	
2	NDC:67877-510-90	90 in 1 BOTTLE; Type 0: Not a Combination Product	12/24/2017	
3	NDC:67877-510-05	500 in 1 BOTTLE; Type 0: Not a Combination Product	12/24/2017	
4	NDC:67877-510-37	05 in 1 CARTON	12/24/2017	

4 NDC:67877-510-33 | 10 in 1 BLISTER PACK; Type 0: Not a Combination Product

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209222	12/24/2017	

Labeler - Ascend Laboratories, LLC (141250469)

Establishment

Name	Address	ID/FEI	Business Operations
Alkem Laboratories Limited		677605851	ANALYSIS(67877-507, 67877-508, 67877-509, 67877-510) , MANUFACTURE(67877-507, 67877-508, 67877-509, 67877-510) , PACK(67877-507, 67877-508, 67877-509, 67877-510)

Revised: 8/2024

Ascend Laboratories, LLC