

NON ASPIRIN PM EXTRA STRENGTH- acetaminophen, diphenhydramine hcl tablet, coated
Chain Drug Consortium, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

DRUG FACTS

Active ingredients (in each geltab)

Acetaminophen 500 mg

Diphenhydramine HCl 25 mg

Purpose

Pain reliever

Nighttime sleep-aid

Uses

temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other products containing diphenhydramine, even one used on skin
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery

Stop use and ask a doctor if

- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness.
- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**
- adults and children 12 years and over: take 2 gels at bedtime. Do not take more than 2 gels of this product in 24 hours.
- children under 12 years: do not use

Other information

- store at room temperature 15°- 30° C (59°- 86° F)
- avoid high humidity and excessive heat

Inactive ingredients

corn starch*, croscarmellose sodium*, D&C red #27 aluminum lake, edible black ink, FD&C blue #1 aluminum lake, gelatin, glycerin, hypromellose*, maltodextrin*, microcrystalline cellulose*, polyethylene glycol*, povidone*, purified water, silicon dioxide*, stearic acid, titanium dioxide
*contains one or more of these ingredients

Questions or comments?

Call toll free 1-877-753-3935 Monday- Friday 9AM- 5PM EST

Principal Display Panel

** Compare to active ingredients in extra strength TYLENOL® PM

extra strength Non- Aspirin PM

ACETAMINOPHEN 500 mg

DIPHENHYDRAMINE HCl 25 mg

- pain reliever
- nighttime sleep-aid

**This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Extra Strength Tylenol® PM

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Product Label

**NO PRINT
NO VARNISH
NO COATING**

**DO NOT USE IF IMPRINTED SAFETY SEAL
UNDER CAP IS BROKEN OR MISSING**

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery

Stop use and ask a doctor if

- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.
- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)
- adults and children 12 years and over: take 2 tablets at bedtime. Do not take more than 2 tablets of this product in 24 hours.
- children under 12 years: do not use

Other Information

- store at room temperature 15°-30°C (59°-86°F)
- avoid high humidity and excessive heat

Inactive ingredients corn starch*, croscarmellose sodium*, D&C red #27 aluminum lake, edible black ink, FD&C blue #1 aluminum lake, gelatin, glycerin, hypromellose*, maltodextrin*, microcrystalline cellulose*, polyethylene glycol*, povidone*, purified water, silicon dioxide*, stearic acid, titanium dioxide

*contains one or more of these ingredients

Questions or comments?
Call toll free 1-877-753-3935
Monday-Friday 9AM-5PM EST

****This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Extra Strength Tylenol® PM.**

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

DISTRIBUTED BY: CHAIN DRUG CONSORTIUM, LLC
3301 NW 80th AVE. SUITE 101
BOCA RATON, FL 33431

If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

PLD-D
FC000991

50 Gelatin Tablets

Acetaminophen 500 mg
Diphenhydramine HCl 25 mg
Pain Reliever * Nighttime Sleep-Aid

Non-Aspirin PM

Extra Strength

Premier Value®

50 Gelatin Tablets

DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each tablet)

Acetaminophen 500 mg.....Pain reliever
Diphenhydramine HCl 25 mg.....Nighttime sleep-aid

Uses temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- glaucoma

10049501

0290620

Acetaminophen PM gels

NON ASPIRIN PM EXTRA STRENGTH

acetaminophen, diphenhydramine hcl tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68016-188
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg

Inactive Ingredients	
Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
D&C RED NO. 27 (UNII: 2LRS185U6K)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
POVIDONES (UNII: FZ989GH94E)	
WATER (UNII: 059QF0KO0R)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
ALUMINUM OXIDE (UNII: LM26O6933)	

Product Characteristics			
Color	BLUE, WHITE	Score	no score
Shape	ROUND	Size	13mm
Flavor		Imprint Code	BPI;50
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68016-188-50	1 in 1 CARTON		
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	10/24/2012	

Labeler - Chain Drug Consortium, LLC (101668460)

Revised: 10/2015

Chain Drug Consortium, LLC