

# **THOMPSON CHEWABLE BISMUTH TABS- bismuth subsalicylate tablet, chewable**

**Thompson Safety LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

## **Thompson Chewable Bismuth Tabs**

### **Drug Facts**

#### **Active ingredient**

Bismuth Subsalicylate 262 mg  
(each tablet contains 102 mg salicylate)

#### **Purpose**

Upset stomach reliever/antidiarrheal

#### **Uses**

relieves

- traveler's diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including: • heartburn, • indigestion, • nausea, • gas, • belching, • fullness

#### **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome a rare but serious illness.

#### **Allergy alert: Contains salicylate. Do not take if you are:**

- allergic to salicylates (including aspirin)
- taking other salicylate products

#### **Do not use if you have**

- bloody or black stool
- an ulcer
- a bleeding problem

#### **Ask a doctor before use if you have**

- fever
- mucus in the stool

#### **Ask a doctor or pharmacist if you are taking any drug for**

- anticoagulation (thinning of the blood)
- diabetes
- gout

- arthritis

**When using this product** a temporary and harmless darkening of the tongue and/or stool may occur.

**Stop use and ask a doctor if**

- symptoms get worse or lasts more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

In case of accidental overdose, contact a physician or poison control center right away (1-800-222-1222)

**Directions**

- chew or dissolve in mouth
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Adults and children: (12 years and over)**

- chew 2 tablets every 1/2 to 1 hour as needed
- do not exceed 16 tablets in 24 hours
- use until diarrhea stops but not more than 2 days

**Children under 12 years:** ask a doctor

**Other information**

- **phenylketonurics:** contains phenylalanine 1.1mg per tablet
- **calcium** content per tablet: 73mg
- store at room temperature 59°-86°F (15°-30°C)
- tamper-evident sealed packets
- do not use any opened or torn packets

**Inactive ingredients**

acacia gum, aspartame, calcium carbonate, D&C red #27 aluminum lake, dextrans, magnesium stearate, maltodextrin, microcrystalline cellulose, peppermint flavor, silicon dioxide

**Questions or comments?** 877.506.4291

Chewable Bismuth Tabs

Upset Stomach / Diarrhea Relief

Thompson

Safety Works | Business Thrives™

Bismuth Subsalicylate 262 MG

36 Tablets

Tamper Evident

### Drug Facts

**Active ingredient (in each tablet)** **Purpose**  
Bismuth Subsalicylate 262 mg .....Upset stomach reliever/antidiarrheal (each tablet contains 102 mg salicylate)

**Uses**  
relieves ■ travelers' diarrhea ■ diarrhea  
■ upset stomach due to overindulgence in food and drink, including:  
■ heartburn ■ indigestion ■ nausea  
■ gas ■ belching ■ fullness

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome a rare but serious illness.

**Allergy alert:** Contains salicylate. Do not take if you are: ■ allergic to salicylates (including aspirin) ■ taking other salicylate products

**Do not use if you have**  
■ bloody or black stool ■ an ulcer ■ a bleeding problem

**Ask a doctor before use if you have** ■ fever ■ mucus in the stool

**Ask a doctor or pharmacist if you are taking any drug for**  
■ anticoagulation (thinning of the blood) ■ diabetes ■ gout ■ arthritis

**When using this product** a temporary and harmless darkening of the tongue and/or stool may occur.

**Stop use and ask a doctor if**  
■ symptoms get worse or lasts for more than 2 days  
■ ringing in the ears or loss of hearing occurs  
■ diarrhea lasts more than 2 days

**Drug Facts (continued)**

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.** In case of accidental overdose, contact a physician or poison control center right away (1-800-222-1222).

**Directions**  
■ chew or dissolve in mouth  
■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Adults and children:** (12 years and older) ■ chew 2 tablets every 1/2 to 1 hour as needed  
■ do not exceed 16 tablets in 24 hours  
■ use until diarrhea stops, but not more than 2 days

**Children under 12 years:** Ask a doctor

**Other information**  
■ phenylketonurics: contains phenylalanine 1.1 mg per tablet  
■ calcium content per tablet: 73 mg  
■ store at room temperature 59°-86°F (15°-30°C)  
■ tamper-evident sealed packets  
■ do not use any opened or torn packets  
*(continued on opposite panel)*

# CHEWABLE BISMUTH TABS

UPSET STOMACH / DIARRHEA RELIEF

## THOMPSON

SAFETY WORKS | BUSINESS THRIVES™

BISMUTH SUBSALICYLATE 262 MG

36 Tablets
TAMPER EVIDENT

**Drug Facts (continued)**

**Inactive ingredients**  
acacia gum, aspartame, calcium carbonate, D&C red #27 aluminum lake, dextrates, magnesium stearate, maltodextrin, microcrystalline cellulose, peppermint flavoring, silicon dioxide

**Questions or comments?**  
877.506.4291

Reorder CS41220

Distributed by  
**Thompson Safety**  
Houston, TX 77073  
877.506.4291  
www.thompson-safety.com

0 73408 4 1220 1

THOMPSON CHEWABLE BISMUTH TABS			
bismuth subsalicylate tablet, chewable			
Product Information			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:73408-407
<b>Route of Administration</b>	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ, BISMUTH CATION - UNII:ZS9CD118YE)	BISMUTH SUBSALICYLATE	262 mg	
Inactive Ingredients			
Ingredient Name	Strength		
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)			
<b>DEXTRATES</b> (UNII: G263MI44RU)			
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)			
<b>CALCIUM CARBONATE</b> (UNII: H0G9379FGK)			
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)			
<b>D&amp;C RED NO. 27</b> (UNII: 2LRS185U6K)			
<b>ACACIA</b> (UNII: 5C5403N26O)			
<b>ASPARTAME</b> (UNII: Z0H242BBR1)			
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)			
Product Characteristics			
<b>Color</b>	pink	<b>Score</b>	no score

<b>Shape</b>	ROUND	<b>Size</b>	16mm
<b>Flavor</b>		<b>Imprint Code</b>	RH;046
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73408-407-20	6 in 1 BOX	08/11/2021	
1		6 in 1 CELLO PACK; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part335	08/11/2021	

**Labeler** - Thompson Safety LLC (080998015)

**Registrant** - Unifirst First Aid Corporation (832947092)