

**TUKOL HONEY DAYTIME COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride syrup**  
**Genomma Lab USA**

-----  
**Tukol® Honey Daytime Cold & Flu**

***Drug Facts***

<b>Active Ingredients (in each 30 mL)</b>	<b>Purpose</b>
<b>Acetaminophen 650 mg</b>	Pain reliever/Fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Phenylephrine HCl 10 mg	Nasal Decongestant

**Uses**

Temporarily relieves cold/flu symptoms:

- minor aches and pain
- headache
- sore throat
- nasal congestion
- cough due to minor throat and bronchial irritation
- temporarily reduces fever

**Warnings**

**Liver warning**

The product contains Acetaminophen. Severe liver damage may occur if you take

- more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert**

Acetaminophen may cause severe skin reactions. Symptoms may include

- skin reddening
- blisters
- rash.

If skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for two weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough accompanied by excessive phlegm (mucus)

**Ask a doctor or pharmacist before use**

- if you are taking the blood thinning drug warfarin

**When using this product**

- do not use more than directed (see overdose warning)
- avoid alcoholic drinks

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- fever gets worse or lasts more than 3 days
- pain or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- redness or swelling is present
- cough comes back, or occurs with rash or headache that lasts.
- new symptoms occur
- These could be signs of a serious condition.

**If pregnant or breast feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning**

Taking more than the recommended dose (overdose) may cause liver damage. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed**
- do not take more than 4 doses in any 24 hours
- this adult strength product is not intended for use in children under 12 years of age
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- TBSP = tablespoon
- dose as follows

<b>age</b>	<b>dose</b>
adults and children 12 years of age and over	30 mL (2 tablespoons) every 4 hours
Children under 12 years	Do not use

- When using Day Time and Night Time products, carefully read each label to ensure correct dosing.

#### **Other information**

- each 30 mL contains sodium: **18 mg**
- store between 15-30°C (59-86°F)
- do not refrigerate

#### **Inactive ingredients**

Citric acid, FD&C Yellow # 6, flavor, glycerin, honey, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

#### **Questions or comments?**

1-877-994-3666 Monday to Friday from 8 am to 6 pm, Central Time.

Distributed by  
Genomma Lab USA Inc.,  
Houston, TX 77098

#### **PRINCIPAL DISPLAY PANEL - 118 ml Bottle Carton**

Day Time / Día  
**Tukol**<sup>®</sup>  
 HONEY MULTI-SYMP TOM  
 MIEL MULTI-SÍNTOMA

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING  
 NO SE USE SI EL SELLO IMPRESO DEBAJO DE LA TAPA ESTA ROTO O FALTA

Day Time / Día  
**Tukol**<sup>®</sup>  
 HONEY MULTI-SYMP TOM  
 MIEL MULTI-SÍNTOMA

**COLD & FLU**  
**GRIPE Y RESFRIADO**

Acetaminophen / Dextromethorphan HBr / Phenylephrine HCl  
 Acetaminofén / Dextrometofano HBr / Fenilefrina HCl

Ages / Edades  
**12+**

SABOR A MIEL  
 NATURAL HONEY FLAVOR

Relieves / Alivia:

HEADACHE, SORE THROAT, FEVER / DOLOR DE CABEZA Y GARGANTA, FIEBRE  
 MINOR ACHE AND PAINS / DOLORES Y MOLESTIAS MENORES  
 NASAL CONGESTION AND COUGH / CONGESTIÓN NASAL Y TOS

4 FL OZ (118 mL)

**Drug Facts**

**Active Ingredients (in each 30 mL)**

Acetaminophen, 650 mg.....Pain reliever/Fever reducer  
 Dextromethorphan HBr, 20 mg.....Cough suppressant  
 Phenylephrine HCl, 10 mg.....Nasal decongestant

**Purposes**

**Uses**

Temporarily relieves these common cold and flu symptoms:  
 ■ minor aches and pain ■ headache  
 ■ sore throat ■ nasal congestion  
 ■ cough due to minor throat and bronchial irritation  
 ■ temporarily reduces fever.

**Warnings**

**Liver warning:** this product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product ■ 3 or more alcoholic drinks every day while using this product.  
**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

**Do not use:** ■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**  
 ■ liver disease ■ heart disease ■ thyroid disease ■ diabetes  
 ■ high blood pressure ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma or emphysema ■ cough accompanied by excessive phlegm (mucus).  
**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

**When using this product** ■ do not use more than directed (see overdose warning) ■ avoid alcoholic drinks.  
**Stop use and ask a doctor if**  
 ■ you get nervous, dizzy or sleepless ■ fever gets worse or lasts more than 3 days ■ pain or cough gets worse or lasts more than 5 days (children) or 7 days (adults) ■ redness or swelling is present ■ cough comes back, or occurs with rash or headache that lasts ■ new symptoms occur.  
 These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children.**  
**Overdose warning:** taking more than the recommended dose (overdose) may cause liver damage. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

**Directions**

do not take more than directed ■ do not take more than 4 doses in any 24 hours ■ this adult strength product is not intended for use in children under 12 years of age ■ measure only with dosing cup provided ■ keep dosing cup with product  
 ■ mL = milliliter ■ TBS = tablespoon ■ dose as follows

age	dose
adults and children 12 years of age and over	30 mL (2 tablespoons) every 4 hours
children under 12 years of age	do not use

**Other information**

each (30 mL) contains: sodium 18 mg  
 store between 15-30°C (59-86°F) ■ do not refrigerate.

**Inactive ingredients**

citric acid, FD&C yellow #6, flavor, glycerin, honey, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose.

**Questions or comments?** 1-877-994-3666 Monday to Friday from 8 am to 6 pm, Central Time.

**Información del Medicamento**

**Ingredientes activos Propósitos (en cada 30 mL)**

Acetaminofén, 650 mg.....Analéptico/Antifebril  
 Dextrometofano HBr, 20 mg.....Supresor de la tos  
 Fenilefrina HCl, 10 mg.....Descongestionante nasal

**Usos**

alivia temporalmente los siguientes síntomas que se presentan con la gripe y el resfriado:  
 ■ dolores y molestias leves ■ dolor de cabeza  
 ■ dolor de garganta ■ congestión nasal  
 ■ tos ocasionada por irritación menor de garganta o bronquios  
 ■ reduce temporalmente la fiebre.

**Advertencias**

**Advertencia acerca del hígado:** este producto contiene acetaminofén. Puede producirse daño hepático grave si toma:  
 ■ más de 4 dosis (de 30 mL cada una) en 24 horas, que es la cantidad máxima diaria para este producto ■ con otros medicamentos que contengan acetaminofén ■ 3 o más bebidas alcohólicas diarias mientras usa este producto.

**Alerta sobre alergia:** el acetaminofén puede provocar reacciones severas en la piel. Los síntomas pueden incluir:  
 ■ enrojecimiento de la piel ■ ampollas ■ sarpullido. Si se produce una reacción en la piel, suspender el uso y busque ayuda médica de inmediato.

**Advertencia acerca del dolor de garganta:** si el dolor de garganta es severo, persiste por más de 2 días, está acompañado o seguido de fiebre, dolor de cabeza, sarpullido, náuseas o vómitos, consultar a un médico de inmediato.

**No utilizarlo:** ■ con ningún otro medicamento que contenga acetaminofén (recetado o de venta libre). Si no está seguro si el fármaco contiene acetaminofén, preguntar a un médico o farmacéutico. ■ si está tomando inhibidores de la monoamino oxidasa (MAO) recetados (ciertos medicamentos para

**Información del Medicamento (continuación)**

la depresión, afecciones psiquiátricas o emocionales, o la enfermedad de Parkinson) o durante 2 semanas después de interrumpir el medicamento MAOI. Si no sabe si el medicamento recetado contiene un MAOI, preguntar a un médico o farmacéutico antes de tomar este producto.

**Consulte a su médico antes de usar si usted tiene**  
 ■ enfermedad hepática ■ enfermedad cardíaca  
 ■ enfermedad tiroidea ■ diabetes ■ hipertensión ■ dificultad para orinar debido al agrandamiento de la próstata ■ tos persistente o crónica tal como ocurre con el tabaquismo, el asma o enfisema ■ tos que se presenta con exceso de flema (mucoosidad).

**Consulte a su médico o farmacéutico antes de usar si está tomando el medicamento anticoagulante warfarina.**  
**Cuando use este producto** ■ no usar más de lo indicado (ver advertencia de sobredosis) ■ evite bebidas alcohólicas.

**Suspenda su uso y consulte a su médico si**  
 ■ siente nerviosismo, mareos o insomnio ■ la fiebre empeora o dura más de 3 días ■ el dolor o la tos empeoran, duran más de 5 días (en niños) o 7 días (en adultos) ■ hay enrojecimiento o hinchazón ■ la tos regresa u ocurre con fiebre, sarpullido o dolor de cabeza persistente ■ se presentan nuevos síntomas. Estos podrían ser signos de una afección grave.

**Si está embarazada o lactando,** preguntar a un profesional de la salud antes de usar.  
**Mantener fuera del alcance de los niños.**

**Advertencia de sobredosis:** tomar más de la dosis recomendada (sobredosis) puede causar daño hepático. En caso de sobredosis accidental, buscar ayuda médica o comuníquese con un Centro de Control de Envenenamiento de inmediato. La atención médica inmediata es crítica tanto para adultos como para niños, incluso si no nota algún signo o sintoma.

**Indicaciones**

no tomar más de lo indicado ■ no tome más de 4 dosis en un periodo de 24 horas ■ este producto de concentración para adultos no está diseñado para usarse en niños menores de 12 años de edad ■ medir solo con la copa dosificadora incluida ■ mantenga la copa dosificadora con este producto  
 ■ mL = mililitro ■ TBS = cucharada ■ administrar dosis según se indica continuación

edad	dosis
adultos y niños de 12 años de edad y mayores	30 mL (2 cucharadas) cada 4 horas
niños menores de 12 años de edad	no usar

**Otra información**  
 cada (30 mL) contiene: 18 mg de sodio  
 ■ almacenar entre 15 - 30°C (59-86°F) ■ no refrigerar.

**Ingredientes inactivos**  
 ácido cítrico, FD&C amarillo #6, sabor, glicerina, miel, propilenglicol, agua purificada, benzoato de sodio, solución de sorbitol, sacarina.

**¿Preguntas o comentarios?** 1-877-994-3666  
 Lunes a viernes de 8 am a 6 pm, Hora del Centro.



Distributed by / Distribuido por:  
 Genomma Lab USA Inc.,  
 Houston, TX 77098

2000088888V-2

Rev. 1

101-X1



EXP:

LOT NO./LOTE:

Tukol®

HONEY MULTI SYMPTOM

COLD & FLU

Acetaminophen / Dextromethorphan HBr  
/ Phenylephrine HCl

Ages

12+

NATURAL

HONEY

FLAVOR

Relieves:

- HEADACHE, SORE THROAT, FEVER
- MINOR ACHES AND PAINS
- NASAL CONGESTION AND COUGH

4 FL OZ (118 ml)

## TUKOL HONEY DAYTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride syrup

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50066-302
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Acetaminophen</b> (UNII: 362O9ITL9D) (Acetaminophen - UNII:362O9ITL9D)	Acetaminophen	21.667 mg in 1 mL
<b>Dextromethorphan Hydrobromide</b> (UNII: 9D2RTI9KYH) (Dextromethorphan - UNII:7355X3ROTS)	Dextromethorphan Hydrobromide	0.667 mg in 1 mL
<b>Phenylephrine Hydrochloride</b> (UNII: 04JA59TNSJ) (Phenylephrine - UNII:1WS297W6MV)	Phenylephrine Hydrochloride	0.333 mg in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>Water</b> (UNII: 059QF0KO0R)	
<b>Glycerin</b> (UNII: PDC6A3C0OX)	
<b>Propylene Glycol</b> (UNII: 6DC9Q167V3)	
<b>Sorbitol</b> (UNII: 506T60A25R)	
<b>Sodium benzoate</b> (UNII: OJ245FE5EU)	
<b>Sucralose</b> (UNII: 96K6UQ3ZD4)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	

**HONEY** (UNII: Y9H1V576FH)

### Product Characteristics

<b>Color</b>	YELLOW (Amber to yellow)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50066-302-04	1 in 1 CARTON	02/09/2020	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	02/09/2020	

**Labeler** - Genomma Lab USA (832323534)

Revised: 1/2024

Genomma Lab USA