

**COLD AND FLU NIGHTTIME- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid**  
**QUALITY CHOICE (Chain Drug Marketing Association)**

-----  
**Drug Facts**

**Active ingredients (in each 30 mL)**

**Acetaminophen 650 mg**

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

**Purposes**

**Pain reliever/fever reducer**

Cough suppressant

Antihistamine

**Uses**

- temporarily relieves these common cold/flu symptoms:
  - minor aches and pains
  - headache
  - sore throat
  - fever
  - runny nose and sneezing
  - cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 doses of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is

accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to an enlarged prostate gland

### **Ask a doctor or pharmacist before use if you are taking**

- sedatives or tranquilizers
- the blood thinning drug warfarin

### **When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 7 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in any 24-hours period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL= milliliter
- keep dosing cup with product
- adults and children 12 years and over: 30 mL every 6 hours
- children under 12 years of age: do not use
- **When using other Day Time or Night Time products, carefully read each label to ensure correct dosing**

## Other information

- **each 30 mL contains:** potassium 5 mg
- **each 30 mL contains:** Sodium 19 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

## Inactive ingredients

acesulfame potassium, alcohol, citric acid, D&C yellow #10, FD&C green #3, FD&C yellow #6,, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

## Questions or comments?

Call **1-248-449-9300 Monday-Friday 9AM-5PM EST**

## Principal Display Panel

\*Compare to the active ingredients in VICKS® NYQUIL® Cold & Flu

## Nighttime Cold & Flu

### Acetaminophen

Dextromethorphan

Doxylamine succinate

For Relief of:

Aches | Fever | Cough

Runny Nose & Sneezing

For Ages 12 Years & Over

Alcohol 10 %

Original Flavor

FL OZ (mL)

\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and NyQuil® are registered trademarks of The Procter & Gamble Company.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.**

Distributed by C.D.M.A., Inc.©

43157 W. 9 Mile Rd

Novi, MI 48376-0995

www.qualitychoice.com

**Product Label**

**Drug Facts (continued)**

**Stop use and ask a doctor if** ■ pain or cough gets worse or lasts more than 7 days  
 ■ new symptoms occur ■ fever gets worse or lasts more than 3 days  
 ■ redness or swelling is present ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions** ■ do not take more than directed (see Overdose warning)  
 ■ do not take more than 4 doses in any 24-hour period  
 ■ measure only with dosing cup provided. Do not use any other dosing device.  
 ■ mL = milliliter ■ keep dosing cup with product ■ adults and children 12 years and over: 30 mL every 6 hours ■ children under 12 years of age: do not use  
 ■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

**Other information** ■ each 30 mL contains: potassium 5 mg  
 ■ each 30 mL contains: sodium 19 mg  
 ■ store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** acesulfame potassium, alcohol, citric acid, D&C yellow #10, FD&C green #3, FD&C yellow #6, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

**Questions or comments?**  
 Call 1-248-449-9300 Monday-Friday 9AM-5PM EST

\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and NyQuil® are registered trademarks of The Procter & Gamble Company.

Distributed by C.D.M.A., Inc.©  
 43157 W 9 Mile Rd  
 Novi, MI 48376  
 www.qualitychoice.com  
 Questions: 248-449-9300

PLD-8407B LB004094

**PARENTS:**  
Learn how to use medicine safely  
 www.StopTheCircus.com

6 35515 95332 1

NDC 63868-140-12

**QC QUALITY CHOICE**

\*Compare to the active ingredients in VICKS® NYQUIL® Cold & Flu

# Nighttime

## Cold & Flu

**Acetaminophen**  
**Dextromethorphan HBr**  
**Doxylamine Succinate**

**For Relief of:**  
 Aches | Fever | Cough  
 Runny Nose & Sneezing

For Ages 12 Years & Over  
 Alcohol 10%

**Original Flavor**



**12 FL OZ (355 mL)**

**Drug Facts** TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

**Active ingredients (in each 30 mL)**

Acetaminophen 650 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg.....	Cough suppressant
Doxylamine succinate 12.5 mg.....	Antihistamine

**Uses** ■ temporarily relieves common cold/flu symptoms  
 ■ minor aches and pains ■ headache ■ fever ■ sore throat  
 ■ runny nose and sneezing ■ cough due to minor throat and bronchial irritation

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours  
 ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma  
 ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ trouble urinating due to enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**  
 ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin

**When using this product**  
 ■ excitability may occur, especially in children ■ marked drowsiness may occur  
 ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness

**QUALITY CHOICE Nighttime Cold & Flu**

**COLD AND FLU NIGHTTIME**

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63868-140
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL

<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>HIGH FRUCTOSE CORN SYRUP</b> (UNII: XY6UN3QB6S)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C GREEN NO. 3</b> (UNII: 3P3ONR6O1S)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-140-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/31/2016	
2	NDC:63868-140-12	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/31/2016	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/31/2016	

**Labeler** - QUALITY CHOICE (Chain Drug Marketing Association) (011920774)

Revised: 5/2024

QUALITY CHOICE (Chain Drug Marketing Association)