

**DG HEALTH ANTI DIARRHEAL- loperamide hydrochloride tablet, film coated
Dolgenercorp, LLC**

Dolgenercorp, LLC Anti-Diarrheal Drug Facts

Active ingredient (in each caplet)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- store at 20° -25° C (68° -77° F)
- see end panel for lot number and expiration date

Inactive ingredients

anhydrous lactose, carnauba wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

Questions or comments?

1-888-309-9030

Principal Display Panel

SEE NEW WARNINGS

Compare to the active ingredient of Imodium® A-D

Loperamide Hydrochloride Tablets, 2 mg

Anti-Diarrheal

Controls the symptoms of diarrhea

Anti-Diarrheal

12 Caplets*

*Capsule-Shaped Tablets

Actual Caplet Size

2 mg

OPEN OTHER END
CODE AREA

22453 VT G1

Drug Facts (continued)
 Inactive ingredients: anhydrous lactose, croscarmellose, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch.
 100% Satisfaction Guarantee (BBB) 309-9030
 Questions or comments? 1-888-309-9030
Drug Facts (continued)

Drug Facts (continued)
Warnings
 Do not use if you have ever had a severe allergic reaction to loperamide HCl. Heart at risk: Taking more than directed can cause serious heart problems or death. Do not use if you have bloody or black stools. Ask a doctor before use if you have:
 ■ a history of abnormal heart rhythm
 ■ a history of liver disease
 ■ fever
 ■ muscle pain
 Ask a doctor before use if you have:
 ■ a history of abnormal heart rhythm
 ■ a history of liver disease
 ■ fever
 ■ muscle pain
 If you are taking a prescription drug, ask a doctor or pharmacist before use. Loperamide may interact with certain prescription drugs.
 When using this product, the drowsiness or dizziness may occur. Be careful when driving or operating machinery.
 Stop use and ask a doctor if:
 ■ symptoms get worse
 ■ diarrhea lasts for more than 2 days
 ■ you get abdominal swelling or bloating.
 These may be signs of a serious condition. If pregnant or breastfeeding, ask a health professional before use.

Directions
 Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).
 Adults and children 12 years and over: 2 caplets after the first loose stool, 1 caplet thereafter each subsequent loose stool, but no more than 4 caplets in 24 hours.
 Children 9-11 years: 1 caplet after the first loose stool, 1/2 caplet thereafter each subsequent loose stool, but no more than 3 caplets in 24 hours.
 Children 6-8 years: 1 caplet after the first loose stool, 1/2 caplet thereafter each subsequent loose stool, but no more than 2 caplets in 24 hours.
 Children 2-5 years: ask a doctor.
 Children under 2 years (up to 33 lbs): do not use.

Other information
 ■ Store at 20°-25°C (68°-77°F).
 ■ See expiration date for lot number and expiration date.

Drug Facts (continued)

3 7003016946 7
 A1378

DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN
 DISTRIBUTED BY: QLD EAST MAIN CO., 100 MISSION RIDGE, GOODLETTSVILLE, TN 37072
 *This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Imodium® A-D.

Loperamide Hydrochloride Tablets, 2 mg

Anti-Diarrheal



SEE NEW WARNINGS
 Compare to the active ingredient of Imodium® A-D**

Loperamide Hydrochloride Tablets, 2 mg

Anti-Diarrheal

- Controls the symptoms of diarrhea
- Anti-Diarrheal

2 mg



12 Caplets*

Actual Caplet Size *Capsule-Shaped Tablets



DG HEALTH ANTI DIARRHEAL

loperamide hydrochloride tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-224
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	

Product Characteristics

Color	GREEN	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	L2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-224-67	1 in 1 CARTON	12/28/2009	01/31/2022
1		48 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:55910-224-23	1 in 1 CARTON	09/11/2012	12/31/2021
2		72 in 1 BOTTLE; Type 0: Not a Combination Product		

3	NDC:55910-224-72	1 in 1 CARTON	05/02/2017	07/31/2021
3		60 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:55910-224-62	24 in 1 CARTON	02/14/2020	
4		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
5	NDC:55910-224-53	12 in 1 CARTON	07/06/2020	
5		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
6	NDC:55910-224-91	6 in 1 CARTON	07/08/2020	08/31/2023
6		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075232	12/28/2009	

Labeler - Dolgencorp, LLC (068331990)

Revised: 9/2023

Dolgencorp, LLC