

**DG HEALTH ANTI DIARRHEAL- loperamide hydrochloride tablet, film coated
Dolgenercorp, LLC**

Dolgenercorp, LLC Anti-Diarrheal Drug Facts

Active ingredient (in each caplet)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Do not use

if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease

Ask a doctor or pharmacist before use if you are
taking antibiotics

When using this product

- tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging.

These may be signs of a serious condition.

If pregnant or breast-feeding,
ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor

Other information

- store at 20°-25°C (68°-77°F)
- do not use if printed foil under cap is broken or missing
- see end panel for lot number and expiration date

Inactive ingredients

anhydrous lactose, carnauba wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

Questions or comments?

1-888-309-9030

Principal Display Panel

Compare to active ingredient of Imodium® A-D

Loperamide Hydrochloride Tablets, 2 mg

Anti-Diarrheal

Controls the Symptoms of Diarrhea

48 Caplets*

2 mg

Actual Caplet Size

*Each caplet (capsule shaped tablet) contains 2 mg loperamide hydrochloride

22447 VT C 4

EXP.

LOT NO.



Drug Facts

Active ingredient (in each caplet) Loperamide HCl 2 mg.....Anti-Diarrheal

Use controls symptoms of diarrhea, including Traveler's Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl
Do not use if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease

Ask a doctor or pharmacist before use if you are taking antibiotics

When using this product

- tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.
- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging.
- These may be signs of a serious condition.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging.
- These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Drug Facts (continued)

Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- take right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over: 2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours

children 9-11 years: 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours

children 6-8 years: 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours

children under 6 years (up to 47 lbs): ask a doctor

Other information

- store at 20°-25°C (68°-77°F)
- do not use if printed foil under cap is broken or missing
- see end panel for lot number and expiration date

Inactive ingredients anhydrous lactose, carnauba wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

Drug Facts (continued)

(Questions or comments?)

1-888-309-9030

*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Imodium® A-D.
DISTRIBUTED BY DOLGENCORP, LLC
100 MISSION RIDGE
GOODLETTSVILLE, TN 37072 USA

100% Satisfaction Guaranteed! (888) 309-9030

ACTUAL SIZE



Compare to active ingredient of Imodium® A-D**



Loperamide Hydrochloride Tablets, 2 mg

Anti-Diarrheal

Controls the Symptoms of Diarrhea



48 Caplets*

2 mg

Actual Caplet Size

*Each caplet (capsule shaped tablet) contains 2 mg loperamide hydrochloride

THIS LABEL DOES NOT CONTAIN FULL PRODUCT

DG HEALTH ANTI DIARRHEAL

loperamide hydrochloride tablet, film coated

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:55910-224

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77T135393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6B30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	

Product Characteristics

Color	GREEN	Score	2 pieces
Shape	CAPSULE	Size	10mm
Flavor		Imprint Code	L2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:559 10-224-67	1 in 1 CARTON	12/28/2009	
1		48 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:559 10-224-23	1 in 1 CARTON	09/11/2012	
2		72 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:559 10-224-72	1 in 1 CARTON	05/02/2017	
3		60 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075232	12/28/2009	

Labeler - Dolgencorp, LLC (068331990)