INFLUENZINUM- influenza a virus a/victoria/2570/2019 ivr-215 (h1n1) hemagglutinin antigen (formaldehyde inactivated), influenza a virus a/tasmania/503/2020 ivr-221 (h3n2) hemagglutinin antigen (formaldehyde inactivated), influenza b virus b/washington/02/2019 hemagglutinin antigen (formaldehyde inactivated), influenza b virus b/phuket/3073/2013 hemagglutinin antigen (formaldehyde inactivated) pellet Boiron

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

INFLUENZINUM 6C

INFLUENZINUM 6C HPUS**

(** contains 0.443 mg of the active ingredient per pellet)

INFLUENZA A VIRUS A/VICTORIA/2570/2019 IVR-215 (H1N1) HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED

INFLUENZA A VIRUS A/TASMANIA/503/2020 IVR-221 (H3N2) HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED

INFLUENZA B VIRUS B/WASHINGTON/02/2019 HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED)

INFLUENZA B VIRUS B/PHUKET/3073/2013 HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED)

After effects of flu or flu-like symptoms*

Stop use and ask a doctor if symptoms persist for more than 3 days or worsen.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children.

Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.

Do not use if pellet dispenser seal is broken.

Contains approx 80 pellets.

How to dispense pellets? Turn tube upside down. Twist until 5 pellets are dispensed into cap. Carefully remove the cap and use it to pour pellets under the tongue.

*CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE NOT ACCEPTED MEDICAL EVIDENCE. NOT FDA EVALUATED.

*C,K,CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.

lactose, sucrose

Made in France

1-800-BOIRON-1 (1-800-264-7661), BoironUSA.com Info@boiron.com Distributed by Boiron, Inc. Newtown Square, PA 19073



Drug Facts

Active ingredient**: See product name on front panel (contains 0.443 mg of the active ingredient per pellet).

Uses: See symptoms on front panel.

Warnings: Stop use and ask a doctor if symptoms persist for more than 3 days or worsen. If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children.

Directions: Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a



INFLUENZINUM

influenza a virus a/victoria/2570/2019 ivr-215 (h1n1) hemagglutinin antigen (formaldehyde inactivated), influenza a virus a/tasmania/503/2020 ivr-221 (h3n2) hemagglutinin antigen (formaldehyde inactivated), influenza b virus b/washington/02/2019 hemagglutinin antigen (formaldehyde inactivated), influenza b virus b/phuket/3073/2013 hemagglutinin antigen (formaldehyde inactivated) pellet

Product Information

Product Type

HUMAN OTC DRUG

ORAL

Item Code (Source)

NDC:0220-2718

Route of Administration

| Active Ingredient/Active Moiety | | | | | | | | |
|---|--|--|--|-------------------------|-----------------------|-------------------------------------|--|--|
| | Ingre | Basis of Stre | Basis of Strength | | | | | |
| Ingredient NameBasis of StrINFLUENZA A VIRUS A/VICTORIA/2570/2019 IVR-215 (H1N1)INFLUENZA A VIRUSHEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) (UNII: ERB9SJ9VMQ) (INFLUENZA A VIRUS A/VICTORIA/2570/2019 IVR-215 (H1N1)A/VICTORIA/2570/2010HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) - UNII: ERB9SJ9VMQ)ANTIGEN (FORMALDEHYDE INACTIVATED) - INACTIVATED)ANTIGEN (FORMALD | | | | | | Strength 6 [hp_C] in 6 [hp_C] | | |
| INFLUENZA B VI ANTIGEN (FORM (INFLUENZA B VIR (FORMALDEHYDE | ALDEHYDE INA US B/WASHINGT | N HEMAGGLUTININ AN | INFLUENZA B VIRUS B/WASHINGTON/02/2019 HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) | | | | | |
| INFLUENZA A VI HEMAGGLUTINII CYN792B8TR) (INI HEMAGGLUTININ / UNII:CYN792B8TR | 2) (H3N2) HEMAGGLUT | A/TASMANIA/503/2020 IVR-221 (H3N2) HEMAGGLUTININ ANTIGEN (FORMALDEHYDE | | | | | | |
| INFLUENZA B VIRUS B/PHUKET/3073/2013 HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) (UNII: 9HB0XUS9TM)INFLUENZA B VIRUS B/PHUKET/3073/2013 HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) - UNII:9HB0XUS9TM)INFLUENZA B VIRUS B/PHUKET/3073/2013 HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) - UNII:9HB0XUS9TM) | | | | | | 6 [hp_C] in 6 [hp_C] | | |
| | | | | | | | | |
| Inactive Ingredients | | | | | | | | |
| Ingredient Name | | | | | | Strength | | |
| SUCROSE (UNII: C151H8M554) | | | | | | | | |
| LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G) | | | | | | | | |
| | | | | | | | | |
| Product Cha | racteristics | 5 | | | | | | |
| Color | | white | Score | | | | | |
| Shape | | ROUND | Size | | 4mm | | | |
| Flavor | | | Imprint Code | | | | | |
| Contains | | | | | | | | |
| Packaging | | | | | | | | |
| # Item Code | P | Package Description | | Marketing Start Date | Marketing End Date | | | |
| 1 NDC:0220-271 41 | NDC:0220-2718- 41 6 [hp_C] in 1 TUBE; Type 0: Not a Combination Product 10/01/2021 | | | | | | | |
| | | _ | | | | | | |
| Marketing Information | | | | | | | | |
| Marketing | Informa | tion | | | | | | |
| Marketing Marketing Category | | TION ation Number or N Citation | Monograph | Marketing Start Date | | ting End ate | | |

Registrant - Boiron, Inc. (014892269)

| Establishment | | | | | | | |
|---------------|---------|-----------|----------------------------|--|--|--|--|
| Name | Address | ID/FEI | Business Operations | | | | |
| Boiron | | 282560473 | manufacture(0220-2718) | | | | |

Revised: 8/2021

Boiron