


MODAFINIL- modafinil tablet
Aphena Pharma Solutions - Tennessee, LLC

Modafinil Tablets

DESCRIPTION

Modafinil is a wakefulness-promoting agent for oral administration. Modafinil is a racemic compound. The chemical name for modafinil is 2-[(diphenylmethyl)sulfinyl]acetamide. The molecular formula is C₁₅H₁₅NO₂S and the molecular weight is 273.35.

The chemical structure is:



Modafinil is a white to off-white, crystalline powder that is practically insoluble in water and cyclohexane. It is sparingly to slightly soluble in methanol and acetone. Modafinil tablets contain 100 mg or 200 mg of modafinil and the following inactive ingredients: lactose monohydrate, microcrystalline cellulose, pregelatinized starch, croscarmellose sodium, povidone, and magnesium stearate.

CLINICAL PHARMACOLOGY

The precise mechanism(s) through which modafinil promotes wakefulness is unknown. Modafinil has wake-promoting actions similar to sympathomimetic agents like amphetamine and methylphenidate, although the pharmacologic profile is not identical to that of sympathomimetic amines.

Modafinil has weak to negligible interactions with receptors for norepinephrine, serotonin, dopamine, GABA, adenosine, histamine-3, melatonin, and benzodiazepines. Modafinil also does not inhibit the activities of MAO-B or phosphodiesterases II-V.

Modafinil-induced wakefulness can be attenuated by the α_1 -adrenergic receptor antagonist prazosin; however, modafinil is inactive in other in vitro assay systems known to be responsive to α -adrenergic agonists, such as the rat vas deferens preparation.

Modafinil is not a direct- or indirect-acting dopamine receptor agonist. However, in vitro, modafinil binds to the dopamine transporter and inhibits dopamine reuptake. This activity has been associated in vivo with increased extracellular dopamine levels in some brain regions of animals. In genetically engineered mice lacking the dopamine transporter (DAT), modafinil lacked wake-promoting activity, suggesting that this activity was DAT-dependent. However, the wake-promoting effects of modafinil, unlike those of amphetamine, were not antagonized by the dopamine receptor antagonist haloperidol in rats. In addition, alpha-methyl-p-tyrosine, a dopamine synthesis inhibitor, blocks the action of amphetamine, but does not block locomotor activity induced by modafinil.

In the cat, equal wakefulness-promoting doses of methylphenidate and amphetamine increased neuronal activation throughout the brain. Modafinil at an equivalent wakefulness-promoting dose selectively and prominently increased neuronal activation in more discrete regions of the brain. The relationship of this

finding in cats to the effects of modafinil in humans is unknown.

In addition to its wake-promoting effects and ability to increase locomotor activity in animals, modafinil produces psychoactive and euphoric effects, alterations in mood, perception, thinking, and feelings typical of other CNS stimulants in humans. Modafinil has reinforcing properties, as evidenced by its self-administration in monkeys previously trained to self-administer cocaine. Modafinil was also partially discriminated as stimulant-like.

The optical enantiomers of modafinil have similar pharmacological actions in animals. Two major metabolites of modafinil, modafinil acid and modafinil sulfone, do not appear to contribute to the CNS-activating properties of modafinil.

Pharmacokinetics

Modafinil is a racemic compound, whose enantiomers have different pharmacokinetics (e.g., the half-life of the *l*-isomer is approximately three times that of the *d*-isomer in adult humans). The enantiomers do not interconvert. At steady state, total exposure to the *l*-isomer is approximately three times that for the *d*-isomer. The trough concentration ($C_{\text{min,ss}}$) of circulating modafinil after once daily dosing consists of 90% of the *l*-isomer and 10% of the *d*-isomer. The effective elimination half-life of modafinil after multiple doses is about 15 hours. The enantiomers of modafinil exhibit linear kinetics upon multiple dosing of 200-600 mg/day once daily in healthy volunteers. Apparent steady states of total modafinil and *l*(-)-modafinil are reached after 2-4 days of dosing.

Absorption

Absorption of modafinil tablets is rapid, with peak plasma concentrations occurring at 2-4 hours. The bioavailability of modafinil tablets is approximately equal to that of an aqueous suspension. The absolute oral bioavailability was not determined due to the aqueous insolubility (<1 mg/mL) of modafinil, which precluded intravenous administration. Food has no effect on overall modafinil bioavailability; however, its absorption (t_{max}) may be delayed by approximately one hour if taken with food.

Distribution

Modafinil is well distributed in body tissue with an apparent volume of distribution (~0.9 L/kg) larger than the volume of total body water (0.6 L/kg). In human plasma, in vitro, modafinil is moderately bound to plasma protein (~60%, mainly to albumin). At serum concentrations obtained at steady state after doses of 200 mg/day, modafinil exhibits no displacement of protein binding of warfarin, diazepam or propranolol. Even at much larger concentrations (1000 μM ; > 25 times the C_{max} of 40 μM at steady state at 400 mg/day), modafinil has no effect on warfarin binding. Modafinil acid at concentrations >500 μM decreases the extent of warfarin binding, but these concentrations are >35 times those achieved therapeutically.

Metabolism and Elimination

The major route of elimination is metabolism (~90%), primarily by the liver, with subsequent renal elimination of the metabolites. Urine alkalization has no effect on the elimination of modafinil.

Metabolism occurs through hydrolytic deamidation, S-oxidation, aromatic ring hydroxylation, and glucuronide conjugation. Less than 10% of an administered dose is excreted as the parent compound. In a clinical study using radiolabeled modafinil, a total of 81% of the administered radioactivity was recovered in 11 days post-dose, predominantly in the urine (80% vs. 1.0% in the feces). The largest fraction of the drug in urine was modafinil acid, but at least six other metabolites were present in lower concentrations. Only two metabolites reach appreciable concentrations in plasma, i.e., modafinil acid and modafinil sulfone. In preclinical models, modafinil acid, modafinil sulfone, 2-[(diphenylmethyl)sulfonyl]acetic acid and 4-hydroxy modafinil, were inactive or did not appear to mediate the arousal effects of modafinil.

In adults, decreases in trough levels of modafinil have sometimes been observed after multiple weeks

of dosing, suggesting auto-induction, but the magnitude of the decreases and the inconsistency of their occurrence suggest that their clinical significance is minimal. Significant accumulation of modafinil sulfone has been observed after multiple doses due to its long elimination half-life of 40 hours. Induction of metabolizing enzymes, most importantly cytochrome P-450 (CYP) 3A4, has also been observed in vitro after incubation of primary cultures of human hepatocytes with modafinil and in vivo after extended administration of modafinil at 400 mg/day. (For further discussion of the effects of modafinil on CYP enzyme activities, see **PRECAUTIONS, Drug Interactions**.)

Drug-Drug Interactions: Based on in vitro data, modafinil is metabolized partially by the 3A isoform subfamily of hepatic cytochrome P450 (CYP3A4). In addition, modafinil has the potential to inhibit CYP2C19, suppress CYP2C9, and induce CYP3A4, CYP2B6, and CYP1A2. Because modafinil and modafinil sulfone are reversible inhibitors of the drug-metabolizing enzyme CYP2C19, co-administration of modafinil with drugs such as diazepam, phenytoin and propranolol, which are largely eliminated via that pathway, may increase the circulating levels of those compounds. In addition, in individuals deficient in the enzyme CYP2D6 (i.e., 7-10% of the Caucasian population; similar or lower in other populations), the levels of CYP2D6 substrates such as tricyclic antidepressants and selective serotonin reuptake inhibitors, which have ancillary routes of elimination through CYP2C19, may be increased by co-administration of modafinil. Dose adjustments may be necessary for patients being treated with these and similar medications (See **PRECAUTIONS, Drug Interactions**). An in vitro study demonstrated that armodafinil (one of the enantiomers of modafinil) is a substrate of P-glycoprotein.

Coadministration of modafinil with other CNS active drugs such as methylphenidate and dextroamphetamine did not significantly alter the pharmacokinetics of either drug.

Chronic administration of modafinil 400 mg was found to decrease the systemic exposure to two CYP3A4 substrates, ethinyl estradiol and triazolam, after oral administration suggesting that CYP3A4 had been induced. Chronic administration of modafinil can increase the elimination of substrates of CYP3A4. Dose adjustments may be necessary for patients being treated with these and similar medications (See **PRECAUTIONS, Drug Interactions**).

An apparent concentration-related suppression of CYP2C9 activity was observed in human hepatocytes after exposure to modafinil in vitro suggesting that there is a potential for a metabolic interaction between modafinil and the substrates of this enzyme (e.g., S-warfarin, phenytoin). However, in an interaction study in healthy volunteers, chronic modafinil treatment did not show a significant effect on the pharmacokinetics of warfarin when compared to placebo. (See **PRECAUTIONS, Drug Interactions, Other Drugs, Warfarin**).

Special Populations

Gender Effect: The pharmacokinetics of modafinil are not affected by gender.

Age Effect: A slight decrease (~20%) in the oral clearance (CL/F) of modafinil was observed in a single dose study at 200 mg in 12 subjects with a mean age of 63 years (range 53 – 72 years), but the change was considered not likely to be clinically significant. In a multiple dose study (300 mg/day) in 12 patients with a mean age of 82 years (range 67 – 87 years), the mean levels of modafinil in plasma were approximately two times those historically obtained in matched younger subjects. Due to potential effects from the multiple concomitant medications with which most of the patients were being treated, the apparent difference in modafinil pharmacokinetics may not be attributable solely to the effects of aging. However, the results suggest that the clearance of modafinil may be reduced in the elderly (See **DOSAGE AND ADMINISTRATION**).

Race Effect: The influence of race on the pharmacokinetics of modafinil has not been studied.

Renal Impairment: In a single dose 200 mg modafinil study, severe chronic renal failure (creatinine clearance \leq 20 mL/min) did not significantly influence the pharmacokinetics of modafinil, but exposure to modafinil acid (an inactive metabolite) was increased 9-fold (See **PRECAUTIONS**).

Hepatic Impairment: Pharmacokinetics and metabolism were examined in patients with cirrhosis of the liver (6 males and 3 females). Three patients had stage B or B+ cirrhosis (per the Child criteria) and 6 patients had stage C or C+ cirrhosis. Clinically 8 of 9 patients were icteric and all had ascites. In these patients, the oral clearance of modafinil was decreased by about 60% and the steady state concentration was doubled compared to normal patients. The dose of modafinil should be reduced in patients with severe hepatic impairment (See **PRECAUTIONS** and **DOSAGE AND ADMINISTRATION**).

CLINICAL TRIALS

The effectiveness of modafinil in reducing excessive sleepiness has been established in the following sleep disorders: narcolepsy, obstructive sleep apnea (OSA), and shift work disorder (SWD).

Narcolepsy

The effectiveness of modafinil in reducing the excessive sleepiness (ES) associated with narcolepsy was established in two US 9-week, multicenter, placebo-controlled, two-dose (200 mg per day and 400 mg per day) parallel-group, double-blind studies of outpatients who met the ICD-9 and American Sleep Disorders Association criteria for narcolepsy (which are also consistent with the American Psychiatric Association DSM-IV criteria). These criteria include either 1) recurrent daytime naps or lapses into sleep that occur almost daily for at least three months, plus sudden bilateral loss of postural muscle tone in association with intense emotion (cataplexy) or 2) a complaint of excessive sleepiness or sudden muscle weakness with associated features: sleep paralysis, hypnagogic hallucinations, automatic behaviors, disrupted major sleep episode; and polysomnography demonstrating one of the following: sleep latency less than 10 minutes or rapid eye movement (REM) sleep latency less than 20 minutes. In addition, for entry into these studies, all patients were required to have objectively documented excessive daytime sleepiness, a Multiple Sleep Latency Test (MSLT) with two or more sleep onset REM periods, and the absence of any other clinically significant active medical or psychiatric disorder. The MSLT, an objective daytime polysomnographic assessment of the patient's ability to fall asleep in an unstimulating environment, measures latency (in minutes) to sleep onset averaged over 4 test sessions at 2-hour intervals following nocturnal polysomnography. For each test session, the subject was told to lie quietly and attempt to sleep. Each test session was terminated after 20 minutes if no sleep occurred or 15 minutes after sleep onset.

In both studies, the primary measures of effectiveness were 1) sleep latency, as assessed by the Maintenance of Wakefulness Test (MWT) and 2) the change in the patient's overall disease status, as measured by the Clinical Global Impression of Change (CGI-C). For a successful trial, both measures had to show significant improvement.

The MWT measures latency (in minutes) to sleep onset averaged over 4 test sessions at 2 hour intervals following nocturnal polysomnography. For each test session, the subject was asked to attempt to remain awake without using extraordinary measures. Each test session was terminated after 20 minutes if no sleep occurred or 10 minutes after sleep onset. The CGI-C is a 7-point scale, centered at *No Change*, and ranging from *Very Much Worse* to *Very Much Improved*. Patients were rated by evaluators who had no access to any data about the patients other than a measure of their baseline severity. Evaluators were not given any specific guidance about the criteria they were to apply when rating patients.

Other assessments of effect included the Multiple Sleep Latency Test (MSLT), Epworth Sleepiness Scale (ESS; a series of questions designed to assess the degree of sleepiness in everyday situations), the Steer Clear Performance Test (SCPT; a computer-based evaluation of a patient's ability to avoid hitting obstacles in a simulated driving situation), standard nocturnal polysomnography, and patient's daily sleep log. Patients were also assessed with the Quality of Life in Narcolepsy (QOLIN) scale, which contains the validated SF-36 health questionnaire.

Both studies demonstrated improvement in objective and subjective measures of excessive daytime sleepiness for both the 200 mg and 400 mg doses compared to placebo. Patients treated with either dose of modafinil showed a statistically significantly enhanced ability to remain awake on the MWT (all

p values <0.001) at weeks 3, 6, 9, and final visit compared to placebo and a statistically significantly greater global improvement, as rated on the CGI-C scale (all p values <0.05).

The average sleep latencies (in minutes) on the MWT at baseline for the 2 controlled trials are shown in Table 1 below, along with the average change from baseline on the MWT at final visit.

The percentages of patients who showed any degree of improvement on the CGI-C in the two clinical trials are shown in Table 2 below.

Similar statistically significant treatment-related improvements were seen on other measures of impairment in narcolepsy, including a patient assessed level of daytime sleepiness on the ESS (p<0.001 for each dose in comparison to placebo).

Nighttime sleep measured with polysomnography was not affected by the use of modafinil.

Obstructive Sleep Apnea (OSA)

The effectiveness of modafinil in reducing the excessive sleepiness associated with OSA was established in two clinical trials. In both studies, patients were enrolled who met the International Classification of Sleep Disorders (ICSD) criteria for OSA (which are also consistent with the American Psychiatric Association DSM-IV criteria). These criteria include either, 1) excessive sleepiness or insomnia, plus frequent episodes of impaired breathing during sleep, and associated features such as loud snoring, morning headaches and dry mouth upon awakening; or 2) excessive sleepiness or insomnia and polysomnography demonstrating one of the following: more than five obstructive apneas, each greater than 10 seconds in duration, per hour of sleep and one or more of the following: frequent arousals from sleep associated with the apneas, bradycardia, and arterial oxygen desaturation in association with the apneas. In addition, for entry into these studies, all patients were required to have excessive sleepiness as demonstrated by a score ≥ 10 on the Epworth Sleepiness Scale, despite treatment with continuous positive airway pressure (CPAP). Evidence that CPAP was effective in reducing episodes of apnea/hypopnea was required along with documentation of CPAP use.

In the first study, a 12-week multicenter placebo-controlled trial, a total of 327 patients were randomized to receive modafinil 200 mg/day, modafinil 400 mg/day, or matching placebo. The majority of patients (80%) were fully compliant with CPAP, defined as CPAP use > 4 hours/night on > 70% nights. The remainder were partially CPAP compliant, defined as CPAP use < 4 hours/night on >30% nights. CPAP use continued throughout the study. The primary measures of effectiveness were 1) sleep latency, as assessed by the Maintenance of Wakefulness Test (MWT) and 2) the change in the patient's overall disease status, as measured by the Clinical Global Impression of Change (CGI-C) at week 12 or the final visit. (See **CLINICAL TRIALS**, *Narcolepsy* section above for a description of these tests.)

Patients treated with modafinil showed a statistically significant improvement in the ability to remain awake compared to placebo-treated patients as measured by the MWT (p<0.001) at endpoint [Table 1]. modafinil -treated patients also showed a statistically significant improvement in clinical condition as rated by the CGI-C scale (p<0.001) [Table 2]. The two doses of modafinil performed similarly.

In the second study, a 4-week multicenter placebo-controlled trial, 157 patients were randomized to either modafinil 400 mg/day or placebo. Documentation of regular CPAP use (at least 4 hours/night on 70% of nights) was required for all patients. The primary outcome measure was the change from baseline on the ESS at week 4 or final visit. The baseline ESS scores for the modafinil and placebo groups were 14.2 and 14.4, respectively. At week 4, the ESS was reduced by 4.6 in the modafinil group and by 2.0 in the placebo group, a difference that was statistically significant (p<0.0001).

Nighttime sleep measured with polysomnography was not affected by the use of modafinil.

Shift Work Disorder (SWD)

The effectiveness of modafinil for the excessive sleepiness associated with SWD was demonstrated in a 12-week placebo-controlled clinical trial. A total of 209 patients with chronic SWD were randomized to receive modafinil 200 mg/day or placebo. All patients met the International Classification of Sleep

Disorders (ICSD-10) criteria for chronic SWD (which are consistent with the American Psychiatric Association DSM-IV criteria for Circadian Rhythm Sleep Disorder: Shift Work Type). These criteria include 1) either: a) a primary complaint of excessive sleepiness or insomnia which is temporally associated with a work period (usually night work) that occurs during the habitual sleep phase, or b) polysomnography and the MSLT demonstrate loss of a normal sleep-wake pattern (i.e., disturbed chronobiological rhythmicity); and 2) no other medical or mental disorder accounts for the symptoms, and 3) the symptoms do not meet criteria for any other sleep disorder producing insomnia or excessive sleepiness (e.g., time zone change [jet lag] syndrome).

It should be noted that not all patients with a complaint of sleepiness who are also engaged in shift work meet the criteria for the diagnosis of SWD. In the clinical trial, only patients who were symptomatic for at least 3 months were enrolled.

Enrolled patients were also required to work a minimum of 5 night shifts per month, have excessive sleepiness at the time of their night shifts (MSLT score < 6 minutes), and have daytime insomnia documented by a daytime polysomnogram (PSG).

The primary measures of effectiveness were 1) sleep latency, as assessed by the Multiple Sleep Latency Test (MSLT) performed during a simulated night shift at week 12 or the final visit and 2) the change in the patient's overall disease status, as measured by the Clinical Global Impression of Change (CGI-C) at week 12 or the final visit. Patients treated with modafinil showed a statistically significant prolongation in the time to sleep onset compared to placebo-treated patients, as measured by the nighttime MSLT [Table 1] ($p < 0.05$). Improvement on the CGI-C was also observed to be statistically significant ($p < 0.001$). (See **CLINICAL TRIALS**, *Narcolepsy* section above for a description of these tests.)

Daytime sleep measured with polysomnography was not affected by the use of modafinil.

Table 1. Average Baseline Sleep Latency and Change from Baseline at Final Visit in Adults (MWT and MSLT in minutes)

Disorder	Measure	Modafinil 200 mg*		Modafinil 400 mg*		Placebo	
		Baseline	Change from Baseline	Baseline	Change from Baseline	Baseline	Change from Baseline
Narcolepsy I	MWT	5.8	2.3	6.6	2.3	5.8	-0.7
Narcolepsy II	MWT	6.1	2.2	5.9	2.0	6.0	-0.7
OSA	MWT	13.1	1.6	13.6	1.5	13.8	-1.1
SWD	MSLT	2.1	1.7	-	-	2.0	0.3

*Significantly different than placebo for all trials ($p < 0.01$ for all trials but SWD, which was $p < 0.05$)

Table 2. Clinical Global Impression of Change (CGI-C) (Percent of Adult Patients Who Improved at Final Visit)

Disorder	Modafinil 200 mg*	Modafinil 400 mg*	Placebo
Narcolepsy I	64%	72%	37%
Narcolepsy II	58%	60%	38%
OSA	61%	68%	37%
SWD	74%	-----	36%

*Significantly different than placebo for all trials (p<0.01)

INDICATIONS AND USAGE

Modafinil is indicated to improve wakefulness in adult patients with excessive sleepiness associated with narcolepsy, obstructive sleep apnea, and shift work disorder.

In OSA, modafinil is indicated as an adjunct to standard treatment(s) for the underlying obstruction. If continuous positive airway pressure (CPAP) is the treatment of choice for a patient, a maximal effort to treat with CPAP for an adequate period of time should be made prior to initiating modafinil. If modafinil is used adjunctively with CPAP, the encouragement of and periodic assessment of CPAP compliance is necessary.

In all cases, careful attention to the diagnosis and treatment of the underlying sleep disorder(s) is of utmost importance. Prescribers should be aware that some patients may have more than one sleep disorder contributing to their excessive sleepiness.

The effectiveness of modafinil in long-term use (greater than 9 weeks in Narcolepsy clinical trials and 12 weeks in OSA and SWD clinical trials) has not been systematically evaluated in placebo-controlled trials. The physician who elects to prescribe modafinil for an extended time in patients with Narcolepsy, OSA, or SWD should periodically reevaluate long-term usefulness for the individual patient.

CONTRAINDICATIONS

Modafinil is contraindicated in patients with known hypersensitivity to modafinil, armodafinil or its inactive ingredients.

WARNINGS

Serious Rash, including Stevens-Johnson Syndrome

Serious rash requiring hospitalization and discontinuation of treatment has been reported in adults and children in association with the use of modafinil.

Modafinil is not approved for use in pediatric patients for any indication.

In clinical trials of modafinil, the incidence of rash resulting in discontinuation was approximately 0.8% (13 per 1,585) in pediatric patients (age <17 years); these rashes included 1 case of possible Stevens-Johnson Syndrome (SJS) and 1 case of apparent multi-organ hypersensitivity reaction. Several of the cases were associated with fever and other abnormalities (e.g., vomiting, leukopenia). The median time to rash that resulted in discontinuation was 13 days. No such cases were observed among 380 pediatric patients who received placebo. No serious skin rashes have been reported in adult clinical trials (0 per 4,264) of modafinil.

Rare cases of serious or life-threatening rash, including SJS, Toxic Epidermal Necrolysis (TEN), and Drug Rash with Eosinophilia and Systemic Symptoms (DRESS) have been reported in adults and children in worldwide post-marketing experience. The reporting rate of TEN and SJS associated with modafinil use, which is generally accepted to be an underestimate due to underreporting, exceeds the background incidence rate. Estimates of the background incidence rate for these serious skin reactions in the general population range between 1 to 2 cases per million-person years.

There are no factors that are known to predict the risk of occurrence or the severity of rash associated with modafinil. Nearly all cases of serious rash associated with modafinil occurred within 1 to 5 weeks after treatment initiation. However, isolated cases have been reported after prolonged treatment (e.g., 3 months). Accordingly, duration of therapy cannot be relied upon as

a means to predict the potential risk heralded by the first appearance of a rash.

Although benign rashes also occur with modafinil, it is not possible to reliably predict which rashes will prove to be serious. Accordingly, modafinil should ordinarily be discontinued at the first sign of rash, unless the rash is clearly not drug-related. Discontinuation of treatment may not prevent a rash from becoming life-threatening or permanently disabling or disfiguring.

Angioedema and Anaphylactoid Reactions

One serious case of angioedema and one case of hypersensitivity (with rash, dysphagia, and bronchospasm), were observed among 1,595 patients treated with armodafinil, the R enantiomer of modafinil (which is the racemic mixture). No such cases were observed in modafinil clinical trials. However, angioedema has been reported in postmarketing experience with modafinil. Patients should be advised to discontinue therapy and immediately report to their physician any signs or symptoms suggesting angioedema or anaphylaxis (e.g., swelling of face, eyes, lips, tongue or larynx; difficulty in swallowing or breathing; hoarseness).

Multi-organ Hypersensitivity Reactions

Multi-organ hypersensitivity reactions, including at least one fatality in postmarketing experience, have occurred in close temporal association (median time to detection 13 days: range 4-33) to the initiation of modafinil.

Although there have been a limited number of reports, multi-organ hypersensitivity reactions may result in hospitalization or be life-threatening. There are no factors that are known to predict the risk of occurrence or the severity of multi-organ hypersensitivity reactions associated with modafinil. Signs and symptoms of this disorder were diverse; however, patients typically, although not exclusively, presented with fever and rash associated with other organ system involvement. Other associated manifestations included myocarditis, hepatitis, liver function test abnormalities, hematological abnormalities (e.g., eosinophilia, leukopenia, thrombocytopenia), pruritus, and asthenia. Because multi-organ hypersensitivity is variable in its expression, other organ system symptoms and signs, not noted here, may occur.

If a multi-organ hypersensitivity reaction is suspected, modafinil should be discontinued. Although there are no case reports to indicate cross-sensitivity with other drugs that produce this syndrome, the experience with drugs associated with multi-organ hypersensitivity would indicate this to be a possibility.

Persistent Sleepiness

Patients with abnormal levels of sleepiness who take modafinil should be advised that their level of wakefulness may not return to normal. Patients with excessive sleepiness, including those taking modafinil, should be frequently reassessed for their degree of sleepiness and, if appropriate, advised to avoid driving or any other potentially dangerous activity. Prescribers should also be aware that patients may not acknowledge sleepiness or drowsiness until directly questioned about drowsiness or sleepiness during specific activities.

Psychiatric Symptoms

Psychiatric adverse experiences have been reported in patients treated with modafinil.

Postmarketing adverse events associated with the use of modafinil have included mania, delusions, hallucinations, suicidal ideation and aggression, some resulting in hospitalization. Many, but not all, patients had a prior psychiatric history. One healthy male volunteer developed ideas of reference, paranoid delusions, and auditory hallucinations in association with multiple daily 600 mg doses of modafinil and sleep deprivation. There was no evidence of psychosis 36 hours after drug discontinuation.

In the adult modafinil controlled trials database, psychiatric symptoms resulting in treatment discontinuation (at a frequency $\geq 0.3\%$) and reported more often in patients treated with modafinil compared to those treated with placebo were anxiety (1%), nervousness (1%), insomnia ($<1\%$), confusion ($<1\%$), agitation ($<1\%$), and depression ($<1\%$). Caution should be exercised when modafinil is given to patients with a history of psychosis, depression, or mania. Consideration should be given to the possible emergence or exacerbation of psychiatric symptoms in patients treated with modafinil. If psychiatric symptoms develop in association with modafinil administration, consider discontinuing modafinil.

PRECAUTIONS

Diagnosis of Sleep Disorders

Modafinil should be used only in patients who have had a complete evaluation of their excessive sleepiness, and in whom a diagnosis of either narcolepsy, OSA, and/or SWD has been made in accordance with ICSD or DSM diagnostic criteria (See **CLINICAL TRIALS**). Such an evaluation usually consists of a complete history and physical examination, and it may be supplemented with testing in a laboratory setting. Some patients may have more than one sleep disorder contributing to their excessive sleepiness (e.g., OSA and SWD coincident in the same patient).

General

Although modafinil has not been shown to produce functional impairment, any drug affecting the CNS may alter judgment, thinking or motor skills. Patients should be cautioned about operating an automobile or other hazardous machinery until they are reasonably certain that modafinil therapy will not adversely affect their ability to engage in such activities.

CPAP Use in Patients with OSA

In OSA, modafinil is indicated as an adjunct to standard treatment(s) for the underlying obstruction. If continuous positive airway pressure (CPAP) is the treatment of choice for a patient, a maximal effort to treat with CPAP for an adequate period of time should be made prior to initiating modafinil. If modafinil is used adjunctively with CPAP, the encouragement of and periodic assessment of CPAP compliance is necessary.

Cardiovascular System

Modafinil has not been evaluated in patients with a recent history of myocardial infarction or unstable angina, and such patients should be treated with caution.

In clinical studies of modafinil, signs and symptoms including chest pain, palpitations, dyspnea and transient ischemic T-wave changes on ECG were observed in three subjects in association with mitral valve prolapse or left ventricular hypertrophy. It is recommended that modafinil tablets not be used in patients with a history of left ventricular hypertrophy or in patients with mitral valve prolapse who have experienced the mitral valve prolapse syndrome when previously receiving CNS stimulants. Such signs may include but are not limited to ischemic ECG changes, chest pain, or arrhythmia. If new onset of any of these symptoms occurs, consider cardiac evaluation.

Blood pressure monitoring in short-term (<3 months) controlled trials showed no clinically significant changes in mean systolic and diastolic blood pressure in patients receiving modafinil as compared to placebo. However, a retrospective analysis of the use of antihypertensive medication in these studies showed that a greater proportion of patients on modafinil required new or increased use of antihypertensive medications (2.4%) compared to patients on placebo (0.7%). The differential use was slightly larger when only studies in OSA were included, with 3.4% of patients on modafinil and 1.1% of patients on placebo requiring such alterations in the use of antihypertensive medication. Increased monitoring of blood pressure may be appropriate in patients on modafinil.

Patients Using Steroidal Contraceptives

The effectiveness of steroidal contraceptives may be reduced when used with modafinil tablets and for one month after discontinuation of therapy (See **PRECAUTIONS, Drug Interactions**). Alternative or concomitant methods of contraception are recommended for patients treated with modafinil tablets, and for one month after discontinuation of modafinil.

Patients Using Cyclosporine

The blood levels of cyclosporine may be reduced when used with modafinil (See **PRECAUTIONS, Drug Interactions**). Monitoring of circulating cyclosporine concentrations and appropriate dosage adjustment for cyclosporine should be considered when these drugs are used concomitantly.

Patients with Severe Hepatic Impairment

In patients with severe hepatic impairment, with or without cirrhosis (See **CLINICAL PHARMACOLOGY**), modafinil should be administered at a reduced dose (See **DOSAGE AND ADMINISTRATION**).

Patients with Severe Renal Impairment

There is inadequate information to determine safety and efficacy of dosing in patients with severe renal impairment. (For pharmacokinetics in renal impairment, see **CLINICAL PHARMACOLOGY**.)

Elderly patients

In elderly patients, elimination of modafinil and its metabolites may be reduced as a consequence of aging. Therefore, consideration should be given to the use of lower doses in this population. (See **CLINICAL PHARMACOLOGY** and **DOSAGE AND ADMINISTRATION**).

Information for Patients

Physicians are advised to discuss the following issues with patients for whom they prescribe modafinil.

Modafinil is indicated for patients who have abnormal levels of sleepiness. Modafinil has been shown to improve, but not eliminate this abnormal tendency to fall asleep. Therefore, patients should not alter their previous behavior with regard to potentially dangerous activities (e.g., driving, operating machinery) or other activities requiring appropriate levels of wakefulness, until and unless treatment with modafinil has been shown to produce levels of wakefulness that permit such activities. Patients should be advised that modafinil is not a replacement for sleep.

Patients should be informed that it may be critical that they continue to take their previously prescribed treatments (e.g., patients with OSA receiving CPAP should continue to do so).

Patients should be informed of the availability of a Medication Guide, and they should be instructed to read it prior to taking modafinil. The complete text of the Medication Guide is provided at the end of this labeling.

Patients should be advised to contact their physician if they experience chest pain, rash, depression, anxiety, or signs of psychosis or mania.

Pregnancy

Patients should be advised to notify their physician if they become pregnant or intend to become pregnant during therapy. Patients should be cautioned regarding the potential increased risk of pregnancy when using steroidal contraceptives (including depot or implantable contraceptives) with modafinil and for one month after discontinuation of therapy (See **Carcinogenesis, Mutagenesis, Impairment of Fertility and Pregnancy**).

Nursing

Patients should be advised to notify their physician if they are breast feeding an infant.

Concomitant Medications

Patients should be advised to inform their physician if they are taking, or plan to take, any prescription or over-the-counter drugs, because of the potential for interactions between modafinil and other drugs.

Alcohol

Patients should be advised that the use of modafinil in combination with alcohol has not been studied. Patients should be advised that it is prudent to avoid alcohol while taking modafinil.

Allergic Reactions

Patients should be advised to stop taking modafinil and to notify their physician if they develop a rash, hives, mouth sores, blisters, peeling skin, trouble swallowing or breathing or a related allergic phenomenon.

Drug Interactions

CNS Active Drugs

Methylphenidate - In a single-dose study in healthy volunteers, simultaneous administration of modafinil (200 mg) with methylphenidate (40 mg) did not cause any significant alterations in the pharmacokinetics of either drug. However, the absorption of modafinil may be delayed by approximately one hour when coadministered with methylphenidate.

In a multiple-dose, steady-state study in healthy volunteers, modafinil was administered once daily at 200 mg/day for 7 days followed by 400 mg/day for 21 days. Administration of methylphenidate (20 mg/day) during days 22-28 of modafinil treatment 8 hours after the daily dose of modafinil did not cause any significant alterations in the pharmacokinetics of modafinil.

Dextroamphetamine - In a single dose study in healthy volunteers, simultaneous administration of modafinil (200 mg) with dextroamphetamine (10 mg) did not cause any significant alterations in the pharmacokinetics of either drug. However, the absorption of modafinil may be delayed by approximately one hour when coadministered with dextroamphetamine.

In a multiple-dose, steady-state study in healthy volunteers, modafinil was administered once daily at 200 mg/day for 7 days followed by 400 mg/day for 21 days. Administration of dextroamphetamine (20 mg/day) during days 22-28 of modafinil treatment 7 hours after the daily dose of modafinil did not cause any significant alterations in the pharmacokinetics of modafinil.

Clomipramine - The coadministration of a single dose of clomipramine (50 mg) on the first of three days of treatment with modafinil (200 mg/day) in healthy volunteers did not show an effect on the pharmacokinetics of either drug. However, one incident of increased levels of clomipramine and its active metabolite desmethylclomipramine has been reported in a patient with narcolepsy during treatment with modafinil.

Triazolam – In the drug interaction study between modafinil and ethinyl estradiol (EE₂), on the same days as those for the plasma sampling for EE₂ pharmacokinetics, a single dose of triazolam (0.125 mg) was also administered. Mean C_{max} and AUC_{0-∞} of triazolam were decreased by 42% and 59%, respectively, and its elimination half-life was decreased by approximately an hour after the modafinil treatment.

Monoamine Oxidase (MAO) Inhibitors - Interaction studies with monoamine oxidase inhibitors have not been performed. Therefore, caution should be used when concomitantly administering MAO inhibitors and modafinil.

Other Drugs

Warfarin - There were no significant changes in the pharmacokinetic profiles of R- and S- warfarin in

healthy subjects given a single dose of racemic warfarin (5 mg) following chronic administration of modafinil (200 mg/day for 7 days followed by 400 mg/day for 27 days) relative to the profiles in subjects given placebo. However, more frequent monitoring of prothrombin times/INR is advisable whenever modafinil is coadministered with warfarin (See **CLINICAL PHARMACOLOGY, Pharmacokinetics**, Drug-Drug Interactions).

Ethinyl Estradiol - Administration of modafinil to female volunteers once daily at 200 mg/day for 7 days followed by 400 mg/day for 21 days resulted in a mean 11% decrease in C_{max} and 18% decrease in AUC_{0-24} of ethinyl estradiol (EE_2 ; 0.035 mg; administered orally with norgestimate). There was no apparent change in the elimination rate of ethinyl estradiol.

Cyclosporine - One case of an interaction between modafinil and cyclosporine, a substrate of CYP3A4, has been reported in a 41 year old woman who had undergone an organ transplant. After one month of administration of 200 mg/day of modafinil, cyclosporine blood levels were decreased by 50%. The interaction was postulated to be due to the increased metabolism of cyclosporine, since no other factor expected to affect the disposition of the drug had changed. Dosage adjustment for cyclosporine may be needed.

Potential Interactions with Drugs That Inhibit, Induce, or are Metabolized by Cytochrome P-450 Isoenzymes and Other Hepatic Enzymes

In *in vitro* studies using primary human hepatocyte cultures, modafinil was shown to slightly induce CYP1A2, CYP2B6 and CYP3A4 in a concentration-dependent manner. Although induction results based on *in vitro* experiments are not necessarily predictive of response *in vivo*, caution needs to be exercised when modafinil is coadministered with drugs that depend on these three enzymes for their clearance. Specifically, lower blood levels of such drugs could result (See *Other Drugs*, Cyclosporine above).

The exposure of human hepatocytes to modafinil *in vitro* produced an apparent concentration-related suppression of expression of CYP2C9 activity suggesting that there is a potential for a metabolic interaction between modafinil and the substrates of this enzyme (e.g., S-warfarin and phenytoin). In a subsequent clinical study in healthy volunteers, chronic modafinil treatment did not show a significant effect on the single-dose pharmacokinetics of warfarin when compared to placebo (See *Other Drugs*, Warfarin above).

In vitro studies using human liver microsomes showed that modafinil reversibly inhibited CYP2C19 at pharmacologically relevant concentrations of modafinil. CYP2C19 is also reversibly inhibited, with similar potency, by a circulating metabolite, modafinil sulfone. Although the maximum plasma concentrations of modafinil sulfone are much lower than those of parent modafinil, the combined effect of both compounds could produce sustained partial inhibition of the enzyme. Drugs that are largely eliminated via CYP2C19 metabolism, such as diazepam, propranolol, phenytoin (also via CYP2C9) or S-mephenytoin may have prolonged elimination upon coadministration with modafinil and may require dosage reduction and monitoring for toxicity.

Tricyclic antidepressants - CYP2C19 also provides an ancillary pathway for the metabolism of certain tricyclic antidepressants (e.g., clomipramine and desipramine) that are primarily metabolized by CYP2D6. In tricyclic-treated patients deficient in CYP2D6 (i.e., those who are poor metabolizers of debrisoquine; 7-10% of the Caucasian population; similar or lower in other populations), the amount of metabolism by CYP2C19 may be substantially increased.

Modafinil may cause elevation of the levels of the tricyclics in this subset of patients. Physicians should be aware that a reduction in the dose of tricyclic agents might be needed in these patients.

In addition, due to the partial involvement of CYP3A4 in the metabolic elimination of modafinil, coadministration of potent inducers of CYP3A4 (e.g., carbamazepine, phenobarbital, rifampin) or inhibitors of CYP3A4 (e.g., ketoconazole, itraconazole) could alter the plasma levels of modafinil.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Carcinogenicity studies were conducted in which modafinil was administered in the diet to mice for 78 weeks and to rats for 104 weeks at doses of 6, 30, and 60 mg/kg/day. The highest dose studied is 1.5 (mouse) or 3 (rat) times greater than the recommended adult human daily dose of modafinil (200 mg) on a mg/m² basis. There was no evidence of tumorigenesis associated with modafinil administration in these studies. However, since the mouse study used an inadequate high dose that was not representative of a maximum tolerated dose, a subsequent carcinogenicity study was conducted in the Tg.AC transgenic mouse. Doses evaluated in the Tg.AC assay were 125, 250, and 500 mg/kg/day, administered dermally. There was no evidence of tumorigenicity associated with modafinil administration; however, this dermal model may not adequately assess the carcinogenic potential of an orally administered drug.

Mutagenesis

Modafinil demonstrated no evidence of mutagenic or clastogenic potential in a series of in vitro (i.e., bacterial reverse mutation assay, mouse lymphoma tk assay, chromosomal aberration assay in human lymphocytes, cell transformation assay in BALB/3T3 mouse embryo cells) assays in the absence or presence of metabolic activation, or in vivo (mouse bone marrow micronucleus) assays. Modafinil was also negative in the unscheduled DNA synthesis assay in rat hepatocytes.

Impairment of Fertility

Oral administration of modafinil (doses of up to 480 mg/kg/day) to male and female rats prior to and throughout mating, and continuing in females through day 7 of gestation produced an increase in the time to mate at the highest dose; no effects were observed on other fertility or reproductive parameters. The no-effect dose of 240 mg/kg/day was associated with a plasma modafinil exposure (AUC) approximately equal to that in humans at the recommended dose of 200 mg.

Pregnancy

Pregnancy Category C: In studies conducted in rats and rabbits, developmental toxicity was observed at clinically relevant exposures.

Modafinil (50, 100, or 200 mg/kg/day) administered orally to pregnant rats throughout the period of organogenesis caused, in the absence of maternal toxicity, an increase in resorptions and an increased incidence of visceral and skeletal variations in the offspring at the highest dose. The higher no-effect dose for rat embryofetal developmental toxicity was associated with a plasma modafinil exposure approximately 0.5 times the AUC in humans at the recommended daily dose (RHD) of 200 mg. However, in a subsequent study of up to 480 mg/kg/day (plasma modafinil exposure approximately 2 times the AUC in humans at the RHD) no adverse effects on embryofetal development were observed.

Modafinil administered orally to pregnant rabbits throughout the period of organogenesis at doses of 45, 90, and 180 mg/kg/day increased the incidences of fetal structural alterations and embryofetal death at the highest dose. The highest no-effect dose for developmental toxicity was associated with a plasma modafinil AUC approximately equal to the AUC in humans at the RHD.

Oral administration of armodafinil (the R-enantiomer of modafinil; 60, 200, or 600 mg/kg/day) to pregnant rats throughout the period of organogenesis resulted in increased incidences of fetal visceral and skeletal variations at the intermediate dose or greater and decreased fetal body weights at the highest dose. The no-effect dose for rat embryofetal developmental toxicity was associated with a plasma armodafinil exposure (AUC) approximately one-tenth times the AUC for armodafinil in humans treated with modafinil at the RHD.

Modafinil administration to rats throughout gestation and lactation at oral doses of up to 200 mg/kg/day resulted in decreased viability in the offspring at doses greater than 20 mg/kg/day (plasma modafinil AUC approximately 0.1 times the AUC in humans at the RHD). No effects on postnatal developmental and neurobehavioral parameters were observed in surviving offspring.

There are no adequate and well-controlled studies in pregnant women. Two cases of intrauterine

growth retardation and one case of spontaneous abortion have been reported in association with armodafinil and modafinil. Although the pharmacology of modafinil and armodafinil is not identical to that of the sympathomimetic amines, they do share some pharmacologic properties with this class. Certain of these drugs have been associated with intrauterine growth retardation and spontaneous abortions. Whether the cases reported are drug-related is unknown.

Modafinil should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Pregnancy Registry: A pregnancy registry has been established to collect information on the pregnancy outcomes of women exposed to modafinil. Healthcare providers are encouraged to register pregnant patients, or pregnant women may enroll themselves in the registry by calling 1-866-404-4106 (toll free).

Labor and Delivery

The effect of modafinil on labor and delivery in humans has not been systematically investigated.

Nursing Mothers

It is not known whether modafinil or its metabolites are excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when modafinil tablets are administered to a nursing woman.

Pediatric Use

Safety and effectiveness in pediatric patients, below age 17, have not been established. Serious skin rashes, including erythema multiforme major (EMM) and Stevens-Johnson Syndrome (SJS) have been associated with modafinil use in pediatric patients (see **WARNINGS, Serious Rash, including Stevens-Johnson Syndrome**).

In a controlled 6-week study, 165 pediatric patients (aged 5-17 years) with narcolepsy were treated with modafinil (n=123), or placebo (n=42). There were no statistically significant differences favoring modafinil over placebo in prolonging sleep latency as measured by MSLT, or in perceptions of sleepiness as determined by the clinical global impression-clinician scale (CGI-C).

In the controlled and open-label clinical studies, treatment emergent adverse events of the psychiatric and nervous system included Tourette's syndrome, insomnia, hostility, increased cataplexy, increased hypnagogic hallucinations and suicidal ideation. Transient leukopenia, which resolved without medical intervention, was also observed. In the controlled clinical study, 3 of 38 girls, ages 12 or older, treated with modafinil experienced dysmenorrhea compared to 0 of 10 girls who received placebo.

There were three 7 to 9 week, double-blind, placebo-controlled, parallel group studies in children and adolescents (aged 6-17 years) with Attention-Deficit Hyperactivity Disorder (ADHD, DSM-IV). Two of the studies were flexible-dose studies (up to 425 mg/day), and the third was a fixed-dose study (340 mg/day for patients <30 kg and 425 mg/day for patients ≥30 kg). Although these studies showed statistically significant differences favoring modafinil over placebo in reducing ADHD symptoms as measured by the ADHD-RS (school version), there were 3 cases of serious rash including one case of possible SJS among 933 patients exposed to modafinil in this program.

Modafinil is not approved for use in pediatric patients for any indication, including ADHD (see **WARNINGS, Serious Rash, including Stevens-Johnson Syndrome**).

Geriatric Use

Experience in a limited number of patients who were greater than 65 years of age in clinical trials showed an incidence of adverse experiences similar to other age groups. In elderly patients, elimination of modafinil and its metabolites may be reduced as a consequence of aging. Therefore, consideration should be given to the use of lower doses in this population (See **CLINICAL PHARMACOLOGY**

and **PRECAUTIONS**).

ADVERSE REACTIONS

Modafinil has been evaluated for safety in over 3500 patients, of whom more than 2000 patients with excessive sleepiness associated with primary disorders of sleep and wakefulness were given at least one dose of modafinil. In clinical trials, modafinil has been found to be generally well tolerated and most adverse experiences were mild to moderate.

The most commonly observed adverse events ($\geq 5\%$) associated with the use of modafinil more frequently than placebo-treated patients in the placebo-controlled clinical studies in primary disorders of sleep and wakefulness were headache, nausea, nervousness, rhinitis, diarrhea, back pain, anxiety, insomnia, dizziness, and dyspepsia. The adverse event profile was similar across these studies.

In the placebo-controlled clinical trials, 74 of the 934 patients (8%) who received modafinil discontinued due to an adverse experience compared to 3% of patients that received placebo. The most frequent reasons for discontinuation that occurred at a higher rate for modafinil than placebo patients were headache (2%), nausea, anxiety, dizziness, insomnia, chest pain and nervousness (each $<1\%$). In a Canadian clinical trial, a 35 year old obese narcoleptic male with a prior history of syncopal episodes experienced a 9-second episode of asystole after 27 days of modafinil treatment (300 mg/day in divided doses).

Incidence in Controlled Trials

The following table (Table 3) presents the adverse experiences that occurred at a rate of 1% or more and were more frequent in adult patients treated with modafinil than in placebo-treated patients in the principal, placebo-controlled clinical trials.

The prescriber should be aware that the figures provided below cannot be used to predict the frequency of adverse experiences in the course of usual medical practice, where patient characteristics and other factors may differ from those occurring during clinical studies. Similarly, the cited frequencies cannot be directly compared with figures obtained from other clinical investigations involving different treatments, uses, or investigators. Review of these frequencies, however, provides prescribers with a basis to estimate the relative contribution of drug and non-drug factors to the incidence of adverse events in the population studied.

Table 3. Incidence Of Treatment-Emergent Adverse Experiences In Parallel-Group, Placebo-Controlled Clinical Trials¹ With Modafinil In Adults With Narcolepsy, OSA, and SWD (200mg, 300mg and 400mg)*

Body System	Preferred Term	Modafinil (n = 934)	Placebo (n = 567)
Body as a Whole	Headache	34%	23%
	Back Pain	6%	5%
	Flu Syndrome	4%	3%
	Chest Pain	3%	1%
	Chills	1%	0%
	Neck Rigidity	1%	0%
Cardiovascular	Hypertension	3%	1%
	Tachycardia	2%	1%
	Palpitation	2%	1%
	Vasodilatation	2%	0%
Digestive	Nausea	11%	3%

	Diarrhea	6%	5%
	Dyspepsia	5%	4%
	Dry Mouth	4%	2%
	Anorexia	4%	1%
	Constipation	2%	1%
	Abnormal Liver Function ²	2%	1%
	Flatulence	1%	0%
	Mouth Ulceration	1%	0%
	Thirst	1%	0%
Hemic/Lymphatic	Eosinophilia	1%	0%
Metabolic/Nutritional	Edema	1%	0%
Nervous	Nervousness	7%	3%
	Insomnia	5%	1%
	Anxiety	5%	1%
	Dizziness	5%	4%
	Depression	2%	1%
	Paresthesia	2%	0%
	Somnolence	2%	1%
	Hypertonia	1%	0%
	Dyskinesia ³	1%	0%
	Hyperkinesia	1%	0%
	Agitation	1%	0%
	Confusion	1%	0%
	Tremor	1%	0%
	Emotional Lability	1%	0%
	Vertigo	1%	0%
Respiratory	Rhinitis	7%	6%
	Pharyngitis	4%	2%
	Lung Disorder	2%	1%
	Epistaxis	1%	0%
	Asthma	1%	0%
Skin/Appendages	Sweating	1%	0%
	Herpes Simplex	1%	0%
Special Senses	Amblyopia	1%	0%
	Abnormal Vision	1%	0%
	Taste Perversion	1%	0%
	Eye Pain	1%	0%
Urogenital	Urine Abnormality	1%	0%
	Hematuria	1%	0%

	Pyuria	1%	0%
--	--------	----	----

* Six double-blind, placebo-controlled clinical studies in narcolepsy, OSA, and SWD.

¹ Events reported by at least 1% of patients treated with modafinil that were more frequent than in the placebo group are included;

incidence is rounded to the nearest 1%. The adverse experience terminology is coded using a standard modified COSTART

Dictionary.

Events for which the modafinil incidence was at least 1%, but equal to or less than placebo are not listed in the table. These

events included the following: infection, pain, accidental injury, abdominal pain, hypothermia, allergic reaction, asthenia, fever,

viral infection, neck pain, migraine, abnormal electrocardiogram, hypotension, tooth disorder, vomiting, periodontal abscess,

increased appetite, ecchymosis, hyperglycemia, peripheral edema, weight loss, weight gain, myalgia, leg cramps, arthritis,

cataplexy, thinking abnormality, sleep disorder, increased cough, sinusitis, dyspnea, bronchitis, rash, conjunctivitis, ear pain,

dysmenorrhea⁴, urinary tract infection.

² Elevated liver enzymes.

³ Oro-facial dyskinesias.

⁴ Incidence adjusted for gender.

Dose Dependency of Adverse Events

In the adult placebo-controlled clinical trials which compared doses of 200, 300, and 400 mg/day of modafinil and placebo, the only adverse events that were clearly dose related were headache and anxiety.

Vital Sign Changes

While there was no consistent change in mean values of heart rate or systolic and diastolic blood pressure, the requirement for antihypertensive medication was slightly greater in patients on modafinil compared to placebo (See **PRECAUTIONS**).

Weight Changes

There were no clinically significant differences in body weight change in patients treated with modafinil compared to placebo-treated patients in the placebo-controlled clinical trials.

Laboratory Changes

Clinical chemistry, hematology, and urinalysis parameters were monitored in Phase 1, 2, and 3 studies. In these studies, mean plasma levels of gamma glutamyltransferase (GGT) and alkaline phosphatase (AP) were found to be higher following administration of modafinil, but not placebo. Few subjects, however, had GGT or AP elevations outside of the normal range. Shifts to higher, but not clinically significantly abnormal, GGT and AP values appeared to increase with time in the population treated with modafinil in the Phase 3 clinical trials. No differences were apparent in alanine aminotransferase, aspartate aminotransferase, total protein, albumin, or total bilirubin.

ECG Changes

No treatment-emergent pattern of ECG abnormalities was found in placebo-controlled clinical trials following administration of modafinil.

Postmarketing Reports

The following adverse reactions have been identified during post-approval use of modafinil. Because these reactions are reported voluntarily from a population of uncertain size, it is not possible to reliably estimate their frequency or establish a causal relationship to drug exposure. Decisions to include these reactions in labeling are typically based on one or more of the following factors: (1) seriousness of the reaction, (2) frequency of the reporting, or (3) strength of causal connection to modafinil.

Hematologic: agranulocytosis

DRUG ABUSE AND DEPENDENCE

Controlled Substance Class

Modafinil is listed in Schedule IV of the Controlled Substances Act.

Abuse Potential and Dependence

In addition to its wakefulness-promoting effect and increased locomotor activity in animals, in humans, modafinil produces psychoactive and euphoric effects, alterations in mood, perception, thinking and feelings typical of other CNS stimulants. In *in vitro* binding studies, modafinil binds to the dopamine reuptake site and causes an increase in extracellular dopamine, but no increase in dopamine release. Modafinil is reinforcing, as evidenced by its self-administration in monkeys previously trained to self-administer cocaine. In some studies, modafinil was also partially discriminated as stimulant-like. Physicians should follow patients closely, especially those with a history of drug and/or stimulant (e.g., methylphenidate, amphetamine, or cocaine) abuse. Patients should be observed for signs of misuse or abuse (e.g., incrementation of doses or drug-seeking behavior).

The abuse potential of modafinil (200, 400, and 800 mg) was assessed relative to methylphenidate (45 and 90 mg) in an inpatient study in individuals experienced with drugs of abuse. Results from this clinical study demonstrated that modafinil produced psychoactive and euphoric effects and feelings consistent with other scheduled CNS stimulants (methylphenidate).

Withdrawal

The effects of modafinil withdrawal were monitored following 9 weeks of modafinil use in one US Phase 3 controlled clinical trial. No specific symptoms of withdrawal were observed during 14 days of observation, although sleepiness returned in narcoleptic patients.

OVERDOSAGE

Human Experience

In clinical trials, a total of 151 protocol-specified doses ranging from 1000 to 1600 mg/day (5 to 8 times the recommended daily dose of 200 mg) have been administered to 32 subjects, including 13 subjects who received doses of 1000 or 1200 mg/day for 7 to 21 consecutive days. In addition, several intentional acute overdoses occurred; the two largest being 4500 mg and 4000 mg taken by two subjects participating in foreign depression studies. None of these study subjects experienced any unexpected or life-threatening effects. Adverse experiences that were reported at these doses included excitation or agitation, insomnia, and slight or moderate elevations in hemodynamic parameters. Other observed high-dose effects in clinical studies have included anxiety, irritability, aggressiveness, confusion, nervousness, tremor, palpitations, sleep disturbances, nausea, diarrhea and decreased prothrombin time.

From post-marketing experience, there have been no reports of fatal overdoses involving modafinil

alone (doses up to 12 grams). Overdoses involving multiple drugs, including modafinil, have resulted in fatal outcomes. Symptoms most often accompanying modafinil overdose, alone or in combination with other drugs have included: insomnia; central nervous system symptoms such as restlessness, disorientation, confusion, excitation and hallucination; digestive changes such as nausea and diarrhea; and cardiovascular changes such as tachycardia, bradycardia, hypertension and chest pain.

Cases of accidental ingestion/overdose have been reported in children as young as 11 months of age. The highest reported accidental ingestion on a mg/kg basis occurred in a three-year-old boy who ingested 800-1000 mg (50-63 mg/kg) of modafinil. The child remained stable. The symptoms associated with overdose in children were similar to those observed in adults.

Overdose Management

No specific antidote to the toxic effects of modafinil overdose has been identified to date. Such overdoses should be managed with primarily supportive care, including cardiovascular monitoring. If there are no contraindications, induced emesis or gastric lavage should be considered. There are no data to suggest the utility of dialysis or urinary acidification or alkalization in enhancing drug elimination. The physician should consider contacting a poison-control center on the treatment of any overdose.

DOSAGE AND ADMINISTRATION

The recommended dose of modafinil is 200 mg given once a day.

For patients with narcolepsy and OSA, modafinil should be taken as a single dose in the morning.

For patients with SWD, modafinil should be taken approximately 1 hour prior to the start of their work shift.

Doses up to 400 mg/day, given as a single dose, have been well tolerated, but there is no consistent evidence that this dose confers additional benefit beyond that of the 200 mg dose (See **CLINICAL PHARMACOLOGY** and **CLINICAL TRIALS**).

General Considerations

Dosage adjustment should be considered for concomitant medications that are substrates for CYP3A4, such as triazolam and cyclosporine (See **PRECAUTIONS, Drug Interactions**).

Drugs that are largely eliminated via CYP2C19 metabolism, such as diazepam, propranolol, phenytoin (also via CYP2C9) or S-mephenytoin may have prolonged elimination upon coadministration with modafinil and may require dosage reduction and monitoring for toxicity.

In patients with severe hepatic impairment, the dose of modafinil should be reduced to one-half of that recommended for patients with normal hepatic function (See **CLINICAL PHARMACOLOGY** and **PRECAUTIONS**).

There is inadequate information to determine safety and efficacy of dosing in patients with severe renal impairment (See **CLINICAL PHARMACOLOGY** and **PRECAUTIONS**).

In elderly patients, elimination of modafinil and its metabolites may be reduced as a consequence of aging. Therefore, consideration should be given to the use of lower doses in this population (See **CLINICAL PHARMACOLOGY** and **PRECAUTIONS**).

HOW SUPPLIED

Repackaged by Aphena Pharma Solutions - TN.

See **Repackaging Information** for available configurations.



100 mg Each capsule-shaped, white, uncoated tablet is debossed with "PROVIGIL" on one side and "100 MG" on the other.

NDC 55253-801-30 - Bottles of 30

NDC 55253-801-90 - Bottles of 90

200 mg Each capsule-shaped, white, scored, uncoated tablet is debossed with "PROVIGIL" on one side and "200 MG" on the other.

NDC 55253-802-30 - Bottles of 30

NDC 55253-802-90 - Bottles of 90

Store at 20° to 25° C (68° to 77° F).

Manufactured In Canada By:

Patheon Inc.

Mississauga, Ontario, Canada L5N 7K9

Manufactured For:

Cephalon, Inc.

Frazer, PA 19355

February 2012

MODT-001

PROVIGIL® is a trademark of Cephalon, Inc. or its affiliates.

Iss. 2/2012

Medication Guide

Modafinil Tablets C-IV

Read the Medication Guide that comes with modafinil tablets before you start taking it and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking with your doctor about your condition or treatment.

What is the most important information I should know about modafinil tablets?

Modafinil tablets may cause serious side effects including a serious rash or a serious allergic reaction that may affect parts of your body such as your liver or blood cells. Any of these may need to be treated in a hospital and may be life-threatening.

Stop taking modafinil tablets and call your doctor right away or get emergency help if you have any of these symptoms:

- skin rash, hives, sores in your mouth, or your skin blisters and peels
- swelling of your face, eyes, lips, tongue, or throat
- trouble swallowing or breathing
- fever, shortness of breath, swelling of the legs, yellowing of the skin or whites of the eyes, or dark urine.

If you have a severe rash with modafinil tablets, stopping the medicine may not keep the rash from becoming life-threatening or causing you to be permanently disabled or disfigured.

Modafinil tablets are not approved for use in children for any medical condition.

It is not known if modafinil tablets are safe or if they work in children under the age of 17.

What are modafinil tablets?

Modafinil tablets are a prescription medicine used to improve wakefulness in adults who are very sleepy due to one of the following diagnosed sleep disorders:

- narcolepsy
- obstructive sleep apnea (OSA). Modafinil tablets are used with other medical treatments for this sleep disorder. Modafinil tablets do not take the place of using your CPAP machine or other treatments that your doctor has prescribed for this condition. It is important that you continue to use these treatments as prescribed by your doctor.
- shift work disorder (SWD)

Modafinil tablets will not cure these sleep disorders. Modafinil tablets may help the sleepiness caused by these conditions, but it may not stop all your sleepiness. Modafinil tablets do not take the place of getting enough sleep. Follow your doctor's advice about good sleep habits and using other treatments.

Modafinil tablets are a federally controlled substance (C-IV) because they can be abused or lead to dependence. Keep modafinil tablets in a safe place to prevent misuse and abuse. Selling or giving away modafinil tablets may harm others, and is against the law. Tell your doctor if you have ever abused or been dependent on alcohol, prescription medicines or street drugs.

Who should not take modafinil tablets?

Do not take modafinil tablets if you:

- are allergic to any of its ingredients. See the end of this Medication Guide for a complete list of ingredients in modafinil tablets.
- have had a rash or allergic reaction to either modafinil (PROVIGIL[®]) or armodafinil (NUVIGIL[®]). These medicines are very similar.

What should I tell my doctor before taking modafinil tablets?

Tell your doctor about all of your medical conditions including, if you:

- have a history of mental health problems, including psychosis
- have heart problems or had a heart attack
- have high blood pressure. Your blood pressure may need to be checked more often while taking modafinil tablets.
- have liver or kidney problems
- have a history of drug or alcohol abuse or addiction
- are pregnant or planning to become pregnant. It is not known if modafinil tablets will harm your unborn baby.

Pregnancy Registry: There is a registry for women who become pregnant during treatment with modafinil tablets. The purpose of this registry is to collect information about the safety of modafinil tablets during pregnancy. Contact the registry as soon as you learn that you are pregnant, or ask your doctor to contact the registry for you. You or your doctor can get information and enroll you in the registry by calling 1-866-404-4106.

- are breastfeeding. It is not known if modafinil tablets pass into your milk. Talk to your doctor about

the best way to feed your baby if you take modafinil tablets.

Tell your doctor about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements. Modafinil tablets and many other medicines can interact with each other, sometimes causing side effects. Modafinil tablets may affect the way other medicines work, and other medicines may affect how modafinil tablets work. Your dose of modafinil tablets or certain other medicines may need to be changed.

Especially, tell your doctor if you use or take:

- a hormonal birth control method, such as birth control pills, shots, implants, patches, vaginal rings, and intrauterine devices (IUDs). Hormonal birth control methods may not work while you take modafinil tablets. Women who use one of these methods of birth control may have a higher chance for getting pregnant while taking modafinil tablets, and for one month after stopping modafinil tablets. Talk to your doctor about birth control choices that are right for you while taking modafinil tablets.

Know the medicines you take. Keep a list of them and show it to your doctor and pharmacist when you get a new medicine. Your doctor or pharmacist will tell you if it is safe to take modafinil tablets and other medicines together. Do not start any new medicines with modafinil tablets unless your doctor has told you it is okay.

How should I take modafinil tablets?

- Take modafinil tablets exactly as prescribed by your doctor. Your doctor will prescribe the dose of modafinil tablets that is right for you. Do not change your dose of modafinil tablets without talking to your doctor.
- Your doctor will tell you the right time of day to take modafinil tablets.
 - People with narcolepsy or OSA usually take modafinil tablets one time each day in the morning.
 - People with SWD usually take modafinil tablets about 1 hour before their work shift.
- Do not change the time of day you take modafinil tablets unless you have talked to your doctor. If you take modafinil tablets too close to your bedtime, you may find it harder to go to sleep.
- You can take modafinil tablets with or without food.
- If you take more than your prescribed dose or if you take an overdose of modafinil tablets, call your doctor or poison control center right away.

Symptoms of an overdose of modafinil tablets may include:

- Trouble sleeping
- Restlessness
- Confusion
- Feeling disoriented
- Feeling excited
- Hearing, seeing, feeling, or sensing things that are not really there (hallucinations)
- Nausea and diarrhea
- A fast or slow heartbeat
- Chest pain
- Increased blood pressure

What should I avoid while taking modafinil tablets?

- Do not drive a car or do other dangerous activities until you know how modafinil tablets affect you. People with sleep disorders should always be careful about doing things that could be dangerous. Do not change your daily habits until your doctor tells you it is okay.
- You should avoid drinking alcohol. It is not known how drinking alcohol will affect you when taking modafinil tablets.

What are possible side effects of modafinil tablets?

Modafinil tablets may cause serious side effects. Stop taking modafinil tablets and call your doctor right away or get emergency help if you get any of the following:

- **a serious rash or serious allergic reaction.** (See “**What is the most important information I should know about modafinil tablets?**”)
- **mental (psychiatric) symptoms, including:**
 - depression
 - feeling anxious
 - hearing, seeing, feeling, or sensing things that are not really there (hallucinations)
 - an extreme increase in activity and talking (mania)
 - thoughts of suicide
 - aggressive behavior
 - other mental problems
- **symptoms of a heart problem,** including chest pain, abnormal heart beats, and trouble breathing.

Common side effects that can happen in anyone who takes modafinil tablets include:

-
- | | |
|-------------------|--------------------|
| • back pain | • diarrhea |
| • headache | • feeling anxious |
| • nausea | • dizziness |
| • feeling nervous | • upset stomach |
| • stuffy nose | • trouble sleeping |
-

Modafinil tablets is not approved for use in children for any medical condition. In studies of modafinil tablets in children with narcolepsy, side effects included:

- Tourette’s syndrome
- hostile behavior
- increase in sudden loss of muscle tone and severe muscle weakness
- increase in seeing and hearing things when falling asleep
- increase in suicidal thoughts
- low white blood count
- painful menstrual periods

Tell your doctor if you get any side effect that bothers you or that does not go away while taking modafinil tablets.

These are not all the side effects of modafinil tablets. For more information, ask your doctor or pharmacist.

Some effects of modafinil tablets on the brain are the same as other medicines called “stimulants”. These effects may lead to abuse or dependence on modafinil tablets.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store modafinil tablets?

- Store modafinil tablets at room temperature between 68° and 77° F (20° and 25° C).
- **Keep modafinil tablets and all medicines out of the reach of children.**

General information about modafinil tablets

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use modafinil tablets for a condition for which it was not prescribed. Do not give modafinil tablets to

other people, even if they have the same symptoms you have. It may harm them and it is against the law.

This Medication Guide summarizes the most important information about modafinil tablets. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about modafinil tablets that is written for health professionals. For more information, call 1-800-896-5855.

What are the ingredients in modafinil tablets?

Active Ingredient: modafinil

Inactive Ingredients: lactose monohydrate, microcrystalline cellulose, pregelatinized starch, croscarmellose sodium, povidone, and magnesium stearate.

Manufactured In Canada By:

Patheon Inc.

Mississauga, Ontario, Canada L5N 7K9

Manufactured For:

Cephalon, Inc.

Frazer, PA 19355

February 2012

MODMGT-001

This Medication Guide has been approved by the U.S. Food and Drug Administration.

PROVIGIL[®] and NUVIGIL[®] are trademarks of Cephalon, Inc. or its affiliates.

Iss. 2/2012

Repackaging Information

Please reference the **How Supplied** section listed above for a description of individual tablets or capsules. This drug product has been received by Aphena Pharma - TN in a manufacturer or distributor packaged configuration and repackaged in full compliance with all applicable cGMP regulations. The package configurations available from Aphena are listed below:

Count	200mg
30	43353-925-30
60	43353-925-53

Store between 20°-25°C (68°-77°F). See USP Controlled Room Temperature. Dispense in a tight light-resistant container as defined by USP. Keep this and all drugs out of the reach of children.

Repackaged by:



Cookeville, TN 38506

20121219JH

PRINCIPAL DISPLAY PANEL - 200mg

NDC 43353-925 - **Modafinil** 200mg - Rx Only



MODAFINIL

modafinil tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:43353-925(NDC:55253-802)
Route of Administration	ORAL	DEA Schedule	CIV

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
modafinil (UNII: R3UK8 X3U3D) (modafinil - UNII:R3UK8 X3U3D)	modafinil	200 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
lactose (UNII: J2B2A4N98G)	
magnesium stearate (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
povidone (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	white	Score	2 pieces
Shape	OVAL	Size	16mm
Flavor		Imprint Code	PROVIGIL;200;MG

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:43353-925-30	30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/17/2014	
2	NDC:43353-925-53	60 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/17/2014	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA authorized generic	NDA020717	03/29/2012	

Labeler - Aphenia Pharma Solutions - Tennessee, LLC (128385585)**Establishment**

Name	Address	ID/FEI	Business Operations
Aphenia Pharma Solutions - Tennessee, LLC		128385585	Repack(43353-925)

Revised: 12/2017

Aphenia Pharma Solutions - Tennessee, LLC