INFLUENZINUM- influenza a virus a/victoria/2570/2019 ivr-215 (h1n1) hemagglutinin antigen (formaldehyde inactivated), influenza a virus a/tasmania/503/2020 ivr-221 (h3n2) hemagglutinin antigen (formaldehyde inactivated), influenza b virus b/washington/02/2019 hemagglutinin antigen (formaldehyde inactivated), influenza b virus b/phuket/3073/2013 hemagglutinin antigen (formaldehyde inactivated) pellet Boiron

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

INFLUENZINUM 15C

INFLUENZINUM 15C HPUS**

(** contains 0.443 mg of the active ingredient per pellet)

INFLUENZA A VIRUS A/VICTORIA/2570/2019 IVR-215 (H1N1) HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED

INFLUENZA A VIRUS A/TASMANIA/503/2020 IVR-221 (H3N2) HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED

INFLUENZA B VIRUS B/WASHINGTON/02/2019 HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED)

INFLUENZA B VIRUS B/PHUKET/3073/2013 HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED)

After effects of flu or flu-like symptoms*

Stop use and ask a doctor if symptoms persist for more than 3 days or worsen.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children.

Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.

Do not use if pellet dispenser seal is broken.

Contains approx 80 pellets.


How to dispense pellets? Turn tube upside down. Twist until 5 pellets are dispensed into cap. Carefully remove the cap and use it to pour pellets under the tongue.

*CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE NOT ACCEPTED MEDICAL EVIDENCE. NOT FDA EVALUATED.

*C,K,CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.

lactose, sucrose





Drug Facts

Active ingredient^{**}: See product name on front panel (contains 0.443 mg of the active ingredient per pellet).

Uses: See symptoms on front panel.

Warnings: Stop use and ask a doctor if symptoms persist for more than 3 days or worsen. If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children.

Directions: Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.

Other information: Do not use if pellet dispenser seal is broken.



INFLUENZINUM

influenza a virus a/victoria/2570/2019 ivr-215 (h1n1) hemagglutinin antigen (formaldehyde inactivated), influenza a virus a/tasmania/503/2020 ivr-221 (h3n2) hemagglutinin antigen (formaldehyde inactivated), influenza b virus b/washington/02/2019 hemagglutinin antigen (formaldehyde inactivated), influenza b virus b/phuket/3073/2013 hemagglutinin antigen (formaldehyde inactivated) pellet

Product Information					
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0220-2722		
Route of Administration	ORAL				

	Ingred	lient Name		Basis of Stre	nath	Strength
INFLUENZA A VIRUS A/VICTORIA/2570/2019 IVR-215 (H1N1) INFLUE					9 IVR-215 NIN	15 [hp_C] in 15 [hp_C
INFLUENZA B VIRUS ANTIGEN (FORMALD (INFLUENZA B VIRUS B (FORMALDEHYDE INAC	INFLUENZA B VIRUS B/WASHINGTON/02/2019 HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED)		15 [hp_C] in 15 [hp_			
INFLUENZA A VIRUS HEMAGGLUTININ AN CYN792B8TR) (INFLUE HEMAGGLUTININ ANTI UNII:CYN792B8TR)	INFLUENZA A VIRUS A/TASMANIA/503/2020 IVR-221 (H3N2) HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED)		15 [hp_C] in 15 [hp_			
INFLUENZA B VIRUS B/PHUKET/3073/2013 HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) (UNII: 9HB0XUS9TM) (INFLUENZA B VIRUS B/PHUKET/3073/2013 HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) - UNII:9HB0XUS9TM)			: 9HB0XUS9TM) GLUTININ ANTIGEN	INFLUENZA B VIRUS B/PHUKET/3073/2013 HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED)		15 [hp_C] in 15 [hp_
Inactive Ingred	lients					
•						rength
						rengtn
	1H8M554)					
SUCROSE (UNII: C15: LACTOSE, UNSPECI		I (UNII: J2B2A4N98				
LACTOSE, UNSPECI	FIED FORM					
LACTOSE, UNSPECI Product Charac	FIED FORM	5	G)			
LACTOSE, UNSPECI Product Charac Color	FIED FORM	white	G) Score			
LACTOSE, UNSPECI Product Charac Color Shape	FIED FORM	5	G) Score Size		4m	ım
LACTOSE, UNSPECI Product Charac Color Shape Flavor	FIED FORM	white	G) Score	2	4m	nm
LACTOSE, UNSPECI Product Charac Color Shape	FIED FORM	white	G) Score Size	5	4m	nm
LACTOSE, UNSPECI Product Charac Color Shape Flavor	FIED FORM	white	G) Score Size	5	4m	nm
LACTOSE, UNSPECI Product Charac Color Shape Flavor Contains	FIED FORM	white	G) Score Size Imprint Code	Marketing Start Date	Marke	nm eting End Date
LACTOSE, UNSPECI Product Charac Color Shape Flavor Contains Packaging # Item Code 1 NDC:0220-2722- 1	FIED FORM	white ROUND	G) Score Size Imprint Code	Marketing Start	Marke	ting End
LACTOSE, UNSPECI Product Charac Color Shape Flavor Contains Packaging # Item Code 1 NDC:0220-2722- 1	FIED FORM Cteristics Pa 5 [hp_C] in :	white ROUND	G) Score Size Imprint Code	Marketing Start Date	Marke	ting End
LACTOSE, UNSPECI Product Charac Color Shape Flavor Contains Packaging # Item Code 1 NDC:0220-2722- 1	FIED FORM	white ROUND ackage Descr 1 TUBE; Type 0: N	G) Score Size Imprint Code	Marketing Start Date	Marke	ting End
LACTOSE, UNSPECI	FIED FORM	white ROUND ackage Descr 1 TUBE; Type 0: N tion	G) Score Size Imprint Code Internation Ot a Combination Ot Monograph	Marketing Start Date	Marke	ting End

Labeler - Boiron (282560473)

Registrant - Boiron, Inc. (014892269)

Establishment						
Name	Address	ID/FEI	Business Operations			
Boiron		282560473	manufacture(0220-2722)			

Revised: 8/2021

Boiron