

**ROBITUSSIN CF- dextromethorphan hydrobromide, guaifenesin,
pseudoephedrine hydrochloride liquid**
GlaxoSmithKline Consumer Healthcare Holdings (US) LLC

Robitussin® CF

Active ingredients (in each 5 mL tsp)	Purpose
Dextromethorphan HBr, USP 10 mg	Cough suppressant
Guaifenesin, USP 100 mg	Expectorant
Pseudoephedrine HCl, USP 30 mg	Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily relieves these symptoms occurring with a cold:
 - nasal congestion
 - cough due to minor throat and bronchial irritation

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

When using this product do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Directions

- do not take more than 4 doses in any 24-hour period

age	dose
adults and children 12 years and over	2 teaspoonfuls every 4 hours
children 6 years to under 12 years	1 teaspoonful every 4 hours
under 6 years	Do not use

Inactive ingredients

citric acid, FD&C red no. 40, flavors, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sorbitol, water.

Keep out of reach of children. In case of overdose, get medical help right away.

Other information

- store at 20-25°C (68-77°F)
- dosage cup provided

MADE IN U.S.A.

PRINCIPAL DISPLAY PANEL - 118 mL Bottle Carton

Wyeth®

Robitussin®

CF

NASAL DECONGESTANT
COUGH SUPPRESSANT
EXPECTORANT

- **CLEARs**
NASAL STUFFINESS
- **CONTROLS COUGHS**
- **LOOSENS and RELIEVES**
CHEST CONGESTION

COUGH FORMULA

For Children and Adults

4 FL OZ (118 mL)

4 FL OZ (118 mL)

Robitussin[®]

CF

NASAL DECONGESTANT
COUGH SUPPRESSANT
EXPECTORANT

Wyeth[®]

Robitussin[®]

روبیتوسین[®] سی إف

CF

NASAL DECONGESTANT
COUGH SUPPRESSANT
EXPECTORANT

• **CLEARs**

NASAL STUFFINESS

• **CONTROLS COUGHS**

• **LOOSENS and RELIEVES
CHEST CONGESTION**

COUGH FORMULA

For Children and Adults

4 FL OZ (118 mL)

٤ أونصة سائلة (١١٨ مللتر)



LOT:
EXP:
MFD:

NEW FORMULA

Robitussin[®]

CF

NASAL DECONGESTANT
COUGH SUPPRESSANT
EXPECTORANT

CHOOSE the correct Robitussin Formula for your Symptoms.

Chest Congestion with Dry Cough?

Robitussin

- Loosens Chest Congestion to Help Coughs Clear Chest.

Frequent Coughs + Chest Congestion?

Robitussin - DM

- Controls and Suppresses Coughing.
- Loosens and Relieves Chest Congestion.

Nasal Stuffiness + Chest Congestion with Dry Cough?

Robitussin - PE

- Clears Nasal Passages to Restore Freer Breathing Through the Nose.
- Loosens and Relieves Chest Congestion.

Nasal Stuffiness + Frequent Coughs + Chest Congestion?

Robitussin - CF

- Clears Nasal Passages to Restore Freer Breathing Through the Nose.
- Controls and Suppresses Coughing.
- Relieves Chest Congestion.
- Sugar-Free.

MADE IN U.S.A.

PACKAGED WITH TAMPER-EVIDENT BOTTLE CAP

NEW FORMULA

Robitussin®

CF

NASAL DECONGESTANT
COUGH SUPPRESSANT
EXPECTORANT

Active ingredients (in each 5 mL tsp) **Purpose**
Dextromethorphan HBr, USP 10 mg Cough suppressant
Guaifenesin, USP 100 mg Expectorant
Pseudoephedrine HCl, USP 30 mg Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily relieves these symptoms occurring with a cold:
 - nasal congestion
 - cough due to minor throat and bronchial irritation

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease • high blood pressure • thyroid disease • diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

When using this product do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Directions

- do not take more than 4 doses in any 24-hour period

age	dose
adults and children 12 years and over	2 teaspoonfuls every 4 hours
children 6 years to under 12 years	1 teaspoonful every 4 hours
under 6 years	Do not use

Inactive ingredients citric acid, FD&C red no. 40, flavors, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sorbitol, water.

NEW FORMULA

Robitussin®

CF

NASAL DECONGESTANT
COUGH SUPPRESSANT
EXPECTORANT

Why Are Robitussin Products Chosen More Often?



RECOMMENDED by more Doctors and Pharmacists than any other cough formula.



RELIABLE and EFFECTIVE formulas for the WHOLE FAMILY.

NON-DROWSY and NON-NARCOTIC formulas.

Richmond Division of Wyeth
2248-2300 Darbytown Rd.
Richmond, VA 23231

Keep out of reach of children. In case of overdose, get medical help right away.

Other information

- store at 20-25°C (68-77°F)
- يحزن في درجة حرارة العرفة بين ٢٠ إلى ٢٥ درجة مئوية
- dosage cup provided

الوكلاء الوحيدون شركة النهضة الطبية	info@annahdah.com
	السعر للجمهور ريال هلاله
Reg. No. 30/86/03	٨ ٨٠

PAA019124

ROBITUSSIN CF

dextromethorphan hydrobromide, guaifenesin, pseudoephedrine hydrochloride liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0031-8678
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 5 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0031-8678-82	1 in 1 CARTON	07/01/2015	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
Export only		07/01/2015	

Labeler - GlaxoSmithKline Consumer Healthcare Holdings (US) LLC (079944263)

