

**TRIAMCINOLONE ACETONIDE - triamcinolone acetonide ointment**  
**RPK Pharmaceuticals, Inc.**


-----  
**Triamcinolone Acetonide Ointment USP, 0.5%**

**For Dermatologic Use Only**  
**Not For Ophthalmic Use**

**Rx Only**

**DESCRIPTION**

The topical corticosteroids constitute a class of primarily synthetic steroids used as anti-inflammatory and anti-pruritic agents. Triamcinolone acetonide is designated chemically as 9-fluoro-11b, 16a,17,21-tetrahydroxypregna-1,4-diene-3,20-dione cyclic 16,17-acetal with acetone.  $C_{24}H_{31}FO_6$ , M.W. 434.51; CAS Reg. No. 76-25-5.



Each gram of Triamcinolone Acetonide Ointment USP, 0.5% contains 5 mg triamcinolone acetonide, respectively, in an ointment base of light mineral oil and white petrolatum.

**CLINICAL PHARMACOLOGY**

Topical corticosteroids share anti-inflammatory, anti-pruritic and vasoconstrictive actions. The mechanism of anti-inflammatory activity of the topical corticosteroids is unclear. Various laboratory methods, including vasoconstrictor assays, are used to compare and predict potencies and/or clinical efficacies of the topical corticosteroids. There is some evidence to suggest that a recognizable correlation exists between vasoconstrictor potency and therapeutic efficacy in man.

**Pharmacokinetics**

The extent of percutaneous absorption of topical corticosteroids is determined by many factors including the vehicle, the integrity of the epidermal barrier, and the use of occlusive dressings. Topical corticosteroids can be absorbed from normal intact skin. Inflammation and/or other disease processes in the skin increase percutaneous

absorption. Occlusive dressings substantially increase the percutaneous absorption of topical corticosteroids. Thus, occlusive dressings may be a valuable therapeutic adjunct for treatment of resistant dermatoses (see **DOSAGE AND ADMINISTRATION**). Once absorbed through the skin, topical corticosteroids are handled through pharmacokinetic pathways similar to systemically administered corticosteroids. Corticosteroids are bound to plasma proteins in varying degrees. Corticosteroids are metabolized primarily in the liver and are then excreted by the kidneys. Some of the topical corticosteroids and their metabolites are also excreted into the bile.

## **INDICATIONS AND USAGE**

Topical corticosteroids are indicated for the relief of the inflammatory and pruritic manifestations of corticosteroid-responsive dermatoses.

## **CONTRAINDICATIONS**

Topical corticosteroids are contraindicated in those patients with a history of hypersensitivity to any of the components of the preparations.

## **PRECAUTIONS**

### **General**

Systemic absorption of topical corticosteroids has produced reversible hypothalamic-pituitary-adrenal (HPA) axis suppression, manifestations of Cushing's syndrome, hyperglycemia, and glucosuria in some patients.

Conditions which augment systemic absorption include the application of the more potent steroids, use over large surface areas, prolonged use, and the addition of occlusive dressings.

Therefore, patients receiving a large dose of any potent topical steroid applied to a large surface area or under an occlusive dressing should be evaluated periodically for evidence of HPA axis suppression by using the urinary free cortisol and ACTH stimulation tests, and for impairment of thermal homeostasis. If HPA axis suppression or elevation of the body temperature occurs, an attempt should be made to withdraw the drug, to reduce the frequency of application, substitute a less potent steroid, or use a sequential approach when utilizing the occlusive technique.

Recovery of HPA axis function and thermal homeostasis are generally prompt and complete upon discontinuation of the drug. Infrequently, signs and symptoms of steroid withdrawal may occur, requiring supplemental systemic corticosteroids. Occasionally, a patient may develop a sensitivity reaction to a particular occlusive dressing material or adhesive and a substitute material may be necessary.

Children may absorb proportionally larger amounts of topical corticosteroids and thus be more susceptible to systemic toxicity (see **PRECAUTIONS-Pediatric Use**).

If irritation develops, topical corticosteroids should be discontinued and appropriate therapy instituted.

In the presence of dermatological infections, the use of an appropriate antifungal or antibacterial agent should be instituted. If a favorable response does not occur promptly, the corticosteroid should be discontinued until the infection has been adequately controlled.

These preparations are not for ophthalmic use.

### **Information for Patients**

Patients using topical corticosteroids should receive the following information and instructions:

1. This medication is to be used as directed by the physician. It is for external use only. Avoid contact with the eyes.
2. Patients should be advised not to use this medication for any disorder other than for which it was prescribed.
3. The treated skin area should not be bandaged or otherwise covered or wrapped as to be occlusive unless directed by the physician.
4. Patients should report any signs of local adverse reactions especially under occlusive dressing.
5. Parents of pediatric patients should be advised not to use tight-fitting diapers or plastic pants on a child being treated in the diaper area, as these garments may constitute occlusive dressings.

### **Laboratory Tests**

The following tests may be helpful in evaluating the HPA axis suppression: urinary free cortisol test and ACTH stimulation test.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Long-term animal studies have not been performed to evaluate the carcinogenic potential or the effect on fertility of topical corticosteroids. Studies to determine mutagenicity with prednisolone and hydrocortisone have revealed negative results.

### **Pregnancy:**

Teratogenic Effects:

#### *Pregnancy Category C*

Corticosteroids are generally teratogenic in laboratory animals when administered systemically at relatively low dosage levels. The more potent corticosteroids have been shown to be teratogenic after dermal application in laboratory animals. There are no adequate and well-controlled studies in pregnant women on the teratogenic effects from topically applied corticosteroids. Therefore, topical corticosteroids should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Drugs of this class should not be used extensively on pregnant patients, in large amounts, or for prolonged periods of time.

### **Nursing Mothers**

It is not known whether topical administration of corticosteroids could result in sufficient

systemic absorption to produce detectable quantities in breast milk. Systemically administered corticosteroids are secreted into breast milk in quantities not likely to have a deleterious effect on the infant. Nevertheless, caution should be exercised when topical corticosteroids are administered to a nursing woman.

## **Pediatric Use**

**Pediatric patients may demonstrate greater susceptibility to topical corticosteroid-induced HPA axis suppression and Cushing's syndrome than mature patients because of a larger skin surface area to body weight ratio.**

Hypothalamic-pituitary-adrenal (HPA) axis suppression, Cushing's syndrome and intracranial hypertension have been reported in children receiving topical corticosteroids. Manifestations of adrenal suppression in children include linear growth retardation, delayed weight gain, low plasma cortisol levels, and absence of response to ACTH stimulation. Manifestations of intracranial hypertension include bulging fontanelles, headaches, and bilateral papilledema.

Administration of topical corticosteroids to children should be limited to the least amount compatible with an effective therapeutic regimen. Chronic corticosteroid therapy may interfere with the growth and development of children.

## **ADVERSE REACTIONS**

The following local adverse reactions are reported infrequently with topical corticosteroids, but may occur more frequently with the use of occlusive dressings. These reactions are listed in an approximate decreasing order of occurrence: burning, itching, irritation, dryness, folliculitis, hypertrichosis, acneiform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis, maceration of the skin, secondary infection, skin atrophy, striae, and miliaria.

To report **SUSPECTED ADVERSE REACTIONS**, contact Teligent Pharma, Inc. at 1-856-697-1441, or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

## **OVERDOSAGE**

Topically applied corticosteroids can be absorbed in sufficient amounts to produce systemic effects (see **PRECAUTIONS**).

## **DOSAGE AND ADMINISTRATION**

Apply a thin film of triamcinolone acetonide ointment 0.5%, as appropriate, to the affected area two to three times daily.

## **OCCLUSIVE DRESSING TECHNIQUE**

Occlusive dressings may be used for the management of psoriasis or other recalcitrant conditions. Apply a thin film of ointment to the lesion, cover with a pliable nonporous film, and seal the edges. If needed, additional moisture may be provided by covering the lesion with a dampened clean cotton cloth before the nonporous film is applied or by briefly wetting the affected area with water immediately prior to applying the medication. The frequency of changing dressings is best determined on an individual basis. It may be

convenient to apply triamcinolone acetonide ointment under an occlusive dressing in the evening and to remove the dressing in the morning (i.e., 12-hour occlusion). When utilizing the 12-hour occlusion regimen, additional ointment should be applied, without occlusion, during the day. Reapplication is essential at each dressing change.

If an infection develops, the use of occlusive dressings should be discontinued and appropriate antimicrobial therapy instituted.

Occlusive dressing may be used for the management of psoriasis or recalcitrant conditions. If an infection develops, the use of occlusive dressing should be discontinued and appropriate antimicrobial therapy instituted.

## HOW SUPPLIED

Product: 53002-0151

NDC: 53002-0151-1 15 g in a TUBE

## STORAGE

Store at 20-25°C (68-77°F) [see USP Controlled Room Temperature].

Mfd. and Dist by:

Teligent Pharma, Inc.  
Buena, NJ 08310

Revised: 12/16

## Triamcinolone Acetonide 0.5% Ointment

<small>15 Gm Tube</small> <b>TRIAMCINOLONE ACETONIDE OINTMENT 0.5%</b> LOT# 25545-048-15 EXP: 05-31-2022 ORDER# 015-63 <b>Rx only</b> <small>Small dosage. See patient directions at back of box.</small>	<b>15 Gm Tube Rx# 210961161001 ORDER# 015-63</b> <b>TRIAMCINOLONE 0.5% OINTMENT</b> TELIGENT Generic for ARISTOCORT <b>APPLY TO AFFECTED AREA</b> <b>2-4 TIMES A DAY</b> <b>OR AS DIRECTED.</b>	 NDC 53002-0151-1 15 Gm TRIAMCINOLONE 0.5% OINTMENT EXP: 05-31-2022 LOT# 25545-048-15 Rx# 210961161001 ORDER# 015-63	<b>15 Gm TRIAMCINOLONE 0.5% OINTMENT</b> LOT# 25545-048-15 EXP: 05-31-2022 Rx# 210961161001 FCA-0101 15 Gm TRIAMCINOLONE 0.5% OINTMENT BILLING NDC# 53002-0151-1 Rx# 210961161001 15 Gm TRIAMCINOLONE 0.5% OINTMENT BILLING NDC# 53002-0151-1 Rx# 210961161001 15 Gm TRIAMCINOLONE 0.5% OINTMENT BILLING NDC# 53002-0151-1 Rx# 210961161001 15 Gm TRIAMCINOLONE 0.5% OINTMENT	<b>TRIAMCINOLONE ACETONIDE OINTMENT 0.5%</b> <b>15 Gm Tube</b> <b>DISCARD BY 05-31-2022</b> NDC# 53002-0151-1 Rx# 210961161001 <small>FEDERAL LAW PROHIBITS ANY PERSON FROM REPRODUCING OR TRANSMITTING ANY INFORMATION FROM THIS LABEL OR ANY INFORMATION FROM THE REACH OF CHILDREN. CALL YOUR DOCTOR FOR MEDICAL ADVICE ABOUT THIS PRODUCT. YOU MUST REPORT SIDE EFFECTS TO FDA AT 1-800-FDA-1088.</small> Clinic Name Here PRESCRIBER NAME DATE PATIENT NAME
<small>15 Gm Tube</small> <b>TRIAMCINOLONE ACETONIDE OINTMENT 0.5%</b> LOT# 25545-048-15 EXP: 05-31-2022 ORDER# 015-63 <b>Rx only</b> <small>Small dosage. See patient directions at back of box.</small>	<b>15 Gm Tube Rx# 210961161001 ORDER# 015-63</b> <b>TRIAMCINOLONE 0.5% OINTMENT</b> TELIGENT Generic for ARISTOCORT <b>APPLY TO AFFECTED AREA</b> <b>2-4 TIMES A DAY</b> <b>OR AS DIRECTED.</b>	 NDC 53002-0151-1 15 Gm TRIAMCINOLONE 0.5% OINTMENT EXP: 05-31-2022 LOT# 25545-048-15 Rx# 210961161001 ORDER# 015-63	<b>15 Gm TRIAMCINOLONE 0.5% OINTMENT</b> LOT# 25545-048-15 EXP: 05-31-2022 Rx# 210961161001 FCA-0101 15 Gm TRIAMCINOLONE 0.5% OINTMENT BILLING NDC# 53002-0151-1 Rx# 210961161001 15 Gm TRIAMCINOLONE 0.5% OINTMENT BILLING NDC# 53002-0151-1 Rx# 210961161001 15 Gm TRIAMCINOLONE 0.5% OINTMENT BILLING NDC# 53002-0151-1 Rx# 210961161001 15 Gm TRIAMCINOLONE 0.5% OINTMENT	<b>TRIAMCINOLONE ACETONIDE OINTMENT 0.5%</b> <b>15 Gm Tube</b> <b>DISCARD BY 05-31-2022</b> NDC# 53002-0151-1 Rx# 210961161001 <small>FEDERAL LAW PROHIBITS ANY PERSON FROM REPRODUCING OR TRANSMITTING ANY INFORMATION FROM THIS LABEL OR ANY INFORMATION FROM THE REACH OF CHILDREN. CALL YOUR DOCTOR FOR MEDICAL ADVICE ABOUT THIS PRODUCT. YOU MUST REPORT SIDE EFFECTS TO FDA AT 1-800-FDA-1088.</small> Clinic Name Here PRESCRIBER NAME DATE PATIENT NAME

## TRIAMCINOLONE ACETONIDE

triamcinolone acetonide ointment

### Product Information

HUMAN PRESCRIPTION

Item Code

NDC: 53002-0151/NDC: 53565

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:53002-0151(NDC:52565-048)	
<b>Route of Administration</b>	TOPICAL			
<b>Active Ingredient/Active Moiety</b>				
<b>Ingredient Name</b>		<b>Basis of Strength</b>	<b>Strength</b>	
<b>TRIAMCINOLONE ACETONIDE</b> (UNII: F446C597KA) (TRIAMCINOLONE ACETONIDE - UNII:F446C597KA)		TRIAMCINOLONE ACETONIDE	5 mg in 1 g	
<b>Inactive Ingredients</b>				
<b>Ingredient Name</b>			<b>Strength</b>	
<b>Light Mineral Oil</b> (UNII: N6K5787QVP)				
<b>Petrolatum</b> (UNII: 4T6H12BN9U)				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:53002-0151-1	15 g in 1 TUBE; Type 0: Not a Combination Product	01/01/2021	
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
ANDA	ANDA208590	03/03/2017		

**Labeler** - RPK Pharmaceuticals, Inc. (147096275)

## Establishment

Name	Address	ID/FEI	Business Operations
RPK Pharmaceuticals, Inc.		147096275	RELABEL(53002-0151) , REPACK(53002-0151)

Revised: 3/2022

RPK Pharmaceuticals, Inc.