

**IBUPROFEN AND PSEUDOEPHEDRINE HYDROCHLORIDE- ibuprofen and pseudoephedrine hydrochloride tablet, sugar coated**  
**Ohm Laboratories Inc.**

-----  
**Ibuprofen and Pseudoephedrine hydrochloride**

***Drug Facts***

<b><i>Active ingredients (in each caplet)</i></b>	<b><i>Purposes</i></b>
Ibuprofen, USP 200 mg (NSAID)*	Pain reliever/fever reducer
Pseudoephedrine HCl, USP 30 mg	Nasal decongestant

\* nonsteroidal anti-inflammatory drug

**Uses**

temporarily relieves these symptoms associated with the common cold or flu:

- headache
- fever
- sinus pressure
- nasal congestion
- minor body aches and pains

**Warnings**

**Allergy alert**

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning**

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- in children under 12 years of age
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other product that contains pseudoephedrine or any other nasal decongestant
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- fever gets worse or lasts more than 3 days
- nasal congestion lasts for more than 7 days
- symptoms continue or get worse
- redness or swelling is present in the painful area
- you get nervous, dizzy, or sleepless
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### **Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years of age and over:
  - take 1 caplet every 4 to 6 hours while symptoms persist. If symptoms do not respond to 1 caplet, 2 caplets may be used.
  - do not use more than 6 caplets in any 24-hour period unless directed by a doctor
- children under 12 years of age: do not use

### **Other information**

- store at 20 - 25° C (68 - 77° F). Avoid excessive heat above 40° C (104° F).
- read all warnings and directions before use. Keep carton.

### **Inactive ingredients**

acacia, calcium carbonate, carnauba wax, confectioner's sugar, corn starch, croscarmellose sodium, crospovidone, FD&C Blue no. 2 Aluminum Lake, FD&C Red no. 40 Aluminum Lake, FD&C Yellow no. 6 Aluminum Lake, gelatin, guar gum, hydrogenated vegetable oil, hydroxypropyl cellulose, iron oxide black, kaolin, polyethylene glycol, powdered cellulose, povidone, pregelatinized starch, propylene glycol, shellac, silicon dioxide, sodium benzoate, sucrose, talc, titanium dioxide, white wax

### **Questions?**

call **1-800-406-7984**

Distributed by:  
Ohm Laboratories Inc.  
New Brunswick, NJ 08901

### **PRINCIPAL DISPLAY PANEL - 200 mg/30 mg Tablet Blister Pack Carton**

†Compare To  
the active ingredients of  
Advil® Cold & Sinus

NDC 51660-493-41

See New Warnings Information

ohm®

Non-Drowsy  
Cold & Sinus  
Ibuprofen and Pseudoephedrine  
HCl Tablets, USP

Ibuprofen, USP 200 mg – Pain Reliever/Fever Reducer (NSAID)\*  
Pseudoephedrine HCl, USP 30 mg – Nasal Decongestant

\*nonsteroidal anti-inflammatory drug

- Relieves Sinus Pressure,  
Nasal Congestion and Fever

40 Coated Caplets\*\*  
(\*\*oval-shaped tablets)

5212203

**Drug Facts (continued)**

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use in children under 12 years of age.

If you have ever had an allergic reaction to any other pain reliever/fever reducer.

If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if:

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke

Ask a doctor or pharmacist before use if you are:

- under a doctor's care for any serious condition
- taking any other product that contains pseudoephedrine or any other nasal decongestant
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product:

- take with food or milk if stomach upset occurs

**Drug Facts (continued)**

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
- you have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- fever gets worse or lasts more than 3 days
- nasal congestion lasts for more than 7 days
- symptoms continue or get worse
- redness or swelling is present in the painful area
- you get nervous, dizzy, or sleepless
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years of age and over:
  - take 1 caplet every 4 to 6 hours while symptoms persist. If symptoms do not respond to 1 caplet, 2 caplets may be used.
- do not use more than 6 caplets in any 24-hour period unless directed by a doctor
- children under 12 years of age: do not use

5212203

**Drug Facts**

**Active ingredients (in each caplet) Purposes**

Ibuprofen, USP 200 mg (NSAID)\* Pain reliever/fever reducer

Pseudoephedrine HCl, USP 30 mg Nasal decongestant

\*nonsteroidal anti-inflammatory drug

**Uses** temporarily relieves these symptoms associated with the common cold or flu:

- headache
- fever
- sinus pressure
- minor body aches and pains

**Warnings**

Allergy alert: Ibuprofen may cause a severe allergic reaction, if:

- you have had stomach ulcers or bleeding problems
- you are age 60 or older
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product
- take other drugs containing prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

**Drug Facts (continued)**

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
- you have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- fever gets worse or lasts more than 3 days
- nasal congestion lasts for more than 7 days
- symptoms continue or get worse
- redness or swelling is present in the painful area
- you get nervous, dizzy, or sleepless
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years of age and over:
  - take 1 caplet every 4 to 6 hours while symptoms persist. If symptoms do not respond to 1 caplet, 2 caplets may be used.
- do not use more than 6 caplets in any 24-hour period unless directed by a doctor
- children under 12 years of age: do not use

NDC 51660-493-41

See New Warnings Information

Compare To  
the active ingredients of  
Advil® Cold & Sinus

**Non-Drowsy**

**ohm®**

**Cold & Sinus**

**Ibuprofen and Pseudoephedrine**

**HCl Tablets, USP**

Ibuprofen, USP 200 mg – Pain Reliever/Fever Reducer (NSAID)\*

Pseudoephedrine HCl, USP 30 mg – Nasal Decongestant

\*nonsteroidal anti-inflammatory drug

● Relieves Sinus Pressure,  
Nasal Congestion and Fever

**40 Coated Caplets\*\***

**oval-shaped tablets**

Expiration Date:

Batch No.

Non Varnish Area

**Drug Facts (continued)****Other information**

■ read all warnings and directions before use. Keep carton.

■ store at 20° - 25° C (68° - 77° F). Avoid excessive heat above 40° C (104° F).

**Inactive ingredients**

acacia, calcium carbonate, carnauba wax, confectioner's sugar, corn starch, croscarmellose sodium, crospovidone, FD&C Blue no. 2, Aluminum Lake, FD&C Red no. 40, Aluminum Lake, FD&C Yellow no. 6, Aluminum Lake, gelatin, guar gum, hydrophobated vegetable oil, hydroxypropyl cellulose, iron oxide black, kaolin, polyethylene glycol, powdered cellulose, povidone, pregelatinized starch, polyethylene glycol, shellac, silicon dioxide, sodium benzoate, sodium chloride, sucrose, talc, titanium dioxide, white wax


**Questions? Call 1-800-406-7994**

**⚠️ WARNING: DO NOT USE IF ALISTER UNITS ARE TORN, BROKEN OR SHOW ANY SIGNS OF TAMPERING.**

† All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Advil® Cold & Sinus.

R0920

Distributed by:  
Ohm Laboratories Inc.  
New Brunswick, NJ 08901



5 2 1 6 6 0 4 9 3 4 1 2

IBUPROFEN AND PSEUDOEPHEDRINE HYDROCHLORIDE

ibuprofen and pseudoephedrine hydrochloride tablet, sugar coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51660-493
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name		Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)		200 mg
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)		30 mg

Inactive Ingredients	
Ingredient Name	Strength
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
ACACIA (UNII: 5C5403N26O)	
CALCIUM CARBONATE (UNII: H0G9379FGK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
HYDROXYPROPYL CELLULOSE (1600000 WAMW) (UNII: RFW2ET671P)	
KAOLIN (UNII: 24H4NWX5CO)	
CROSPVIDONE (120 .MU.M) (UNII: 68401960MK)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCROSE (UNII: C151H8M554)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
WHITE WAX (UNII: 7G1J5DA97F)	
GUAR GUM (UNII: E89I1637KE)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
CORN OIL (UNII: 8470G57WFM)	

Product Characteristics			
Color	brown	Score	no score
Shape	OVAL (caplets)	Size	14mm

Flavor		Imprint Code		423
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51660-493-41	40 in 1 BLISTER PACK; Type 0: Not a Combination Product	10/13/2001	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA074567	10/13/2001	

**Labeler** - Ohm Laboratories Inc. (184769029)

Establishment			
Name	Address	ID/FEI	Business Operations
Ohm Laboratories Inc.		184769029	MANUFACTURE(51660-493)